SHORT COMMUNICATION

Transmembrane Respiration-driven H^+ Translocation is Unimpaired in an *eup* Mutant of *Escherichia coli*

By G. DUNCAN HITCHENS, DOUGLAS B. KELL* AND
J. GARETH MORRIS

Department of Botany and Microbiology, University College of Wales, Aberystwyth,
Dyfed SY23 3DA, U.K.

(Received 21 May 1982)

Respiration-driven H^+ translocation has been examined in an *eup* ('energy uncoupled phenotype') mutant of *Escherichia coli* and compared with that observed in its otherwise isogenic wild-type parent. Respiration-driven H^+ translocation was unimpaired in the *eup* mutant strain. It appears that the role of the *eup* gene product lies in the utilization of energized protons pumped across the *E. coli* cytoplasmic membrane.

INTRODUCTION

Certain mutant strains of *Escherichia coli* have been described that, although selected for resistance to the aminoglycoside neomycin, are also rather resistant to colicin K, grow poorly on succinate, and are impaired in the operation of a number of respiration-linked active transport systems (Plate, 1976). Respiratory and ATP hydrolase activities are not decreased in the mutant strains, designated *eup* (for 'energy uncoupled phenotype'), and it was proposed *inter alia* (Plate, 1976) that the *eup* gene codes for a proteinaceous component in the bacterial cytoplasmic membrane that acts to couple the free energy released by respiration to processes such as active transport. Later work (Plate, 1979; Plate & Suit, 1981) showed that the mutant allele maps at 86.5–87 min on the *E. coli* linkage map (Bachmann & Low, 1980), that the *eup* mutation only marginally affects the apparent membrane potential (as judged by the uptake of the methyltriphenylphosphonium cation) and the intracellular pH of these organisms, and that the lesion in some way prevents the mutant strains from coupling energized H^+ movements to solute symport. Physiological and genetic studies are consistent with the view that the *eup* gene is the same as the independently isolated genes *ssd* (Morris & Newman, 1980; Newman *et al.*, 1981, 1982) and *ecfB* (Thorbjarnardóttir *et al.*, 1978).

In view of current controversy concerning the reliability of the methods used for measuring the electrical and pH gradients across the bacterial cytoplasmic membrane (e.g. Kell, 1979; Tedeschi, 1981), we have measured respiration-driven H^+ translocation in an *eup* mutant strain and its otherwise isogenic *eup*⁺ parent. We report here that there is no significant difference in the ability of the two strains to effect respiration-driven H^+ translocation.

METHODS

Strains. *Escherichia coli* CJ49 (*lacZ*(Am) *trp*(Am) *eup*⁺*thi*) and CJ48 (*lacZ*(Am) *trp*(Am) *eup*-5 *thi*) were generously donated by Dr Charles A. Plate, Northwestern University Medical School, Chicago, U.S.A. and grown aerobically at 37 °C in Ozeki minimal medium supplemented with glucose (0.4%, w/v), thiamin (0.5 µg ml⁻¹) and tryptophan (50 µg ml⁻¹) (Plate & Suit, 1981). After each experiment, strains were checked for their appropriate phenotype by assay for aerobic growth on succinate and sensitivity/resistance to neomycin sulphate (10 µg ml⁻¹).

Respiration-driven H^+ translocation. This was measured by the O₂ pulse method of Scholes & Mitchell (1970). Mid-exponential cultures were harvested, washed three times in 150 mM-KCl containing 0.25 mM-glycylglycine,

Fig. 1. Respiration-driven H^+ translocation in *E. coli* CJ48 (*eup*) and CJ49 (*eup*⁺). Measurements were made as described in Methods. The number of cells present, in a final volume of 6 ml, was 1.34×10^{10} for the *eup* strain (●), and 1.74×10^{10} for the *eup*⁺ strain (○). Total cell counts were made microscopically, with an improved Neubauer counting chamber. The amount of oxygen added in each pulse was 23.5 ng-atom, and KSCN was present at the concentrations shown.

pH 6.5, and resuspended at 3 mg dry wt ml⁻¹ in the same solution to which carbonic anhydrase at 80 µg ml⁻¹ was added. KSCN was added to the concentrations shown in Fig. 1. Assays were carried out in a vigorously stirred glass reaction vessel maintained at 37 °C and equipped with a sensitive pH electrode connected to a potentiometric system (Kell & Morris, 1980; Hitchens & Kell, 1982). O₂-free N₂ was passed over the surface of the reaction mixture. O₂ pulses, as air-saturated KCl, and calibrating additions of anaerobic KOH and HCl were delivered in the usual way (Scholes & Mitchell, 1970).

Respiration. This was measured with a Clark-type oxygen electrode (Phillips & Kell, 1981).

RESULTS AND DISCUSSION

The apparent $\rightarrow H^+/O$ ratios shown by the *eup*⁺ and *eup* strains, as obtained by extrapolation of the O₂-induced pH excursions to the half time of O₂ reduction (Scholes & Mitchell, 1970), are shown in Fig. 1. The increase of the apparent $\rightarrow H^+/O$ ratio on the addition of the permeant SCN⁻ ion (Lawford & Haddock, 1973) demonstrates that the respiration-induced H^+ movements were transmembranous in nature, and not purely scalar (Scholes & Mitchell, 1970). Since the *eup* strain was unable to use succinate, and presumably other respiratory substrates, as carbon and energy source for growth, it was necessary to use glucose-grown cells for these experiments. This explains why the absolute limiting $\rightarrow H^+/O$ stoichiometries were lower than those obtained by other workers using *E. coli* cells grown on respiratory substrates and respiring endogenously (Lawford & Haddock, 1973; Gould & Cramer 1977; Jones, 1977). The crucial observation in the present work is that the ability of the *eup* strain to pump H^+ is unimpaired, in comparison with that of its similarly-grown *eup*⁺ parent.

We have recently reviewed the available evidence on the mode of action of membrane-active colicins, together with that concerning the nature of the *eup* gene product, and have proposed (Kell & Morris, 1981; Kell *et al.*, 1981) that the normal function of the *eup* gene product is to direct H^+ , energized (pumped) by respiration, laterally along the cytoplasmic membrane surfaces between their sources and their sinks such as the H^+ -ATPase and H^+ -solute symporters. The finding that the *eup* strain is unimpaired in its ability to pump H^+ across its cytoplasmic membrane is consistent with its ability to generate an apparent protonmotive force (Plate, 1979; Plate & Suit, 1981), and, when taken together with the knowledge that it is unable to utilize energized protons (Plate 1976, 1979; Plate & Suit, 1981), allows us to state quite explicitly that efficient protonmotivated energy coupling requires functional proteins additional to the generally recognized reversible proton pumps themselves, a conclusion in line with various earlier proposals (e.g. Plate, 1976, 1979; Plate & Suit, 1981; Hong, 1977; Kell & Morris, 1981; Kell *et al.*, 1981).

We are grateful to the Science and Engineering Research Council, U.K. for financial support. We are indebted to Dr Charles A. Plate for his generosity in making available to us the *eup*⁺ and *eup* strains with which this work was carried out.

REFERENCES

- BACHMANN, B. J. & LOW, K. B. (1980). Linkage map of *Escherichia coli* K-12. Edition 6. *Microbiological Reviews* **44**, 1–56.
- GOULD, J. M. & CRAMER, W. A. (1977). Relationship between oxygen-induced proton efflux and membrane energisation in cells of *Escherichia coli*. *Journal of Biological Chemistry* **252**, 5875–5882.
- HITCHENS, G. D. & KELL, D. B. (1982). Respiration-driven proton translocation in *Paracoccus denitrificans*: role of the 'permeant' ion. *Biochemical Society Transactions* **10**, 261.
- HONG, J.-S. (1977). An *ecf* mutation in *Escherichia coli* pleiotropically affecting energy coupling in active transport but not generation or maintenance of membrane potential. *Journal of Biological Chemistry* **252**, 8582–8588.
- JONES, C. W. (1977). Aerobic respiratory systems in bacteria. *Symposia of the Society for General Microbiology* **27**, 23–59.
- KELL, D. B. (1979). On the functional proton current pathway of electron transport phosphorylation: an electrodic view. *Biochimica et biophysica acta* **549**, 55–99.
- KELL, D. B. & MORRIS, J. G. (1980). Formulation and some biological uses of a buffer mixture whose buffering capacity is relatively independent of pH in the range pH 4–9. *Journal of Biochemical and Biophysical Methods* **3**, 143–150.
- KELL, D. B. & MORRIS, J. G. (1981). Proton-coupled membrane energy transduction: pathways, mechanisms and control. In *Vectorial Reactions in Electron and Ion Transport in Mitochondria and Bacteria*, pp. 339–347. Edited by F. Palmieri, E. Quagliariello, N. Siliprandi & E. C. Slater. Amsterdam: Elsevier/North Holland.
- KELL, D. B., CLARKE, D. J. & MORRIS, J. G. (1981). On proton-coupled information transfer along the surface of biological membranes and the mode of action of certain colicins. *FEMS Microbiology Letters* **11**, 1–11.
- LAWFORD, H. G. & HADDOCK, B. A. (1973). Respiration-driven proton translocation in *Escherichia coli*. *Biochemical Journal* **136**, 217–220.
- MORRIS, J. F. & NEWMAN, E. B. (1980). Map location of the *ssd* mutation in *Escherichia coli* K-12. *Journal of Bacteriology* **143**, 1504–1505.
- NEWMAN, E. B., MORRIS, J. F., WALKER, C. & KAPOOR, V. (1981). A mutation affecting L-serine and energy metabolism in *E. coli* K-12. *Molecular and General Genetics* **182**, 143–147.
- NEWMAN, E. B., MALIK, N. & WALKER, C. (1982). L-Serine degradation in *Escherichia coli* K-12: directly isolated *ssd* mutants and their intragenic revertants. *Journal of Bacteriology* **150**, 710–715.
- PHILLIPS, M. K. & KELL, D. B. (1981). A benzoxazole inhibitor of NADH dehydrogenase in *Paracoccus denitrificans*. *FEMS Microbiology Letters* **11**, 111–113.
- PLATE, C. A. (1976). Mutant of *Escherichia coli* defective in response to colicin K and in active transport. *Journal of Bacteriology* **125**, 467–474.
- PLATE, C. A. (1979). Colicin K as a probe of energy coupling in *Escherichia coli*. In *Microbiology – 1979*, pp. 58–61. Edited by D. Schlessinger. Washington, D.C.: American Society for Microbiology.
- PLATE, C. A. & SUIT, J. L. (1981). The *eup* genetic locus of *Escherichia coli* and its role in H⁺ solute symport. *Journal of Biological Chemistry* **256**, 12974–12980.
- SCHOLES, P. & MITCHELL, P. (1970). Respiration-driven proton translocation in *Micrococcus denitrificans*. *Journal of Bioenergetics* **1**, 309–323.
- TEDESCHI, H. (1981). The transport of cations in mitochondria. *Biochimica et biophysica acta* **639**, 157–196.
- THORBJARNARDÓTTIR, S. H., MAGNÚSDÓTTIR, R. Á., EGGERTSON, G., KAGAN, S. A. & ANDRÉSSON, O. S. (1978). Mutations determining generalised resistance to aminoglycoside antibiotics in *Escherichia coli*. *Molecular and General Genetics* **161**, 89–98.