Phylogenetic analysis of some *Aerococcus*-like organisms from urinary tract infections: description of *Aerococcus urinae* sp. nov.

MARIA AGUIRRE and MATTHEW D. COLLINS*

Department of Microbiology, AFRC Institute of Food Research, Reading Laboratory, Shinfield, Reading RG2 9AT, UK

(Received 15 August 1991; revised 26 September 1991; accepted 21 October 1991)

Partial 16S ribosomal ribonucleic acid sequences of five *Aerococcus*-like organisms originally isolated from patients with urinary tract infections were determined using reverse transcriptase in order to clarify their taxonomic position. Analysis of the sequence data revealed that the clinical isolates represent a hitherto unknown line of descent within the genus *Aerococcus*. A new species, *Aerococcus urinae*, is proposed for these isolates. The type strain is NCFB 2893.

Introduction

The genus *Aerococcus* was proposed by Williams *et al.* (1953) to accommodate some Gram-positive, microaerophilic, catalase-negative bacteria which differed from streptococci primarily by their characteristic tetrad cellular arrangements. Currently, the genus comprises a single species, *Aerococcus viridans* (Evans, 1986). Aerococci are widely distributed and occur naturally in air, dust and vegetation; they also form part of the flora of meat-curing brines (Deibel & Niven, 1960). Although aerococci are generally considered saprophytic, some are pathogenic for lobsters (Hitchner & Snieszko, 1947) and there have been various reports of *Aerococcus*-like organisms from human infections (e.g. subacute bacterial endocarditis: Colman, 1967; Janosek *et al.*, 1980; Parker & Ball, 1976; urinary tract infections: Colman, 1967; bacteraemia: Taylor & Trueblood, 1985; and meningitis: Nathavitharana *et al.*, 1983). The identification of aerococci is complex, due to the paucity of reliable discriminatory phenotypic traits. Their most characteristic feature is their tendency to divide in two planes at right angles to form tetrads. Aerocci also exhibit many biochemical and physiological similarities with pediococci, enterococci, lactococci and leuconostocs, and are often confused with streptococci (Facklam *et al.*, 1989). Problems with the identification of aerococci are further

compounded by evidence of some genetic heterogeneity within the species (Bosley *et al.*, 1990; Wiik *et al.*, 1986).

Christensen *et al.* (1989, 1991) recently reported the isolation and characterization of some *Aerococcus*-like organisms (ALOs) from urine samples of patients with urinary tract infections. These clinical ALOs differed from typical *Aerococcus viridans* strains in several phenotypic tests and their precise taxonomic status remained uncertain. 16S rRNA sequencing studies have done much to clarify the inter- and intra-generic relationships of aerococci, pediococci, streptococci and other lactic acid bacteria (Bentley *et al.*, 1991; Collins *et al.*, 1989, 1990, 1991; Martinez Murcia & Collins, 1990; Williams *et al.*, 1991; Yang & Woese, 1989). Therefore, in the present study we have investigated the partial 16S rRNA sequences of five ALOs from urinary tract infections (Christensen *et al.*, 1989, 1991) in an attempt to determine their true phylogenetic position.

Methods

Cultures and cultivation. Five strains, designated E2, E3, B3, I3 and ES4, originally isolated from urine specimens, were received from Dr J. J. Christensen (Bispebjerg Hospital, Copenhagen, Denmark). The isolates were grown on blood agar plates and, for nucleic acid analysis, in glucose/yeast extract/phosphate broth (Garvie, 1978) at 37 °C. Cultures were harvested by centrifugation in late exponential phase and washed in deionized water.

Analysis of 16S rRNA. Cells (about 2 g) were mechanically broken using a Braun homogenizer, and RNA purified according to the procedure of Embley *et al.* (1988). Nucleotide sequences were determined by the dideoxynucleotide chain-termination method (Sanger *et al.*, 1977) directly from rRNA using AMV reverse transcriptase (Lane *et al.*, 1985). Nucleotide fragments produced in the

* Author for correspondence. Tel. (0734) 883103; fax (0734) 884763.

Abbreviation: ALO, *Aerococcus*-like organism.

The nucleotide sequence data reported in this paper have been submitted to GenBank and have been assigned the accession number M77819.

chain-elongation reactions were separated on 55 cm wedge-shaped 6% (w/v) acrylamide/7 M urea gels at 55 °C using an LKB MacroPhor 2010 sequencing unit operated at 50 W per gel.

The sequences generated were aligned, and homology values determined by the Beckman Microgenie Program (Queen & Korn, 1984). Nucleotide substitution rates (K_{nuc} values) were calculated and an unrooted phylogenetic tree produced using the algorithm of Fitch & Margoliash (1967).

Biochemical tests. Biochemical tests were performed on the five clinical strains and the type strain of *Aerococcus viridans* (NCDO 1225) by using the API 20 STREP and 50 CH systems according to the manufacturer's instructions. Blood agar was used as basal medium. Test preparations were incubated at 37 °C and read after 4, 24 and 48 h.

Results and Discussion

The partial 16S rRNA nucleotide sequence from ALO strain E2 was determined using reverse transcriptase. The sequence is shown in Fig. 1; it consisted of a continuous stretch of 1481 nucleotides representing about 96% of the total molecule. The first and last nucleotides in the sequence correspond to positions 8 and 1482 of the *Escherichia coli* numbering system (Brosius *et al.*, 1978). The primary structures of two short fragments (approx. positions 50 to 450, which includes variable regions V1 and V2; and positions 950 to 1210, which includes variable region V6; see Neefs *et al.*, 1990, for nomenclature) of the 16S rRNA of four other ALOs (strains E3, B3, I3 and ES4) were also determined and found to be identical to that of strain E2, thereby confirming the genetic relatedness of the five clinical isolates.

In order to determine the taxonomic position of the clinical ALOs, the sequence of strain E2 was aligned and compared with those of *Aerococcus viridans* and 27 reference strains representing 14 low-mol% G+C genera. Percentage sequence similarities, shown in Table 1, were calculated from a continuous region of 1340 nucleotides (ranging from positions 107 to 1433 of the *E.*

coli sequence). Approximately 100 nucleotides at the 5' end of the rRNA molecule were not included in the calculation to eliminate possible alignment errors due to the extremely variable V1 region (Neefs *et al.* 1990) and to ensure that only homologous positions were compared. The unknown isolate exhibited the highest similarity (94.5%) with *A. viridans*. High sequence similarities (approx. 91 to 92%) were also shown with carnobacteria, enterococci and vagococci, but significantly lower values (approx. 88 to 90%) were shared with lactobacilli, pediococci and streptococci. Some representative sequence homologies are shown in Table 1. A phylogenetic tree constructed from a matrix of derived K_{nuc} values using the Fitch and Margolish algorithm is shown in Fig. 2. In this scheme, the unknown clinical bacterium showed a close affinity with *A. viridans*, and clearly represents a second line within the genus *Aerococcus*. The five ALOs possessed identical biochemical profiles using the API 20 STREP and 50 CH systems. The type strain of *A. viridans* differed from these strains in several tests. Differential tests are shown in Table 2.

Phylogenetic studies have shown that the genus *Aerococcus* represents a distinct line of descent within the lactic acid bacteria (Collins *et al.*, 1990, 1991). Currently the genus consists of a single species, *A. viridans* (Evans, 1986). There is, however, indication of some heterogeneity within aerococci. In an investigation of aerococci from clinical sources Bosley *et al.* (1990) reported that two groups were discernible within *A. viridans* on the basis of DNA-DNA pairing. The two groups, however, possessed very high overall chromosomal DNA sequence relatedness (60 to 78% homology under optimal hybridization conditions) and were phenotypically indistinguishable. 16S rRNA sequencing studies confirmed the close affinity between these two groups, with only two base differences detected for a comparison of 1486 nucleotides between *A. viridans* and a representative strain of the second DNA homology group (Collins *et al.*,

```

8   AGAGUUUGAUCCUGGCUAGGACGAACGCGUGGCGGCGGCCUAAUACAUGCAAGUCGAGCGAACCGACGAAGUCUUGCAGUACGUUAGCGGCGGA
110 CGGGUGAGUAAACACGUAAAGGAACCUACCGAUAAAGCGGGGACAAACUCCGAAACGGGUGCUAAUACCGCAUAGGAAAGGUCACCNCAUGGUGACCUUUG
202 GAAAGACGGCUUUGUGUCACUUAUCGAUGGCCUUGCNGUGCAUUAAGCUCGUUGGUGGGUAAACGGCCUACCAAGGCAUAGAUGCAUAGCCGACCGAGAGA
301 GGGUAAUCGGCCACAUUGGGACUGAGACACGGCCCAACUCCUACGGGAGGCAGCAGUAGGGAAUCUUCGCAUUGGGCGCAAGCCUGACGGAGCAACGC
401 CGCGUGAGUGAAGAAGGUUUUCGGAUCGUAAAGCUCUGUUGUAGAGAAGAAACAAUUGGAGAGUAAACUGCUCUUGACGGUUAUCUUAACAGAAAG
501 CCACGGCUAAACUAGUGCCAGCAGCCGCGUAAUACGUAGGUGGCAAGCGUUGUCCGGAUUUUAUUGGGCGUAAAGGGGCGCAGGCGGUUUCUUAAGUCU
601 GAUGUGAAAGCCACGGCUUUAACCGUGGAAGUGCAUUGGAAACUGGGNAACUUGAGUACAGAGAGAGAAAGUGGAAACUCCAUGUGUAGCGGUGGAAUGCG
701 UAGAUAUUAUGGAAGAACACAGUGGGCGAAGGCGACUUCUGGUCUGUCACUGACGCGUGAGGCGCGAAAGCGUGGGUAGCAAACAGGAUUAAGAUACCCUGG
801 UAGUCCACGGCGUAAACGAUGAGUGCUANGUGUUGGAGGGUUUCCACCCUUCAGUGCCGGAGUUAACGCAUUAAGCACUCCGCCUGGGGAGUACGGCCGC
900 AAGGCUAGAAACUAAAGGAUUGACGGGGACCGCACAAAGCGGUGGAGCAUGUGGUUUAUUCGAAGCAACGCGAAGAACCUAACCAAGUCUUGACAUCCG
1001 UUGACCACUCUAGAGAUAGAGCUUUCUUGGGGACNAAGUGACAGGUGGUGCAUGGUUGUCUGCAGCUCGUGUGAGAUUUGGGUUAAGUCCCGCAA
1103 CGAGCGCAACCCUUAUUGUUAUGUUGCCAGCAUUNAGUUGGGCACUUCUAGCGAGACUGCCGGUGACAAACCGGAGGAAGGCGGGGAGUACGUAUUAUC
1204 AUGCCCCUUAUGACUUGGGCUACACAGUGCUACAAUGGAUGGUUACAAAGAGUCAGCGACCUUGUGAAAGCAAGCGAAUCUUAAGGCAUUCUCAGUU
1303 CGGAUUGUAGUCUGCAACUCGACUACAUGAAGCGGAAUCGCUAGUAAUCGCGGAGCAGCAGCCCGGUGAAUACGUUCCCGGUCUUGUACACACCGC
1403 CCGUCACACCACGAGAGUUUGUAAACCCUGAAGUGCGGUGAGGUAAACCUUUGGAGCCAGCCGCGAAGGUNNACAGAUUU

```

Fig. 1. Primary structure of 16S rRNA from *Aerococcus urinae* (NCFB 2893) determined using reverse transcriptase. N, undetermined nucleotide. The number of the nucleotide at the beginning of each line in the sequence corresponds to the *E. coli* numbering system (Brosius *et al.*, 1978).

Table 1. Homology (lower left-hand triangle) and K_{nuc} values (upper right-hand triangle) for a 1340-nucleotide region of 16S rRNAs of *Aerococcus urinae* and other taxa

	1	2	3	4	5	6	7	8	9	10	11	12	13	14
1. <i>Aerococcus urinae</i> NCFB 2893 ^T (= ALO E2)	·	0.0571	0.1225	0.1284	0.0812	0.0891	0.1260	0.1513	0.1428	0.1120	0.1085	0.1143	0.1260	0.0925
2. <i>Aerococcus viridans</i> NCDO 1225 ^T	94.5	·	0.1355	0.1190	0.0790	0.0812	0.1476	0.1260	0.1367	0.1050	0.1143	0.1004	0.1225	0.0768
3. <i>Bacillus subtilis</i> rrNB	88.7	87.6	·	0.1143	0.1062	0.1004	0.1274	0.1537	0.1636	0.0868	0.1225	0.0891	0.1476	0.1131
4. <i>Brochothrix thermosphacta</i> NCDO 1676 ^T	88.2	89.0	89.4	·	0.1050	0.0925	0.1525	0.1464	0.1649	0.0724	0.1213	0.1096	0.1331	0.0936
5. <i>Carnobacterium divergens</i> NCDO 2763 ^T	92.3	92.5	90.1	90.2	·	0.0560	0.1331	0.1343	0.1476	0.0846	0.0835	0.1096	0.1154	0.0658
6. <i>Enterococcus cecorum</i> NCDQ 2674 ^T	91.6	92.3	90.6	91.3	94.6	·	0.1154	0.1004	0.1331	0.0824	0.0835	0.1050	0.1027	0.0593
7. <i>Lactobacillus acidophilus</i> NCDO 1748 ^T	88.4	86.6	88.3	86.2	87.8	89.3	·	0.1661	0.1296	0.1428	0.1131	0.1574	0.1416	0.1272
8. <i>Lactococcus garvieae</i> NCDO 2155 ^T	86.3	88.4	86.1	86.7	87.7	90.6	85.1	·	0.1599	0.1440	0.1201	0.1699	0.0902	0.1331
9. <i>Leuconostoc mesenteroides</i> NCDO 523 ^T	87.0	87.5	85.3	85.2	86.6	87.8	88.1	85.6	·	0.1537	0.1476	0.1686	0.1488	0.1428
10. <i>Listeria innocua</i> NCTC 11288 ^T	89.6	90.2	91.8	93.1	92.0	92.2	87.0	86.9	86.1	·	0.1108	0.1027	0.1320	0.0947
11. <i>Pediococcus acidilactici</i> NCFB 2767 ^T	89.9	89.4	88.7	88.8	92.1	92.1	89.5	88.9	86.6	89.7	·	0.1272	0.1225	0.0925
12. <i>Staphylococcus xylosus</i> NCTC 11043 ^T	89.4	90.6	91.6	89.8	89.8	90.2	85.8	84.8	84.9	90.4	88.3	·	0.1550	0.1120
13. <i>Streptococcus pyogenes</i> NCDO 2381 ^T	88.4	88.7	86.6	87.8	89.3	90.4	87.1	91.5	86.5	87.9	88.7	86.0	·	0.1260
14. <i>Vagococcus fluvialis</i> NCDO 2497 ^T	91.3	92.7	89.5	91.2	93.7	94.3	88.3	87.8	87.0	91.1	91.3	89.6	89.0	·

Table 2. Characteristics useful in differentiating *A. urinae* and *A. viridans*

Data were obtained using the API 20STREP and 50CH systems; results are for five strains of *A. urinae* and the type strain of *A. viridans*.

Test	<i>A. urinae</i>	<i>A. viridans</i>
Production of:		
β -Galactosidase	—	+
β -Glucuronidase	+	—
Pyrrolidonylarylamidase	—	+
Leucine aminopeptidase	+	—
Acid production from:		
Lactose	—	+
Glycerol	—	+
Sorbitol	+	—

1990). The results of the present study clearly demonstrate that the human ALOs isolated by Christensen *et al.* (1989, 1991) from urinary tract infections are distinct from the *A. viridans* and genotype 2 of Bosley *et al.* (1990). The 16S rRNA of ALO strain E2 showed 74 base differences with that of the type strain of *A. viridans* (based on a comparison of the same 1486 nucleotide stretch). The ALOs from urine are also phenotypically quite distinct from *A. viridans* (see Table 2). Therefore, on the basis of the present findings it is formally

proposed that the ALOs of Christensen and associates be classified as a new species of the genus *Aerococcus*, as *Aerococcus urinae* (u.ri'nae: L. fem. n. *urinae*, pertaining to urine).

Description of *Aerococcus urinae* sp. nov.

Cells are Gram-positive, coccoid, occurring mostly in clusters but also in pairs and tetrads. Non-pigmented and non-motile. Microaerophilic. Catalase-negative. Growth does not occur at 10 and 45 °C. Growth occurs in 6.5% (w/v) NaCl, and on 5% (v/v) horse blood agar, producing α -haemolytic reaction. Acid is produced from D-arabitol (slow reaction), D-glucose, D-fructose, mannitol, ribose (slow reaction), sorbitol, sucrose and xylitol. Acid is not produced from amygdalin, arabinose, galactose, glycerol, glycogen, inositol, lactose, maltose, D-raffinose and trehalose. Hippurate is hydrolysed. Aesculin is not hydrolysed. β -Glucuronidase and leucine aminopeptidase positive. α -Galactosidase, β -galactosidase, pyrrolidonylarylamidase, alkaline phosphatase and arginine dihydrolase negative. H₂S is not produced. Nitrate is not reduced to nitrite. The G+C content of DNA is 44.4 mol%, as determined by melting temperature. Isolated from human sources (urinary tract specimens). The type strain is NCFB 2893 (NCTC 12142).

Fig. 2. Unrooted phylogenetic tree showing the position of *Aerococcus urinae* within a radiation of lactic acid bacteria. Genus abbreviations: A., *Aerococcus*; B., *Bacillus*; Br., *Brochothrix*; C., *Carnobacterium*; E., *Enterococcus*; Lb., *Lactobacillus*; Lc., *Lactococcus*; Leu., *Leuconostoc*; List., *Listeria*; P., *Pediococcus*; S., *Streptococcus*; Staph., *Staphylococcus*; T., *Tetragenococcus*; V., *Vagococcus*. Reference species corresponded to *C. piscicola* NCDO 2762^T, *E. hirae* NCDO 1258^T, *Lb. brevis* NCDO 1749^T, *Lb. casei* NCDO 161^T, *Lb. delbrueckii* NCDO 213^T, *Lb. plantarum* NCDO 1752^T, *Lb. reuteri* NCDO 2589^T, *Lc. lactis* NCDO 2118^T, *Leu. paramesenteroides* NCDO 803^T, *P. damnosus* NCDO 1832^T, *S. sanguis* NCTC 7863^T, *S. parauberis* NCDO 651, *S. porcinus* NCDO 600, *T. halophilus* NCDO 1635^T, and *V. salmoninarum* NCFB 2777^T. Strain details of other species examined are given in Table 1.

We are grateful to Dr J. J. Christensen (Copenhagen) for the gift of cultures.

References

- BENTLEY, R. W., LEIGH, J. A. & COLLINS, M. D. (1991). Intrageneric structure of *Streptococcus* based on comparative analysis of small-subunit rRNA sequences. *International Journal of Systematic Bacteriology* **41**, 487–494.
- BOSLEY, G. S., WALLACE, P. L., MOSS, C. W., STEIGERWALT, A. G., BRENNER, D. J., SWENSON, J. M., HEBERT, G. A. & FACKLAM, R. R. (1990). Phenotypic characterization, cellular fatty acid composition, and DNA relatedness of aerococci and comparison to related genera. *Journal of Clinical Microbiology* **28**, 416–421.
- BROSIOUS, J., PALMER, M. L., KENNEDY, J. P. & NOLLER, H. F. (1978). Complete nucleotide sequence of a 16S ribosomal RNA gene from *Escherichia coli*. *Proceedings of the National Academy of Sciences of the United States of America* **75**, 4801–4805.
- CHRISTENSEN, J. J., KORNER, B. & KJAEGAARD, H. (1989). *Aerococcus*-like organism – an unnoticed urinary tract pathogen. *APMIS* **97**, 539–546.
- CHRISTENSEN, J. J., VIBITS, H., URSING, F. & KORNER, B. (1991). *Aerococcus*-like organism, a newly recognized potential urinary tract pathogen. *Journal of Clinical Microbiology* **29**, 1049–1053.
- COLLINS, M. D., ASH, C., FARROW, J. A. E., WALLBANKS, S. & WILLIAMS, A. M. (1989). 16S ribosomal ribonucleic acid sequences analyses of lactococci and related taxa. Description of *Vagococcus fluvialis* gen. nov., sp. nov. *Journal of Applied Bacteriology* **67**, 453–460.
- COLLINS, M. D., WILLIAMS, A. M. & WALLBANKS, S. (1990). The phylogeny of *Aerococcus* and *Pediococcus* as determined by 16S rRNA sequence analysis: description of *Tetragenococcus* gen. nov. *FEMS Microbiology Letters* **70**, 255–262.
- COLLINS, M. D., RODRIGUES, U., ASH, C., AGUIRRE, M., FARROW, J. A. E., MARTINEZ MURCIA, A., PHILLIPS, B. A., WILLIAMS, A. M. & WALLBANKS, S. (1991). Phylogenetic analysis of the genus *Lactobacillus* and related lactic acid bacteria as determined by reverse transcriptase sequencing of 16S rRNA. *FEMS Microbiology Letters* **77**, 5–12.
- COLMAN, G. (1967). *Aerococcus*-like organisms isolated from human infections. *Journal of Clinical Pathology* **20**, 294–297.
- DEIBEL, R. H. & NIVEN, C. F. (1960). Comparative study of *Gaffkya homari*, *Aerococcus viridans*, tetrad-forming cocci from meat curing brines, and the genus *Pediococcus*. *Journal of Bacteriology* **79**, 175–180.
- EMBLEY, T. M., SMIDA, J. & STACKEBRANDT, E. (1988). Reverse transcriptase sequencing of 16S ribosomal RNA from *Faenia rectivirgula*, *Pseudonocardia thermophila* and *Saccharopolyspora hirsuta*, three wall IV actinomycetes which lack mycolic acids. *Journal of General Microbiology* **134**, 961–966.

- EVANS, J. B. (1986). Genus *Aerococcus*. In *Bergey's Manual of Systematic Bacteriology*, vol. 2, p. 1080. Edited by P. H. A. Sneath, N. S. Mair, M. E. Sharpe & J. G. Holt. Baltimore: Williams & Wilkins.
- FACKLAM, R. R., HOLLIS, D. & COLLINS, M. D. (1989). Identification of gram-positive coccid and coccobacillary vancomycin-resistant bacteria. *Journal of Clinical Microbiology* **27**, 724–730.
- FITCH, W. M. & MARGOLISH, E. (1967). Construction of phylogenetic trees: a method based on mutation distances as estimated from cytochrome c sequences is a general applicability. *Science* **155**, 279–284.
- GARVIE, E. I. (1978). *Streptococcus raffinolactis* (Orla-Jensen and Hansen): a group N streptococcus found in raw milk. *International Journal of Systematic Bacteriology* **28**, 190–193.
- HITCHNER, E. R. & SNIESZKO, S. F. (1947). A study of a microorganism causing a bacterial disease of lobsters. *Journal of Bacteriology* **54**, 48.
- JANOSEK, J., ECKERT, J. & HUDAC, A. (1980). *Aerococcus viridans* as a causative agent of infectious endocarditis. *Journal of Clinical Pathology* **20**, 92–96.
- LANE, D. J., PACE, B., OLSEN, G. J., STAHL, D. A., SOGIN, M. L. & PACE, N. R. (1985). Rapid determination of 16S ribosomal RNA sequences for phylogenetic analyses. *Proceedings of the National Academy of Sciences of the United States of America* **82**, 6955–6959.
- MARTINEZ MURCIA, A. & COLLINS, M. D. (1990). A phylogenetic analysis of the genus *Leuconostoc* based on reverse transcriptase sequencing of 16S rRNA. *FEMS Microbiology Letters* **70**, 73–84.
- NATHAVITHARANA, K. A., ARSECULERATNE, S. N., APOUNSO, H. A., VIJERATNAM, R., JAYASENA, L. & NAVARATMAN, C. (1983). Acute meningitis in early childhood caused by *Aerococcus viridans*. *British Medical Journal* **286**, 1248.
- NEEFS, J. M., VAN DE PEER, Y., HENDRIKS, L. & DE WACHTER, R. (1990). Compilation of small ribosomal subunit RNA sequences. *Nucleic Acids Research Supplement* **18**, 2237–2242.
- PARKER, M. T. & BALL, L. C. (1976). Streptococci and aerococci associated with systemic infection in man. *Journal of Medical Microbiology* **9**, 275–302.
- QUEEN, C. & KORN, L. J. (1984). A comprehensive sequence analysis program for the IBM personal computer. *Nucleic Acids Research* **12**, 581–599.
- SANGER, F., NICKLEN, S. & COULSON, A. R. (1977). DNA sequencing with chain-terminating inhibitors. *Proceedings of the National Academy of Sciences of the United States of America* **74**, 5463–5467.
- TAYLOR, P. V. & TRUEBLOOD, M. C. (1985). Septic arthritis due to *Aerococcus viridans*. *Journal of Rheumatology* **12**, 1604–1605.
- WIJK, R., TORSVIK, V. & EGIDIUS, E. (1986). Phenotypic and genotypic comparisons among strains of the lobster pathogen *Aerococcus viridans* and other marine *Aerococcus viridans*-like cocci. *International Journal of Systematic Bacteriology* **36**, 431–434.
- WILLIAMS, A. M., RODRIGUES, U. M. & COLLINS, M. D. (1991). Intrageneric relationships of enterococci as determined by reverse transcriptase sequencing of small subunit rRNA. *Research in Microbiology* **142**, 67–74.
- WILLIAMS, R. E. O., HIRSCH, A. & COWAN, S. T. (1953). *Aerococcus*, a new bacterial genus. *Journal of General Microbiology* **8**, 475–480.
- YANG, D. & WOESE, C. R. (1989). Phylogenetic structure of the leuconostocs: an interesting case of a rapidly evolving organism. *Systematic and Applied Microbiology* **12**, 145–149.