Alan E. Wheals,*¹ Maxine E. Smith¹ and J. Richard Dickinson²

¹School of Biology and Biochemistry, University of Bath, Claverton Down, Bath, BA2 7AY, UK.

²Department of Pure and Applied Biology, University of Wales College of Cardiff, PO Box 915, Cardiff, CF1 3TL, UK.

*For correspondence. Tel: +44 1225 826826 ext. 4278. Fax: +44 1225 826779. e-mail: A.E.Wheals@bath.ac.uk

1. Baroni, M.D., Monti, P. & Alberghina, L. (1994). Repression of growth-regulated G1 cyclin expression by cyclic AMP in budding yeast. *Nature* **371**, 339–342.
2. Brown, B.L., Albano, J.D.M., Ekins, R.P. & Sgherzi, A.M. (1971). A simple and sensitive saturation assay method for the measurement of adenosine 3', 5'-cyclic monophosphate. *Biochem J* **121**, 561–562.
3. François, J., Eraso, P. & Gancedo, C. (1987). Changes in the concentration of cAMP, fructose 2,6-bisphosphate and related metabolites and enzymes in *Saccharomyces cerevisiae* during growth on glucose. *Eur J Biochem* **164**, 369–373.
4. Sherman, F. (1991). Getting started with yeast. In *Guide to Yeast Genetics and Molecular Biology*, pp. 3–21. Edited by C. Guthrie & G. Fink. London: Academic Press.
5. Smith, M., Dickinson, J.R. & Wheals, A.E. (1990). Intracellular and extracellular levels of cyclic AMP during the cell cycle of *Saccharomyces cerevisiae*. *Yeast* **6**, 53–60.
6. Tokiwa, G., Tyers, M., Volpe, T. & Fletcher, B. (1994). Inhibition of G1 cyclin activity by the RAS/cAMP pathway in yeast. *Nature* **371**, 342–345.
7. Uno, I. (1992). Role of signal transduction systems in cell proliferation in yeast. *Int Rev Cytol* **139**, 309–332.

Paper Commentary

Oxidase names: to '3' or not to '3'?

In an earlier issue of *Microbiology*, we presented data on the spectral kinetic properties of a terminal quinol oxidase of *Escherichia coli* (7). This oxidase in particular has become the subject of intense interest, especially from bioenergeticists, biochemists and spectroscopists who recognize the attractions of a bacterial oxidase that serves as an amenable model for the cytochrome *c* oxidase (EC 1.9.3.1) of eukaryotes, an enzyme of enormous complexity and difficulty. In that paper, we suggested that due care be given to the nomenclature of this and other bacterial oxidases. The father of cytochrome studies, David Keilin, certainly held this view (3), writing: 'As the position of absorption bands of cytochrome observed in intact cells of micro-organisms show certain variations, it is important when referring to these bands to use as uniform a terminology as possible'. However, there are now appearing in the burgeoning literature (10) names that are not only unrecognized in current official nomenclature recommendations but which, in our view, are etymologically inaccurate. The purpose of this commentary on our own paper is to expand the argument advanced there, point out the pitfalls of extending the 'new', unofficial nomenclature to other cytochromes (which seems to be happening) and to draw the attention of the community studying these enzymes to the proper historical background.

The discovery of 'cytochrome o'

In 1954, Lucile Smith, LaRoy Castor and one of us (B.C.) discovered a photodissociable carbon monoxide compound in several types of bacteria, the absorption bands of which were similar to haemoglobin. In view of this similarity, and recognizing the claim by Keilin & Tissièrè of the presence of haemoglobin in certain micro-organisms, functional studies were made. Using an improvement of Warburg's photochemical method, Castor & Chance (2) showed the mysterious haemoprotein to be an oxidase and named it cytochrome *o*. It is important to recognize that the name, *o*, signifies oxidase. In an otherwise very clear summary of haem nomenclature (1), an explanation of 'cytochrome *o*' illustrates the present misunderstanding of the name's origin: 'The original designation of an *o*-type cytochrome meant that it contained a heme B, which can bind to carbon monoxide (or O₂, extrapolating for oxidases)'. Firstly, the name was intended to describe function (*o* for oxidase), not haem type. At the time, haem O (see below) had not been discovered but the spectral similarity with haem B (5) justified the supposition that the ligand-reactive haem was B. Secondly, this description implies that the characteristic of cytochrome *o* is CO binding, whereas Chance & co-workers coined the name only after the oxidase function, i.e. the reaction with oxygen, was demonstrated by the light reversibility of inhibition of respiration (2). The direct demonstration of the binding of oxygen with cytochrome *o* to give a compound spectrally resembling the CO compound was obtained by us only much later (6), with the availability of low temperature photolysis methods.

A new era of oxidase research.....

The explosion of interest in research on bacterial oxidases is largely the consequence of molecular cloning of structural genes, purification of oxidase complexes and the recognition of a much greater diversity of oxidase types than had hitherto been suspected. In addition to the long recognized existence of oxidases containing cytochromes *o*, *a*₁, *a*₃, *d* and *d*₁ as ligand-binding haems, it has become clear that cytochrome *b* can act in this capacity and that certain oxidases contain previously unrecognized haem types. For example, the cytochrome *o*-type oxidase of *Escherichia coli* is now known to contain a novel haem, haem O (8). To add to the possibility of confusion (or interest, depending on one's point of view) it is now known that a haem site in an oxidase protein can be occupied by more than one class of haem. For example, in the cytochrome *o*-type oxidase of *E. coli*, a variant of the oxidase is known in which both haems are the O-type, in contrast to the 'normal' oxidase in which one haem is O (ligand-binding) and the other is B (9).

Quite naturally, those working on these oxidases have sought a way of distinguishing, in the name, between haems in a given oxidase. It is particularly valuable to demonstrate in the name which of the haems is, or are, capable of

binding oxygen (the substrate) and other ligands such as CO and cyanide. Precisely this requirement arose in the 1930s, when Keilin (3) described 'two...compounds having the same haem nucleus. One of these compounds (*a*) is not autoxidizable and does not combine with potassium cyanide or carbon monoxide, while the other (*a*₃) is autoxidizable and combines with both'. Why was the name *a*₃ chosen? Again, Keilin himself makes this perfectly clear. 'We shall designate this new component as *a*₃ in order to distinguish it from cytochromes *a*₁ (589 nm) and *a*₂ (630 nm) which are known to occur only in a few bacteria devoid of cytochrome *a*' (We know now that certain bacteria do possess cytochrome *a*₃ as well as 'a₂', but the principle of nomenclature remains valid.) Cytochrome *a*₁ was the name given to a cytochrome observed in *Acetobacter* and a few other bacteria. This name is now little used because in many bacteria, but not all (10), cytochrome *a*₁ has been shown to be a high-spin *b*-type cytochrome that is a component of the cytochrome *bd* complex. Cytochrome *a*₂ was the early name for cytochrome *d*. Like most types of '*a*', it was subsequently shown not to contain haem A but, in this case, a chlorin haem that gives rise to the unique spectral features of this type of oxidase.

....but do we need new names, and are some of them really new?

It is with supposed deference to the name '*aa*₃' that the names *bo*₃, *bb*₃ and others have been adopted by some workers in this field over the past few years (for a survey, see 10). The subscript '3' denotes the ligand-binding haem. The name was first used for the *E. coli* cytochrome *o*-type oxidase by Puustinen & Wikström (8), who suggested that this oxidase be called *bo*₃ (when the haems are B and O) or *oo*₃ (when both haems are O). It is claimed that the nomenclature is useful because it shows the haem types (we agree), that it shows which is the ligand-binding haem (it does, but a system for doing so already exists, see below), and that it is 'historically pertinent' (we disagree). If '*o*₃' has the same etymological root as *a*₃, where are cytochromes *o*₁ and *o*₂?

We accept that the subscript is useful in signifying which haem binds oxygen. This is a particularly valuable feature in cases like cytochromes *oo* and *bb* where the low-spin (electron-conducting) and high-spin (ligand-binding) haems are of the same chemical structure differing only in their axial ligation to the protein and thus their ability to bind oxygen. However, there exists already an internationally validated recommendation (4) that serves precisely this function. It is simple and unambiguous, namely that the 5-coordinate ligand-binding haem is primed as in *bo*' (for the *E. coli* oxidase) and *aa*' (for the mitochondrial oxidase).

Unfortunately, this nomenclature has never 'caught on' amongst students of the mitochondrial oxidase, probably because the official recommendation lagged many years behind the

Keilin nomenclature. In contrast, the names oo_3 , bo_3 and bb_3 are recent and not yet, we hope, too embedded in the literature. We agree with the IUB Nomenclature Committee that current usage of aa_3 is unlikely to be abandoned and suggest that this term, being universally understood, be retained as the acceptable exception to the rule. However, our view is that in other cases IUB recommendations should be implemented. Thus the cytochrome o -containing oxidase of *E. coli*, for example, would be called cytochrome bo' or oo' , depending on the haem composition.

Finally, we recognize that there may be those who will argue that the subscript '3' will be recognized by all as meaning ligand-binding and that concern over nomenclature is unnecessary and pedantic. Consider the name ' b_3 ', which is becoming more widely used to describe oxidases in which the ligand binding haem is not A or O, but B. Oxidases appearing to have two haems B are now recognized in several bacteria (10) and are especially interesting because one such oxidase may be the elusive high-affinity oxidase of nitrogen-fixing bacteroids. However, unlike the cases of bo' and aa_3 , the evidence for haem B being oxygen-binding in such oxidases, as demonstrated at low temperature or in photochemical action spectra, is sadly lacking. Furthermore, do the inventors of this name know (i) that the term ' b_3 ' has already been used to describe a microsomal cytochrome in non-photosynthetic plant tissues and (ii) that the IUB Nomenclature Committee has already suggested that 'this category is not sufficiently defined to warrant further use'?

An even more serious problem may lie ahead if the 'subscript 3' nomenclature is to be extended. How would protagonists of the bo_3 and bb_3 names describe a hypothetical oxidase in which the ligand-binding haem is C (11)? If the answer is ' c_3 ' then this would indeed be ill-advised. Cytochrome c_3 is already a well-

established name for a low-potential non-oxidase cytochrome, which already warrants a paragraph in the IUB nomenclature recommendations (4).

Summary

The table below summarizes our own views on the most appropriate names for some of the oxidases now being intensively investigated in many laboratories.

Preferred name	Other name(s)
aa_3 or aa' ; also caa_3 or cad'	
bo'	o , bo , bo_3 , $b_{562}o$
oo'	oo_3
bb' ; also cbb'	bb_3 , cbb_3

In conclusion, we suggest here, as in (7), that existing nomenclature recommendations be respected and adopted and that new names of doubtful historical relevance should not be used unless they have been ratified by the international biochemical community. The worst examples are those names which are second-hand or reincarnations of other names.

Call things by their right names...Glass of brandy and water! That is the current but not the appropriate name: ask for a glass of liquid fire and distilled damnation.

From Gregory's *Life of Hall*

Robert K. Poole*¹ and Britton Chance²

¹Division of Life Sciences, King's College London, Campden Hill Road, London W8 7AH, UK.

²Department of Biochemistry and Biophysics, School of Medicine, University of Pennsylvania, Philadelphia, PA 19104-6089, USA.

*For correspondence. Tel: +44 171 333 4275. Fax: +44 171 333 4500.

1. Calhoun, M.W., Thomas, J.W. & Gennis, R.B. (1994). The cytochrome oxidase superfamily of redox-driven proton pumps. *TIBS* **19**, 325–330.
2. Castor, L.N. & Chance, B. (1959). Photochemical determinations of the oxidases of bacteria. *J Biol Chem* **234**, 1587–1592.
3. Keilin, D. (1970). *The History of Cell Respiration and Cytochrome*. Cambridge: Cambridge University Press.
4. Nomenclature Committee of the International Union of Biochemistry (NCIUB) (1991). Nomenclature of electron-transfer proteins. Recommendations 1989. *Eur J Biochem* **200**, 599–611.
5. Poole, R.K. (1994). Oxygen reactions with bacterial oxidases and globins: binding, reduction and regulation. *Antonie Leeuwenhoek* **65**, 289–310.
6. Poole, R.K., Waring, A.J. & Chance, B. (1979). The reaction of cytochrome o in *Escherichia coli* with oxygen. Low-temperature kinetic and spectral studies. *Biochem J* **184**, 379–389.
7. Poole, R.K., Salmon, I. & Chance, B. (1994). The high-spin cytochrome o' component of the cytochrome bo -type quinol oxidase in membranes from *Escherichia coli*: formation of the primary oxygenated species at low temperature is characterized by a slow 'on' rate and low dissociation constant. *Microbiology* **140**, 1027–1034.
8. Puustinen, A. & Wikström, M. (1991). The heme groups of cytochrome o from *Escherichia coli*. *Proc Natl Acad Sci USA* **88**, 6122–6126.
9. Puustinen, A., Morgan, J.E., Verkhovsky, M., Thomas, J.W., Gennis, R.B. & Wikström, M. (1992). The low-spin heme site of cytochrome o from *Escherichia coli* is promiscuous with respect to heme type. *Biochemistry* **31**, 10363–10369.
10. Van der Oost, J., de Boer, A.P.N., de Gier, J.-W.L., Zumft, W.G., Stouthamer, A.H. & van Spanning, R. (1994). The heme-copper oxidase family consists of three distinct types of terminal oxidases and is related to nitric oxide reductase. *FEMS Microbiol Lett* **121**, 1–9.
11. Wood, P.M. (1984). Bacterial proteins with CO-binding b- or c-type haem. Functions and absorption spectroscopy. *Biochim Biophys Acta* **768**, 293–317.