***Trichoderma harzianum* genes induced during growth on *Rhizoctonia solani* cell walls**

Valérie Vasseur,[†] Marc Van Montagu and Gustavo H. Goldman[‡]

Author for correspondence: Marc Van Montagu. Tel: +32 9 2645170. Fax: +32 9 2645349.
e-mail: BITNET mamon@gengenp.rug.ac.be

Laboratorium voor
Genetica, Universiteit
Gent, K. L.
Ledeganckstraat 35,
B-9000 Gent, Belgium

***Trichoderma harzianum* is a biocontrol agent that attacks a range of economically important phytopathogenic fungi. In an attempt to identify genes specifically expressed by *T. harzianum* during growth on cell walls of *Rhizoctonia solani*, we carried out differential screening of an induced cDNA library. In this paper we report the analysis of the sequence and expression of two cDNA clones that encode putative mycoparasitism-related proteins of *T. harzianum*. One of these clones corresponds to a gene, *inda1*, that encodes a protein of 570 amino acids with a predicted molecular mass of 62853 Da. The predicted amino acid sequence of *inda1* showed a high degree of similarity with amino acid permeases from several other organisms. The other cDNA clone corresponds to a gene, *indc11*, that encodes a novel protein of 340 amino acids with a predicted molecular mass of 37010 Da. The use of this methodology should provide specific genetic markers to follow mycoparasitism by *Trichoderma* spp.**

Keywords: *Trichoderma harzianum*, *Rhizoctonia solani* cell walls, mycoparasitism

INTRODUCTION

Modern agriculture is very strongly dependent on fungicides. The repeated use of these pesticides has not only polluted the environment, but many pesticides have also become useless owing to the development of resistance among the target organisms. One possible alternative to synthetic fungicides is the use of biological control agents. Many natural biocontrol agents have been shown to be effective in controlling plant pathogens (see reviews by Chet, 1990; Manocha, 1991). However, biocontrol is not often used in commercial agricultural production because control of plant diseases with microbial agents has been less effective and reliable than with synthetic fungicides. The improvement of biocontrol efficacy requires a better knowledge of the mechanisms involved in the control of plant pathogens.

The fungus *Trichoderma harzianum* has been used as a biocontrol agent and shown to attack a range of econ-

omically important aerial and soil-borne fungal plant pathogens. In soil, *T. harzianum* strains have been used as antagonists against several plant-pathogenic fungi, for example *Rhizoctonia solani*, *Botrytis cinerea* and *Sclerotium rolfsii* (Chet, 1987). A great deal of information about the factors involved in the antagonistic properties of *T. harzianum* has been identified. These include volatile and non-volatile antibiotics, and hydrolytic enzymes such as chitinases, glucanases and proteases (Dennis & Webster, 1971a, b; Elad *et al.*, 1982; Ridout *et al.*, 1986, 1988; Claydon *et al.*, 1991; Geremia *et al.*, 1991; Graeme-Cook & Faull, 1991; Ulhoa & Peberdy, 1991; Avent *et al.*, 1992; De La Cruz *et al.*, 1992; Di Pietro *et al.*, 1993; Harman *et al.*, 1993; Lorito *et al.*, 1993). Recently, two mycoparasitism-related genes from *T. harzianum*, encoding an alkaline proteinase (Geremia *et al.*, 1993) and an endochitinase (Hayes *et al.*, 1994), have been isolated and characterized.

When mycoparasitic strains of *Trichoderma* spp. grow on host cell walls, induction and/or derepression of genes involved in mycoparasitism has been found to occur. In addition, glucose represses the activity and/or expression of hydrolytic enzymes important for the degradation of the cell walls (Geremia *et al.*, 1991, 1993). Our approach to identify factors determining mycoparasitic abilities of *T. harzianum* was to isolate genes of this species that show induced or increased expression during growth on R.

[†] Present address: ESMISAB-Laboratoire de Microbiologie et Biochimie, Technopôle Brest-Iroise, F-29280, Plouzané, France.

[‡] Present address: Universidade de São Paulo, Faculdade de Ciências Farmacêuticas de Ribeirão Preto, Via do Café S/N, 14040-903, Ribeirão Preto, SP, Brazil.

The EMBL accession numbers for the *inda1* and *indc11* nucleotide sequences reported in this paper are Z22594 and Z22221, respectively.

solani cell walls. Identification of the proteins encoded by these genes might help elucidate some of the molecular and cellular events involved in mycoparasite–host interactions.

To isolate the putative mycoparasitism genes, we carried out differential screening of an induced cDNA library. In this paper we report the analysis of the sequence and expression of two cDNA clones from this library. One corresponds to a gene encoding a putative amino acid permease (*inda1*), and the other corresponds to a gene encoding a novel polypeptide (*indc11*).

METHODS

Fungal strains and growth media. *T. harzianum* (IMI 206040) was grown in Erlenmeyer flasks containing 50 ml minimal medium (MM: 0.2% asparagine, 0.2% NaNO₃, 0.05% MgSO₄·7H₂O, 0.05% KCl, 0.1% KH₂PO₄, pH 6.0) supplemented with 2% (w/v) glucose on a rotatory shaker (150 r.p.m.) at 28 °C for 3 d. The mycelium was then harvested, rinsed with sterile water and transferred to Erlenmeyer flasks with the same medium containing either 2% glucose or 0.1% cell walls of *R. solani* (prepared as described by Goldman *et al.*, 1992a). Mycelium was harvested after 24, 48 and 72 h, lyophilized, and stored at –20 °C.

cDNA library. The cDNA library was constructed using poly(A)⁺-RNA obtained from *T. harzianum* grown for 24 h in MM containing cell walls of *R. solani* (1 mg ml^{–1}) as sole carbon source (Goldman *et al.*, 1992b). Total RNA and poly(A)⁺-RNA were isolated from fungal mycelia according to a previously described protocol (Jones *et al.*, 1985), adapted for the large-scale isolation of RNA and using a poly-A tract mRNA isolation kit (Promega).

Differential screening. This was carried out essentially as described by Gasser *et al.* (1989) and M. H. de S. Goldman *et al.* (1992). Bacterial colonies (3000) were stored in microtitre plates containing Luria–Bertani (LB) medium supplemented with ampicillin. Duplicate nylon replica filters (Hybond-N, Amersham) were used on each of the plates and treated as recommended by the manufacturer. Radioactive plus/minus cDNA probes were synthesized from poly(A)⁺-RNA of mycelium grown on glucose ('minus') or cell wall ('plus') medium. Hybridization probes were synthesized from 1 µg of poly(A)⁺-RNA in a reaction similar to the first-strand cDNA synthesis. Modifications introduced were the use of 100 µCi (3.7 MBq) [³²P]dCTP for the first hour of reaction and the use of random hexanucleotides as primers. Unlabelled dCTP was then added to a final concentration of 0.5 mM and the reaction was continued for 30 min. The labelled cDNA/RNA hybrids were purified on Nensorb columns (Nensorb). In preparation for hybridization, the probes were denatured and the RNA was hydrolysed by adding 60 µl 1.0 M NaOH, followed by 30 min incubation at room temperature. The base was neutralized by adding 60 µl 1.0 M HCl and 60 µl 20× SSPE (1× SSPE: 0.15 M NaCl, 10 mM sodium phosphate, 1 mM EDTA, pH 7.4). Filters were prehybridized in a solution containing 6× SSC (1× SSC: 0.15 M NaCl, 0.015 M sodium citrate, pH 7.0), 1% (w/v) SDS, 100 µg denatured carrier DNA ml^{–1}, 0.25% milk powder for 1 h at 68 °C. The labelled cDNA was added to the prehybridization solution to a final concentration of 10⁶ c.p.m. ml^{–1} and incubated at 68 °C for 36–48 h. The probe-specific activities were adjusted to equality and equivalent amounts of radioactivity per ml of hybridization solution were used. Filters were washed sequen-

tially at 68 °C in 6× SSC, 0.1% SDS for 30 min, and in 2× SSC, 0.1% SDS for 30 min. Filters were exposed to Kodak-X Omat films overnight between intensifying screens at –70 °C.

DNA/RNA manipulations. Restriction enzyme digests and DNA ligation were performed according to the recommendations of the suppliers (Boehringer and Amersham). Plasmid DNA isolation and Southern blotting were all performed using standard procedures (Sambrook *et al.*, 1989). DNA probes were made using a primer system according to the manufacturer (Boehringer). The complete sequence of the *inda1* and *indc11* genes was determined by the dideoxy-chain termination method (Sanger *et al.*, 1977) from both strands using synthetic oligonucleotide primers.

For Northern analysis, *T. harzianum* was grown by inoculating MM supplemented with 2% (w/v) glucose with 3 × 10⁴ conidiospores per ml medium. The cultures were incubated in a reciprocal shaker at 28 °C for 3 d, harvested by filtration through a Whatman filter (no. 1), and washed thoroughly with TES buffer (50 mM Tris, 50 mM NaCl, 5 mM EDTA, pH 8.0). Equal amounts of mycelium were transferred to MM supplemented with 2% glucose or cell walls (1 mg ml^{–1}) of the phytopathogenic fungus *R. solani*. This last condition was defined throughout this work as cell-wall-grown, i.e. growth in liquid medium having a cell wall preparation of *R. solani* as sole carbon source (Goldman *et al.*, 1992a). Mycelium was collected, disrupted and total RNA extracted. In all Northern analysis experiments, RNA concentration was normalized by hybridization with a β -tubulin gene from *Trichoderma viride* (*tub1*; Goldman *et al.*, 1993).

RESULTS

Isolation of specific *T. harzianum* cDNA clones grown on cell walls

We isolated specific cDNA clones from a *T. harzianum* cell-wall-grown cDNA library by differential screening against probes from *T. harzianum* grown on glucose or on cell walls of *R. solani*, as described by Gasser *et al.* (1989) and M. H. de S. Goldman *et al.* (1992). The initial screening resulted in the identification of 65 clones (from 3000 clones of a non-amplified library) and a secondary screening in 15 cDNA clones that hybridized only to cDNA probes derived from *T. harzianum* grown on cell walls. For further analysis, we decided to choose only the clones which in Northern analysis experiments hybridized with mRNAs isolated from cell-wall-grown *T. harzianum* (see below). These preliminary experiments showed that only eight clones hybridized to mRNA obtained from *T. harzianum* grown on cell walls. Southern analysis experiments indicated that three of these clones cross-hybridized at high stringency (data not shown). Thus, we obtained at the end of our screening six different clones which are expressed specifically during growth on cell walls of *R. solani*. Two of these cDNA clones, *INDA1* and *INDC11*, were chosen for further analysis.

inda1 and *indc11* genes are expressed during growth of *T. harzianum* on cell walls

To clarify the conditions under which the *inda1* and *indc11* genes were expressed, RNA was isolated from *T. harzi-*

Fig. 1. Northern blot of mRNA obtained from *T. harzianum* grown on MM plus glucose (MM+G) or *R. solani* cell walls (MM+CW). Transcript levels of *prb1* (a), *inda1* (b), *indc11* (c) and *tub1* (d) in mycelia harvested 24 (lane 1), 48 (lane 2) and 72 (lane 3) h after transfer into MM+G or MM+CW.

anum grown on cell walls and Northern analyses were carried out (Fig. 1). One gene expressed during mycoparasitism by *T. harzianum* is the *prb1* gene, which encodes an alkaline proteinase (Geremia *et al.*, 1993). Its expression was repressed in the presence of glucose and increased during cell-wall-grown *T. harzianum* (Fig. 1a). The *inda1* gene specifies a single transcript of about 2.4 kb whereas the *indc11* gene specifies a single transcript of about 1.3 kb (Figs 1b and 1c). β -Tubulin mRNA expression was constant during growth on glucose and cell walls (Fig. 1d). Both genes hybridized only with mRNA obtained from *T. harzianum* grown on cell walls. They did not hybridize with mRNA isolated during growth of *T. harzianum* in the presence of glucose. These data indicated that *inda1* and *indc11* genes are specifically expressed in cell-wall-grown *T. harzianum* and repressed during growth on glucose.

The IND1A1 cDNA clone corresponds to a gene that encodes a putative amino acid permease

The IND1A1 cDNA clone contained an insert of 2.1 kb. The complete sequence of the *inda1* gene of *T. harzianum* is shown in Fig. 2. The cDNA has an ORF of 1710 nucleotides. To determine whether *inda1* is a member of a gene family, we performed DNA blot hybridizations with genomic DNA prepared from *T. harzianum* (Fig. 3a). After digestion of the genomic DNA with appropriate

Fig. 2. Nucleotide sequence and predicted amino acid sequence of the *T. harzianum* *inda1* gene (2.1 kb *Sfi*I–*Not*I fragment).

restriction enzymes, few fragments could be identified in the restriction map of the clone hybridized to the IND1A1 cDNA, indicating that *inda1* is probably a single-copy gene. The *inda1* ORF encodes a protein of 574 amino acid residues with a predicted molecular mass of 62853 Da and a calculated pI of 7.81. The codon usage of the *inda1* gene resembles that of other *Trichoderma* genes (Vanhanen, 1991; Goldman *et al.*, 1992a). Of the 61 codons, 9 are not used at all and 22 are used more than 10 times, coding for about 77% of the amino acid residues of IND1A1 (Table 1). Thus, no codon bias is apparent, a result in agreement with other genes expressed at low levels (Bennetzen & Hall, 1982).

Comparison of the amino acid sequence of IND1A1 with those of amino acid permeases from several other organisms revealed a striking identity (Fig. 4): 43.4% with GAP1 from *Saccharomyces cerevisiae*; 35.9% with HIP1 from *S. cerevisiae* (Tanaka & Fink, 1985); 34.1% with YCC5 from *S. cerevisiae* (Tanaka & Fink, 1985); 32.9%

Fig. 3. Southern blot of *T. harzianum* genomic DNA. DNA was prepared according to Garber & Yoder (1983) and digested with restriction endonucleases *Pst*I (lane 1), *Hind*III (lane 2), *Eco*RI (lane 3) or *Bam*HI (lane 4). Blots were probed with a 32 P-labelled 2.1 kb fragment from an *INDA1* cDNA clone (a) or a 1.2 kb fragment from an *INDC11* cDNA clone (b).

with LYSP from *Escherichia coli* (Steffes *et al.*, 1992); and 27.2% with CAN1 from *S. cerevisiae* (Hoffmann, 1985). The identity to the GAP1 protein from *S. cerevisiae* is the

highest value of similarity yet reported for amino acid permeases.

Experiments on the topology of membrane proteins have shown that structures predicted from hydropathy profiles are consistent with experimental methods of determining the structure of permeases (Ahmad & Bussey, 1986; Gött & Boos, 1988; Sophianopoulou & Scazzocchio, 1989). We analysed the hydropathy profile of the *INDA1* polypeptide (Fig. 5). The *INDA1* putative transporter lacks a hydrophobic N-terminal signal peptide, the whole N-terminal region of 75 amino acids being hydrophilic. The C-terminus of 39 amino acids is also hydrophilic. The remainder of the protein is highly hydrophobic, having 10 distinct putative transmembrane segments of an average hydropathy index greater than 30 (residues 98–114, 151–167, 182–198, 212–228, 300–316, 352–368, 401–417, 421–437, 477–493 and 511–527; Fig. 5). We also examined the hydropathy profile of the *INDA1* permease using a number of the sequenced prokaryotic and eukaryotic transport proteins mentioned above. The hydropathy profile of the *INDA1* permease was similar to all of the permeases, even though the number of putative transmembrane segments was not identical in all cases (data not shown; Fig. 5). Thus, based on the sequence homology and hydropathy profile of the *INDA1*, we propose that the *inda1* gene encodes a putative amino acid permease.

The *INDC11* cDNA clone corresponds to a gene that encodes a novel polypeptide

The *INDC11* cDNA clone contains an insert of approximately 1.2 kb. The cDNA has an ORF of 1020 nucleotides

Table 1. Codon usage of the *T. harzianum* *inda1* and *indc11* genes

Amino acid	Codon	Occurrence*		Amino acid	Codon	Occurrence*		Amino acid	Codon	Occurrence*		Amino acid	Codon	Occurrence*	
		<i>inda1</i>	<i>indc11</i>			<i>inda1</i>	<i>indc11</i>			<i>inda1</i>	<i>indc11</i>			<i>inda1</i>	<i>indc11</i>
Phe	TTT	9	4	Ser	TCT	13	1	Tyr	TAT	8	1	Cys	TGT	1	0
	TTC	36	11		TCC	10	8		TAC	17	14		TGC	4	2
Leu	TTA	0	0	Pro	TCA	2	1	Stop	TAA	1	1	Stop	TGA	0	0
	TTG	6	1		TCG	5	2		TAG	0	0		TGG	18	6
	CTT	12	0		CCT	6	4		CAT	2	2		CGT	5	0
	CTC	21	15		CCC	16	15		CAC	9	5		CGC	8	9
Ile	CTA	0	1	Thr	CCA	2	1	Gln	CAA	0	1	Ser	CGA	3	0
	CTG	16	9		CCG	3	5		CAG	8	7		CGG	0	1
	ATT	15	1		ACT	5	2	Asn	AAT	2	2	Arg	AGT	1	0
	ATC	26	17		ACC	14	9		AAC	13	10		AGC	4	7
Met	ATA	1	1	Ala	ACA	5	1	Lys	AAA	2	3	Gly	AGA	3	1
	ATG	12	9		ACG	3	5		AAG	22	17		AGG	0	2
Val	GTT	7	2		GCT	28	0	Asp	GAT	5	3		GGT	29	1
	GTC	32	18		GCC	26	2		GAC	16	14		GGC	30	21
	GTA	0	1		GCA	4	2		GAA	4	4		GGA	3	6
	GTG	3	5		GCG	1	2		GAG	17	11		GGG	0	1

* Total occurrence of this codon in the deduced amino acid sequence of the *inda1* and *indc11* genes.

GAP1	MSNTS---SYEKNPNLKNHGITI-----DS-EFLTQEPITIPNSGSAVSIDETSGSGKWQDFK--DSFKRVKPIEVDP-----	69
HIP1	MRRNPLKKEYWADVDFKPATSPAFENEKEST-TFVTELTSTKDSAPLSSKDSPGINQTTNDITSSDRFRNEDTEQED-----	80
YCC5	MSSS--KSLY---ELKDLKNSSTEIHATGDNEIEYFETGSDRPSQPHLYGHEHNTSAVRRFF--DSFKRADQGPQDEVEATQMNDLTSISPSSRQ	92
INDA1	MS-----KEESGHVTPKGDNVVDY-----QASTTVLPSEGPEDANW-----FTR-----NGL-----N	45
CAN1	MTNSK-----EDADIEEKHMY-----NEPVTLFHDVEASQTHHRRGSIPLKDEKSKELYPL-RSFPTRVNG	61
LYSP	M-----	1
GAP1	--NLSEAEKVAIIITAQT-----PLKHLKLNRLQMIAGGAGTGLLVGSGTALRTGGP-SLLIGWGSTGTHIYAMVMALGELAVIFP-ISGGFTTYATR	160
HIP1	---NLSGDLVSRHLLTAVGGAGTGLVYNTGAALSTGGPASLVIDWVIISTCLFTVINSLGELSAAFP-VVGGFNYSYHR	161
YCC5	AOELEKNSSDNIGANTGHKSDSLKKTIPRRHVLMIAGTGTGLLVGNGTALVHAGPAGLLIYAIMGSIYLCIIQACGEMALVYSNLTGGVYAYPV	192
INDA1	VDSFKKKHYGPGHVE-----LERPMKARLHMIAIGGSIAGGFVGGGALAKGGPGLFVDFLIIGIMFNVVYALGELAIMYP-VSGSFYTSAR	136
CAN1	EDTFSMEDGIDGDEGEVQNAE-VKRELKQRHIGMIALGGTIGTGLFGLSTPLNAGPVGALISYLFMGSLAYSVTQSLGEMAT-FIPVTSSTFTVSQR	159
LYSP	-----VSETKTTEAPGLRRELKARLHMTIAIGGSIAGTGLFVASGATISQAGPGGALLSYMLIGLMVYFLMTSLGELA-RYMPVSGSFATYGGN	88
GAP1	FIDESFGYANNFYMLQWLVLPLEIVAASITVNFVGTDPKYGDFVALFWLAIVIIMFVGKGYGEAEFVFSFIKVITVVGFIILGIIILNCGGPTGG-	259
HIP1	FIEPSFAFVNLNLYLAQWLVLPLELVAASITIKYWNKIN-SDAWVAIFYATIALANMLDVKSFGGETFVLNMIKILSIIGFTILGIVLSCGGPHGG-	259
YCC5	LMMVFGFAVAVWYCLQWLVCPLLEVTASMTIKYWTTSVN-PDVVFIIFYVLVITINIFGARGYAEAEFFNCCCKILMMTGFFILGIIIDVGGAGNDG-	290
INDA1	FIDPAAGFAMGWNVYLAQWAAVLPLELTVCGTITSYWNSEIT-TAAWISLFLGVIIINLFGALGYAEFFHASCFLAATVIFMIAFVLVGGGPKDGR	235
CAN1	FLSPAFAANGYMYFWSAITFALELSVVGQVIFQWTKYVPLA-AWISIFWVITIMNLFPVKYGEFEFVWASIKVLAIIIGFLIYCFCHVCGAGVTP-	257
LYSP	YVEEGFGFALGWNYWYNAVTIAVDLVAALVMSWFPDTP-GWISALFLGVIFLLNYISVRGFGAEYWFSLIKVTTVIVFIIVGVLMITGI-----	181
GAP1	---YIGGKYWHDPGA-----FAGDTPGAKFGVCSVFVTAASFAGSELVGLAASESVEPRKSVPKAAKQVFWIRITLFYILSLLMVLGLVPYNDKSLI-G	350
HIP1	---YIGGKYWHDPGA-----FVGHSSGTQFKGLCSVFVTAAFYSGIEMTAVSAAESKNPRETIKAAKRTFWLITASYVTITLITGLVPSNDPRLLNG	351
YCC5	---YIGGKYWHDPGA-----FNGKHAIDRFKGVVATLVTAFAFGGSEFIAITTAEGSNPKAIPGAQKQMIYRILFLFLATIIILGLFVPSNDQL-G	381
INDA1	YHEYWGARYWYDPGA-----FKNG-----FKGFCSVFVTAASFSGTELVGLAAEAESTNPTKNMPGAIKQVFWIRITIFYILGLFFVGLLINSDDALLSS	325
CAN1	---VGFYWRNPAGAGPGIISKDKNEGRFLGWSSLIINAAFTFGTGLVGTAGEAANPRKSVPRATKKVFRILTFYIGLSLFIIGLLVPSNDPKLTQS	353
LYSP	---FKGA-----QPAGSNWTIGEAPFAGGFAAMIGVAMIVGFSFGTGLIGIAAGESEDPAKNIPRAVRQVFWIRILLFYVFAILIISLIIPYDPSLLRN	274
GAP1	ASSVDAASPFIKIAIKTHGKGLPSVNVVILIAVLVSGNSAIYACSRMVALAEQRFLEIFSYVDRKGRPLVGIATVSAFGLIAFVAASKKEGEVFNW	450
HIP1	SSSVDAASSPLVIAIENGIGKGLPSLMNATILIAVSVANSVAVYACSRCHVAMAHIGNLPKFLNRVDKGRPMNAILTLFGLLSFVAASDKQAEVFTW	451
YCC5	TSGGGTASPYVIAVASHGVRVVPHFINAVILLSVLSMANSSVYSSARLFLTLSEGGYAPKVFYSYIDRAGRPLIAMGVSALFAVIAFCAASPKKEGVFTW	481
INDA1	AAAYDSKASPYVIVAGYAGLKGDFHMLVILASVLSIGVSGVSGSRTLTALAQGGYAPKLFYIDKSGRPLPSVIFLILFGFIAYVSLDATGPVVDW	425
CAN1	TSYVST--SPFIIAENSQTKVLPHFIFNAVILTTIIISAANSNIYVGSRTILFGLSKNKLAPKFLSRTTKGGVPYIAVFVTAAGALAYMETSGGDKVFEW	451
LYSP	DV-KDISVSPFTLVFHAGLLSAAAVMNAVILTAVL SAGNSGMYASTRMLYLACDQKAPRI FAKLSRGVPRKAAVCTTVIAGLCFLTSMFGNQTVYLW	373
GAP1	LLALSGLSLFTWGGICICHIRFRKALAAQGRGLDELFSKSPTVGWGSYGLFMVIMFIAQFYVALF--PVGDS-PAEGFFEAYLSFPLVMVYIGNK	547
HIP1	LSALSGLSTIFCUMAINLSHIFRQAMKQVQERSLDELPIISQTVGKGSYGFIVLFLVLIASFW-----TFSVPI-----R	522
YCC5	LLAISGLSGLFTWTAICLSHLRFRAMKQVQGRSLGELGFSQTVGWGSYACIMMILIAQFWAIA--PIGEGKLDQAQFFENYLAAMPILIALYVGYK	579
INDA1	LLAISGLAALFTWGSVCLAHIRFRKAKYHGHITLDEIPFKAAGGVGYSGYLGLFCIVILMAQFYTAIAAPPSPGSGVGTAEFFKQYLAAPVVLGFWIVGW	525
CAN1	LLNITGVAGFFAWLFIISISHIRFMAKLYRGISRDELFPKAKLMPGLAYATFMTIIIIIGGTFAT-----APKFNGVSFAAAYISVFLFVAVILFQ	545
LYSP	LLNTSGMTGFIATWGLIAISHYFRRGYVLQGHINDLPYRSGFFLPGIFAFILCLITLQGNYEAFKDTIDWGGVAA-----TYIGIPLFLIIWFGYK	468
GAP1	-IYKRNWKLFIAPAEKMDIDTGRREVLDLLKQEI AEEKAIMATKPRWYRIWNF-WC	601
HIP1	-RFRSQRRIIL-----	532
YCC5	-VWMDKWLFIADKIDLDSDHRIIDFELIKQEDDEYRERLRNGPYKRVVAF-WC	633
INDA1	-LWKQ--PFLRTKNIDVDTGLREFDWD-----EINAERTRIAPLPAWRRIIHTFX	574
CAN1	CIFRC--RFIWKIGDVIDSDRRDIE-AIVWEDHE-----PKTFMDKFWNVV-A	590
LYSP	LIGKTH--FVRYSEMKFPQNDKK-----	489

Fig. 4. Comparison of the amino acid sequence deduced for the *T. harzianum* INDA1 protein with amino acid permeases from *S. cerevisiae* (GAP1, HIP1, YCC5 and CAN1), and *E. coli* (LYSP). See Results for source of data. Asterisks indicate identical residues, dots indicate similar residues and hyphens indicate gaps inserted into the sequences to maximize alignment.

(Fig. 6) which encodes a protein of 340 amino acid residues with a predicted molecular mass of 37010 Da and a calculated pI value of 8.25. The *indc11* nucleotide and amino acid sequences do not display a high degree of homology with any sequence deposited in the databanks. A typical N-terminal sequence resembling signal sequences for excreted proteins was not found (Intelligenetics), suggesting that the INDC11 is not a secreted protein. The DNA sequence surrounding the translation initiation site ATG is CCACAATGGC, similar to the consensus sequence CCACCATGGC (Kozak, 1984). The consensus sequence AATAAA, which is thought to be involved in the 3'-end formation and polyadenylation of the precursor mRNA (see review by Ballance, 1991), was found in the *indc11* gene at nucleotide position 1351, 19 nucleotides upstream from the apparent poly(A) addition site. No other significant sequence similarities were identified in

computer-assisted searches of the databanks. Codon usage data for the *T. harzianum indc11* gene are presented in Table 1. The coding region of the *indc11* gene of *T. harzianum* contains approximately 55 mol% G + C. Of the 61 codons, 6 are not used at all and 12 are used more than 10 times, coding for about 56% of the amino acid residues of INDC11. Thus codon bias is apparent, a result in agreement with other genes expressed at high levels (Bennetzen & Hall, 1982; Ballance, 1991). The bases G and C are generally preferred at the third position of the codons. The codon usage of the *T. harzianum indc11* gene resembles that of other *Trichoderma* genes (Vanhanen, 1991; Goldman *et al.*, 1992a).

To determine whether *indc11* is a member of a gene family, we performed DNA blot hybridizations with genomic DNA prepared from *T. harzianum* (Fig. 3b). After

Fig. 5. Hydropathy profiles of the predicted amino acid sequence of the *T. harzianum* IND1 amino acid transporter (a), *S. cerevisiae* GAP1 general amino acid permease (b), and *A. nidulans* PRNB proline transport protein (c). The profiles of the amino acid transporters are plotted according to Kyte & Doolittle (1982) from the N-terminus to the C-terminus by averaging hydropathy values over a window of 15 residues.

digestion of the genomic DNA with appropriate restriction enzymes, few fragments that could be identified in the restriction map of the clone hybridized to the IND11 cDNA, suggesting that *indc11* is a single-copy gene.

DISCUSSION

We have described the isolation and characterization of two cDNA clones that correspond to genes encoding novel putative mycoparasitism-related proteins of *T. harzianum*. These two clones were obtained by differential screening of an induced library and correspond to genes that are expressed during growth on *R. solani* cell walls. One of them corresponds to a gene, *inda1*, that encodes a putative amino acid permease whereas the other corresponds to a gene, *indc11*, which is expressed at high levels during mycoparasitism. The *indc11* gene encodes a novel polypeptide that does not show any similarity with other proteins deposited in the databanks.

Although the *inda1* gene encodes a presumptive amino acid transporter that shows high similarity to a general amino acid permease of *S. cerevisiae* (see Results), the specificity of the permease activity of IND1 remains to be elucidated. The hydropathy profile of the IND1 transport protein is similar to many different permeases (see Results). There is no classical cleavage leader sequence in the IND1 amino acid permease. All fungal and many bacterial and mammalian transporters lack an N-terminal hydrophobic signal peptide. Blobel (1980) suggested that one or more of the hydrophobic segments of these proteins might function as an 'internal signal peptide'. Further investigation is required to determine the actual role of IND1 in mycoparasitic interactions, and how its regulation can be affected by a diversity of environmental factors.

It has been proposed that the mycoparasitic activity of *Trichoderma* proceeds in three major steps (Chet, 1987). Initially, the mycoparasite hyphae grow towards the host hyphae (Chet *et al.*, 1981). Then the parasite attaches to the target hyphae, presumably using a host lectin (Elad *et al.*, 1983a, b; Barak *et al.*, 1985), and appressoria-like structures coil around the attacked cells (Chet *et al.*, 1981; Elad *et al.*, 1983b). Concurrently, degradation of 1,3- β -glucans and chitin from the host cell wall is observed (Elad *et al.*, 1983b). Finally, the mycoparasite penetrates and/or lyses the host hyphae (Chet *et al.*, 1981), releasing the cellular contents and providing nutrients to sustain its growth. Both mechanical pressure and cell-wall-degradation by hydrolytic enzymes are probably involved in the process of penetration. Extracellular enzymes that hydrolyse the main chemical constituents of the fungal cell wall, i.e. chitin, glucans and proteins, were detected when *T. harzianum* was grown on *R. solani* mycelia or cell walls as sole carbon source (Ridout *et al.*, 1988; Geremia *et al.*, 1991). The enzymes appeared sequentially: first an alkaline proteinase was produced followed by glucanases and chitinases (Geremia *et al.*, 1993; G. H. Goldman & D. Jacobs, unpublished results). We have recently purified the alkaline proteinase specifically induced by *R. solani* cell walls and chitin (Geremia *et al.*, 1991) and cloned the corresponding gene (*prb1*; Geremia *et al.*, 1993). We suggested that PRB1 was involved in the degradation of the phytopathogen cell walls, membranes and possibly proteins released after the lysis of the host, which could then be used as nutrients by the mycoparasite. Thus, the

- Dennis, C. & Webster, J. (1971a). Antagonistic properties of species-groups of *Trichoderma*. I. Production of non-volatile antibiotics. *Trans Br Mycol Soc* **57**, 25–39.
- Dennis, C. & Webster, J. (1971b). Antagonistic properties of species-groups of *Trichoderma*. II. Production of volatile antibiotics. *Trans Br Mycol Soc* **57**, 41–48.
- Di Pietro, A., Lorito, M., Hayes, C. K., Broadway, R. M. & Harman, G. E. (1993). Endochitinase from *Gliocladium virens*: isolation, characterization, and synergistic antifungal activity in combination with gliotoxin. *Phytopathology* **83**, 308–313.
- Elad, Y., Chet, I. & Henis, Y. (1982). Degradation of plant pathogenic fungi by *Trichoderma harzianum*. *Can J Microbiol* **28**, 719–725.
- Elad, Y., Barak, R. & Chet, I. (1983a). Possible role of lectins in mycoparasitism. *J Bacteriol* **154**, 1431–1435.
- Elad, Y., Chet, I., Boyle, P. & Henis, Y. (1983b). Parasitism of *Trichoderma* spp. on *Rhizoctonia solani* and *Sclerotium rolfsii* – scanning electron microscopy and fluorescence microscopy. *Phytopathology* **73**, 85–88.
- Garber, R. C. & Yoder, O. C. (1983). Isolation of DNA from filamentous fungi and separation into nuclear, mitochondrial, ribosomal, and plasmid components. *Anal Biochem* **135**, 416–422.
- Gasser, C. S., Budelier, K. A., Smith, A. G., Shah, D. M. & Fraley, R. T. (1989). Isolation of tissue specific cDNAs from tomato pistils. *Plant Cell* **1**, 15–24.
- Geremia, R., Jacobs, D., Goldman, G. H., Van Montagu, M. & Herrera-Estrella, A. (1991). Induction and secretion of hydrolytic enzymes by the biocontrol agent *Trichoderma harzianum*. In *Biotic Interactions and Soil-borne Diseases* (Developments in Agricultural and Managed-Forest Ecology Series vol. 23), pp. 181–186. Edited by A. B. R. Beemster, G. J. Bollen, M. Gerlagh, M. A. Ruissen, B. Schippers & A. Tempel. Amsterdam: Elsevier.
- Geremia, R., Goldman, G. H., Jacobs, D., Ardiles, W., Vila, S. B., Van Montagu, M. & Herrera-Estrella, A. (1993). Molecular characterization of the proteinase-encoding gene, *prb1*, related to mycoparasitism by *Trichoderma harzianum*. *Mol Microbiol* **8**, 603–613.
- Goldman, G. H., Geremia, R. A., Caplan, A. B., Vila, S. B., Villaruel, R., Van Montagu, M. & Herrera-Estrella, A. (1992a). Molecular characterization and regulation of the phosphoglycerate kinase gene from *Trichoderma viride*. *Mol Microbiol* **6**, 1231–1242.
- Goldman, G. H., Demolder, J., Dewaele, S., Herrera-Estrella, A., Geremia, R. A., Van Montagu, M. & Contreras, R. (1992b). Molecular cloning of the imidazoleglycerolphosphate dehydratase gene of *Trichoderma harzianum* by genetic complementation in *Saccharomyces cerevisiae* using a direct expression vector. *Mol & Gen Genet* **234**, 481–488.
- Goldman, G. H., Temmerman, W., Jacobs, D., Contreras, R., Van Montagu, M. & Herrera-Estrella, A. (1993). A nucleotide substitution in one of the β -tubulin genes of *Trichoderma viride* confers resistance to the antimetabolic drug methyl benzimidazole-2-yl-carbamate. *Mol & Gen Genet* **240**, 73–80.
- Goldman, M. H. de S., Pezzotti, M., Seurinck, J. & Mariani, C. (1992). Developmental expression of tobacco pistil-specific genes encoding novel extensin-like proteins. *Plant Cell* **4**, 1041–1051.
- Gött, P. & Boos, W. (1988). The transmembrane topology of the sn-glycerol-3-phosphate permease of *Escherichia coli* analysed by *pboA* and *lacZ* protein fusions. *Mol Microbiol* **2**, 655–663.
- Graeme-Cook, K. A. & Faull, J. L. (1991). Effect of ultraviolet-induced mutants of *Trichoderma harzianum* with altered antibiotic production on selected pathogens *in vitro*. *Can J Microbiol* **37**, 659–664.
- Harman, G. E., Hayes, C. K., Lorito, M., Broadway, R. M., Di Pietro, A., Peterbauer, C. & Tronsmo, A. (1993). Chitinolytic enzymes of *Trichoderma harzianum*: purification of chitobiosidase and endochitinase. *Phytopathology* **83**, 313–318.
- Hayes, C. K., Klemsdal, S., Lorito, M., Di Pietro, A., Peterbauer, C., Nakas, J. P., Tronsmo, A. & Harman, G. E. (1994). Isolation and sequence of an endochitinase-encoding gene from a cDNA library of *Trichoderma harzianum*. *Gene* **138**, 143–148.
- Hoffmann, W. (1985). Molecular characterization of the *CAN1* locus in *Saccharomyces cerevisiae*. A transmembrane protein without N-terminal hydrophobic signal sequence. *J Biol Chem* **260**, 11831–11837.
- Jones, J. D. G., Dunsmuir, P. & Bedbrook, J. (1985). High level expression of introduced chimaeric genes in regenerated transformed plants. *EMBO J* **4**, 2411–2418.
- Kozak, M. (1984). Selection of initiation sites by eucaryotic ribosomes: effect of inserting AUG triplets upstream from the coding sequence for preproinsulin. *Nucleic Acids Res* **12**, 3873–3893.
- Kyte, J. & Doolittle, R. F. (1982). A simple method for displaying the hydropathic character of a protein. *J Mol Biol* **157**, 105–132.
- Lorito, M., Harman, G. E., Hayes, C. K., Broadway, R. M., Tronsmo, A., Woo, S. L. & Di Pietro, A. (1993). Chitinolytic enzymes produced by *Trichoderma harzianum*: antifungal activity of purified endochitinase and chitobiosidase. *Phytopathology* **83**, 302–307.
- Manocha, M. S. (1991). Physiology and biochemistry of biotrophic mycoparasitism. In *Soil and Plants* (Handbook of Applied Mycology vol. 1), pp. 273–300. Edited by D. K. Arora, B. Rai, G. K. Mukerji & G. R. Knudsen. New York: Marcel Dekker.
- Ridout, C. J., Coley-Smith, J. R. & Lynch, J. M. (1986). Enzyme activity and electrophoretic profile of extracellular protein induced in *Trichoderma* spp. by cell walls of *Rhizoctonia solani*. *J Gen Microbiol* **132**, 2345–2352.
- Ridout, C. J., Coley-Smith, J. R. & Lynch, J. M. (1988). Fractionation of extracellular enzymes from a mycoparasitic strain of *Trichoderma harzianum*. *Enzyme Microb Technol* **10**, 180–187.
- Sambrook, J., Fritsch, E. F. & Maniatis, T. (1989). *Molecular Cloning: a Laboratory Manual*, 2nd edn. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory.
- Sanger, F., Nicklen, S. & Coulson, A. R. (1977). DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci USA* **74**, 5463–5467.
- Sophianopoulou, V. & Scazzocchio, C. (1989). The proline transport protein of *Aspergillus nidulans* is very similar to amino acid transporters of *Saccharomyces cerevisiae*. *Mol Microbiol* **3**, 705–714.
- Steffes, C., Ellis, J., Wu, J. & Rosen, B. P. (1992). The *lysP* gene encodes the lysine-specific permease. *J Bacteriol* **174**, 3242–3249.
- Tanaka, J.-i. & Fink, G. R. (1985). The histidine permease gene (*HIP1*) of *Saccharomyces cerevisiae*. *Gene* **38**, 205–214.
- Ulhoa, C. J. & Peberdy, J. F. (1991). Regulation of chitinase synthesis in *Trichoderma harzianum*. *J Gen Microbiol* **137**, 2163–2169.
- Vanhanen, S. (1991). *Isolation and characterization of genes involved in basic metabolism of the filamentous fungus Trichoderma reesei*. Academic dissertation, University of Helsinki (Technical Research Centre of Finland Publications no. 75).

Received 22 September 1994; accepted 5 December 1994.