Microbiology Comment provides a platform for readers of *Microbiology* to communicate their personal observations and opinions in a more informal way than through the submission of papers.

Most of us feel, from time to time, that other authors have not acknowledged the work of our own or other groups or have omitted to interpret important aspects of their own data. Perhaps we have observations that, although not sufficient to merit a full paper, add a further dimension to one published by others. In other instances we may have a useful piece of methodology that we would like to share.

The Editors hope that readers will take full advantage of this new section and use it to raise matters that hitherto have been confined to a limited audience.

Jon Saunders
Editor-in-Chief

An energetic move for Microbiology

In January 1996, the Editors of *Microbiology* are introducing a new subject category – *Bioenergetics and Transport* – for papers published in the journal.

Why introduce another section? The proposal stems from the fact that much of the most exciting work in contemporary bioenergetics is performed with micro-organisms. Of course, bacteria and yeasts, in particular, have always been at the forefront of bioenergetics research. Consider the recognition of fermentation and the unravelling of its diverse pathways, the discovery of cytochromes, the elucidation of chemiosmotic principles, determination of the path of carbon in photosynthesis, and kinetic and mechanistic descriptions of transport. All these and more were achieved as a result of exploiting the experimental advantages of micro-organisms.

Bioenergetics has entered a new era, largely as a result of the availability and imaginative use of molecular genetic tools. An example of a current 'hot topic' is the dissection of structure-function relationships in complex respiratory proteins like terminal oxidases. These enzymes display an extraordinary similarity of

function and mechanism in bacteria and mitochondria, but the bacterial versions are structurally much simpler and, of course, much easier to handle. The possibility of solving seemingly intractable problems has attracted many 'mitochondriacs' to study bacteria. There have been equally dramatic leaps forward in our understanding of the transport of nutrients into cells (and of antibiotics, unwanted ions and polypeptides out), the energetics of intact cells, and photosynthetic electron transfer proteins, culminating in 1988 with a Nobel prize to Deisenhofer, Huber and Michel for elucidation of the structure of the membrane-bound photosynthetic reaction centre of the purple bacterium *Rhodospseudomonas viridis*. Rapidly developing areas are the bioenergetics of pathogens, to which a symposium was devoted at the Society for General Microbiology's Golden Jubilee meeting at Bath, and the regulation of expression of genes encoding components of energetic mechanisms.

Microbiology is already very active in publishing papers in this field. A survey of papers published in the first 14 months of the new journal revealed more than 50 papers dealing with various aspects of energetics and transport. About half of these appeared in the rather general categories of *Biochemistry* and *Physiology and Growth*. The Editors hope that the new section will provide an appropriate focus for developments in this important area and attract first-rate papers in the field. We invite all those working on any aspects of microbial energetics and transport to consider publishing their work in *Microbiology* and earmarking it for the new section.

Robert K. Poole (Editor)

Microbiology Group, Division of Life Sciences, King's College London, Campden Hill Road, London W8 7AH, UK.

Tel: +44 171 333 4275. Fax: +44 171 333 4500.

Soft X-rays – a novel technique for the study of bacteria

Pesticides, apart from controlling pests, have given rise to numerous environmental concerns. However, microbial species belonging to the genera *Arthrobacter*, *Bacillus*, *Nocardia* and *Pseudomonas* help by utilizing a number of

pesticides by oxidative, reductive or hydrolytic processes (1). It is sometimes tedious and time consuming to spot these few pesticide-degrading strains. In this work, we report for the first time the study of some pesticide-degrading *Pseudomonas* spp. and a number of non-degrading organisms with soft X-rays.

Cultivation of *Pseudomonas ovalis*, *P. tralucida* Pt⁺, *P. tralucida* Ptm⁺ and *Bacillus subtilis* was as described previously (4). *Rhizobium meliloti* and *R. leguminosarum* bv. *trifolii* were cultivated in TY medium, *Salmonella typhimurium* was cultivated in nutrient broth and *Escherichia coli* in Luria broth in accordance with standard conditions. *Frankia* sp. was cultivated as described elsewhere (2).

The bacterial cells, after reaching the exponential phase of growth, were pelleted at low speed (5000 g) for 10 min. The cells were then treated with different fixatives and stains, such as ethyl alcohol, formaldehyde, glutaraldehyde, ruthenium red, osmium tetroxide and uranyl acetate. Different concentrations (0.05–3.0 %) and various combinations were also tried. The solutions were allowed an infiltration time of 120 min at 25 °C.

Smears of the organisms were then made on aluminium foil (40 × 40 × 0.04 mm) and dried overnight using calcium chloride as desiccant. The organisms treated thus were studied with a SOFTEX ISTV 25 soft X-ray system

► GUIDELINES

Communications should be in the form of letters and should be brief and to the point. A single small Table or Figure may be included, as may a limited number of references (cited in the text by numbers, and listed in alphabetical order at the end of the letter). A short title (fewer than 50 characters) should be provided.

Approval for publication rests with the Editor-in-Chief, who reserves the right to edit letters and/or to make a brief reply. Other interested persons may also be invited to reply. The Editors of *Microbiology* do not necessarily agree with the views expressed in *Microbiology Comment*.

Contributions should be addressed to the Editor-in-Chief via the Editorial Office.

hooked up to a Sigma II real time analyser for digital processing of the radiographic image.

The ruthenium preference index was computed by developing an equation given below:

$$\text{ruthenium preference index} = \frac{S \cdot M}{A}$$

where S = stain-induced levels, M = magnification and A = area of bacterial radiodigital image (mm^2). Observations at 17 kV, 3.5 mA and integration of 64 frames were found to be the best for all organisms.

Observations of radiographs of the organisms treated only with fixatives like formaldehyde and glutaraldehyde produced underdeveloped radiographs. The identification of the organisms was not possible since the radiographs were too transparent. However, glutaraldehyde was relatively better than formaldehyde. Stains like 0.05 % uranyl acetate and ruthenium red gave better results where dark and light levels became visible. The exception was in the case of 0.05 % osmium tetroxide, where an opaque overdeveloped radiograph in some places coupled with transparent patches was obtained.

However, double preparatory procedures gave different results. Combinations of 0.05 % ruthenium red and 0.05 % uranyl acetate produced three levels on the radiographs of the organisms. Glutaraldehyde (2%, w/v) with 0.5% uranyl acetate and 0.05 % osmium tetroxide made the radiographs too dark for observation. Glutaraldehyde (2%) and 0.05 % ruthenium red in a 1 : 1 ratio gave the best results. Here the radiographs of micro-organisms produced four grey levels. Hence, this combination was used for pre-treatment of all samples.

The radiograph shows *P. ovalis* as an elongated spindly shadow (Fig. 1) while *R. meliloti* produced a broad stump-like shadow. It should be noted here that radiographs are density oriented and not morphology oriented.

The shadows were broken up into pixels for digital image analysis. The I and IV pixel range

was found to vary amongst hexachlorocyclohexane-degrading and non-degrading strains. The IV pixel percentage of degrading organisms ranged from 55 to 60 and that of non-degrading organisms from 32 to 45 (Table 1). *P. ovalis*, a hexachlorocyclohexane-degrading organism, showed IV pixel percentage of 60. *S. typhimurium* showed IV pixel percentage of 40. The values of II and III pixels merge for degrading and non-degrading organisms. The data substantiated that IV pixel is important for demarcation of the organisms studied. The increase in IV pixel percentage might be attributed to more ruthenium absorption of degrading *Pseudomonas* spp. The absorption would be greater when polysaccharides are produced. This is because ruthenium, in addition to binding with cells, would bind polyanions of acidic polysaccharides (3).

The above results were confirmed by calculating the ruthenium preference index. These calculations showed degrading *Pseudomonas* spp. to have a high ruthenium preference index (see Table). This was due to higher binding to ruthenium. The results also indicated that ruthenium could act as a probe stain for observation under soft X-rays.

Fig. 1. Radiograph of *P. ovalis*.

Table 1. Ruthenium preference amongst hexachlorocyclohexane-degrading and non-degrading organisms

Organism	IV pixel percentage	Ruthenium preference index
Degrading:		
<i>P. ovalis</i>	60	25
<i>P. tralucida</i> Pt ⁺	56	9
<i>P. tralucida</i> Ptm ⁺	55	11
Non-degrading:		
<i>B. subtilis</i>	32	6.5
<i>R. meliloti</i>	39	3.5
<i>R. leguminosarum</i> bv. <i>trifoli</i>	33	3.8
<i>S. typhimurium</i>	40	4.6
<i>E. coli</i>	40	5.0
<i>Frankia</i> sp.	45	3.0

Note: SD values not above 0.6.

The results showed that soft X-rays could be used as a tool to study bacteria. It might prove interesting to study the site of binding of pesticide-degrading organisms and of numerous other organisms.

G. Ganesh* and J. S. Venugopal

19, Devasikamani Road, Muthamiz Nagar, Pammal, Pallavaram, Madras 600 075, India.

*For correspondence.

1. Ganesh, G. (1990). Ultrastructure and morphological variations of some strains of *Pseudomonas* sp. with special reference to pesticide biodegradation. M Phil thesis, Mangalore University, India.
2. Ganesh, G., Misra, A. K., Chapelon, C. & Normand, P. (1994). Morphological and molecular characterization of *Frankia* sp. isolates from nodules of *Alnus nepalensis* Don. *Arch Microbiol* **161**, 152–156.
3. Hayat, M. A. (1975). *Positive Staining for Electron Microscopy*. New York: Van Nostrand Reinhold.
4. Venugopal, J. S. & Ganesh, G. (1991). A novel method for the detection of pesticide degrading *Pseudomonas* by interference microscopy. *Sci Cult* **57**, 139–140.