or not the chromosome is a topologically constrained covalently closed circle. The bacterial nucleoid is thought to be folded into domains of about 100 kb which are topologically separate. RNA and protein scaffolding is proposed to maintain this structure. If there are scaffold attachment points which define these domains then sequences in different domains should not be able to contact each other by sliding. Pat Higgins described a test of this hypothesis using Tn1000 resolvase sites introduced via Mu into a non-essential part of the *Salmonella typhimurium* chromosome. He showed that there do not appear to be barriers to distant points on the chromosome meeting each other, which goes against a fixed nucleoid structure.

Another important area of bacterial genetics is recombination and mutagenesis and particularly the way that stress can induce a higher rate of mutation, particularly changes which have an adaptive advantage. The mechanism by which chi sites alter RecD to convert the RecBCD complex from destructive to recombinogenic was an exciting topic described by Frank Stahl. Maurice Fox reported that conjugation can stimulate adaptive mutation, which ties in with the observation reported by Susan Rosenberg that adaptive mutation requires recombination enzymes and supports a model when single-strand invasion primes mutagenic DNA synthesis.

Apart from these aspects of bacterial genetics, most of the meeting focused on gene expression and regulation. There is much interest in the details of repressors and activators, of the contacts RNA polymerase makes with DNA and regulator proteins, and the ways that regulatory proteins modulate DNA structure, including looping. The process of transcription initiation and elongation is also receiving much attention, as is the story of Gre factors which rescue abortive transcription events. Polyadenylation of bacterial mRNA also seems to be accepted now as a result of demon-

stration via PCR with poly-T primers that many messengers are modified in this way. Termination and antitermination are still topical and will soon be understood much more clearly.

However, in trying to bring the genetics of bacteria alive to students it is probably important to emphasize the questions still to be answered which are of importance to society as a whole. The evolution and adaptation of bacteria in soil and clinical contexts and the fundamental understanding of how they evolved and how we can combat them when they cause disease are all still wide open and will draw on the ingenuity of geneticists for many years to come.

Christopher M. Thomas

School of Biological Sciences, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK.

Tel: +44 121 414 5903. Fax: +44 121 414 5925.
e-mail: C.M.Thomas@BHAM.AC.UK

Plasmid-mediated exchange between molecular biologists and ecologists

'Workshop' is a term often used for a scientific meeting with a limited number of participants, but so often these end up as mini conferences – a series of oral presentations with little time for questions or discussion. As a result, there is limited opportunity to air different approaches to a problem or to discuss how different approaches may come together to provide a fuller picture of a complex phenomenon. One of the purposes of the Network on 'Molecular biology and ecology of plasmid-mediated genetic exchange' is to stimulate the interaction between molecular biologists and microbial ecologists. These groups have been

attending some of the same meetings for a number of years, for example the Bacterial Genetics and Ecology (BAGECO) series, but a gulf still remains. A major task of the Network workshop is to stimulate this interaction or break down the communication barrier between the two groups. We therefore dispensed with the presentation of results except through poster presentations and concentrated on discussion on defined topics that would bring together both sides. Introductory talks set the scene for the four sessions. We then split down into six groups of ten (total attendees equalled 60) to discuss our different interests in the topics. Finally we got back together to share ideas.

This style of forced discussion seemed to work well for junior and senior scientists alike. The groups were picked on the basis of my prior knowledge of personalities and interests. Many people who would rarely participate in a large group found themselves able to discuss their work. Our impression was that many barriers were broken down and we hope to see evidence of this in future meetings.

The main outcome of the meeting was an agreement to set up a computer bulletin board to facilitate exchange of information and to set up a plasmid database to act as a resource for accumulating information on well-studied and newly isolated plasmids. European laboratories interested in participating in the use of this resource who do not already belong to the Network should contact the author at the address shown below.

Christopher M. Thomas

School of Biological Sciences, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK.

Tel: +44 121 414 5903. Fax: +44 121 414 5925.
e-mail: C.M.Thomas@BHAM.AC.UK