A protein having similarity with methylmalonyl-CoA mutase is required for the assimilation of methanol and ethanol by *Methylobacterium extorquens* AM1

Loraine M. Smith,^{1,2} Wim G. Meijer,² Lubbert Dijkhuizen² and Pat M. Goodwin^{1†}

Author for correspondence: Pat M. Goodwin. Tel: +44 171 611 8690. Fax: +44 171 611 8363.
e-mail: p.goodwin@wellcome.ac.uk

¹ School of Cell and Molecular Biology, NEScot, Reigate Road, Epsom KT17 3DS, UK

² Department of Microbiology, Groningen Biomolecular Sciences and Biotechnology Institute (GBB), University of Groningen, Kerklaan 30, 9751 NN Haren, The Netherlands

A 4.0 kb region of *Methylobacterium extorquens* AM1 DNA which complements three mutants unable to convert acetyl-CoA to glyoxylate (and therefore defective in the assimilation of methanol and ethanol) has been isolated and sequenced. It contains two ORFs and the 3'-end of a third one. The mutations in all three mutants mapped within the first ORF, which was designated *meaA*; it encodes a protein having similarity with methylmalonyl-CoA mutase. However, methylmalonyl-CoA mutase was measured in extracts of one of the mutants and the specific activity was found to be similar to that in extracts of wild-type cells. Furthermore, although the predicted *meaA* gene product has the proposed cobalamin-binding site, it does not contain a highly conserved sequence (RIARNT) which is present in all known methylmalonyl-CoA mutases; *meaA* may therefore encode a novel vitamin-B₁₂-dependent enzyme. The predicted polypeptide encoded by the second ORF did not have similarity with any known proteins. The partial ORF encoded a protein with similarity with the 3-oxoacyl-[acyl-carrier-protein] reductases; it was not essential for growth on methanol or ethanol.

Keywords: *Methylobacterium*, methanol, mutase, serine pathway, C₁ assimilation

INTRODUCTION

Methylobacterium extorquens AM1 is a pink-pigmented facultative methylotroph which assimilates C₁ compounds by the serine pathway (Fig. 1) (Anthony, 1982). The first step in this pathway involves the condensation of a C₁-tetrahydrofolate derivative with glycine to form serine, catalysed by serine hydroxymethyltransferase. The serine is then converted to C₃ and C₄ carboxylic acids which can be assimilated into cell material. This involves four key enzymes – serine glyoxylate aminotransferase, hydroxypyruvate reductase, glycerate kinase and an acetyl-CoA-independent phosphoenolpyruvate carboxylase. There must also be a means of regenerating the C₁ acceptor, glycine. Its immediate precursor is glyoxylate and acetate is an intermediate in the recycling pathway. In the so-

called Icl⁺ serine pathway organisms, malate thiokinase, malyl-CoA lyase, isocitrate lyase and some of the tri-carboxylic acid cycle enzymes are required for the conversion of acetate to glyoxylate (Bellion & Hersh, 1972).

In *M. extorquens* AM1 there is evidence that malyl-CoA lyase and malate thiokinase are essential for the assimilation of C₁ compounds (Salem *et al.*, 1973a, 1974; Chistoserdova & Lidstrom, 1994b). However, isocitrate lyase has never been detected in this organism, nor in a number of other serine pathway methylotrophs (Dunstan *et al.*, 1972a; Kortstee, 1981); such organisms are known as Icl[–] serine pathway methylotrophs. Although several different routes for the conversion of acetate to glyoxylate in these organisms have been proposed (Dunstan *et al.*, 1972b; Kortstee, 1981; Shimizu *et al.*, 1984), none of them have been confirmed. Several mutants have been described which are unable to grow on C₁ compounds unless the medium is supplemented with glyoxylate, but the biochemical basis of this phenotype has not been elucidated (Dunstan *et al.*, 1972b; Salem *et al.*, 1973b;

† Present address: The Wellcome Trust, 183 Euston Road, London NW1 2BE, UK.

The GenBank accession number for the nucleotide sequence reported in this paper is U28335.

Fig. 1. Icl^- variant of the serine pathway showing the key enzymes and the route (broken line) for the regeneration of glyoxylate proposed by Shimizu *et al.* (1984). GK, glycerate kinase; GM, glutamate mutase; HPRT, hydroxypyruvate reductase; MA, β -methylaspartase; MCL, maly-CoA lyase; MH, mesaconyl-CoA hydratase; MMC, methylmalonyl-CoA mutase; MML, methylmalyl-CoA lyase; MT, mesaconate thiokinase; MTK, malate thiokinase; PCC, propionyl-CoA carboxylase; PEPC, phosphoenolpyruvate carboxylase; SGAT, serine:glyoxylate aminotransferase; STHM, serine transhydroxymethylase; THF, tetrahydrofolate.

Bolbot & Anthony, 1980a, b; Stone & Goodwin, 1989). Mutants with this phenotype are also unable to grow on ethanol and on other substrates assimilated after conversion to acetate, including 1,2-propanediol, β -hydroxybutyrate and malonate. This suggests that the enzyme(s) defective in these mutants is essential for utilization of these compounds as sole carbon source (Dunstan *et al.*, 1972b; Dunstan & Anthony, 1973; Bolbot & Anthony, 1980a, b).

Recently, Chistoserdova & Lidstrom (1994a) have isolated a mutant defective in serine hydroxymethyltransferase. Surprisingly, it was able to grow on succinate, but not on methanol or ethanol. Furthermore, when the medium was supplemented with glyoxylate it grew on ethanol, but not on methanol. This indicates that serine hydroxymethyltransferase is required for the conversion of acetyl-CoA to glyoxylate, as well as for the fixation of formaldehyde during growth on C_1 compounds. The protein encoded by an ORF (*orf4*) which is found within

a cluster of seven other serine pathway genes also has two functions during growth on C_1 compounds, one being that it is necessary for the conversion of acetyl-CoA to glyoxylate. The deduced amino acid sequence of this protein does not exhibit similarity with any known proteins (Chistoserdova & Lidstrom, 1994b).

Three other regions of the *M. extorquens* AM1 chromosome encode serine pathway genes; one complements mutants defective in glycerate kinase and the other two complement mutants which are specifically blocked in the conversion of acetyl-CoA to glyoxylate (Stone & Goodwin, 1989). In this paper, we describe the isolation and characterization of another mutant of the latter type and demonstrate that it, and two of the previously described mutants, are defective in a protein with similarity with methylmalonyl-CoA mutase.

METHODS

Bacterial strains. The bacterial strains and plasmids used in this study are listed in Table 1.

Growth of organisms. *M. extorquens* AM1 was grown at 30 °C on MacLennan's minimal salts medium (MacLennan *et al.*, 1971). Carbon substrates were added to give the following final concentrations: 0.4% (v/v) for methanol; 0.5% (v/v) for 1,2-propanediol; 5 mM for glyoxylate; 0.2% (v/v) for ethanol; 0.2% (w/v) for other substrates. *Escherichia coli* strains were grown in Luria-Bertani broth at 37 °C (Sambrook *et al.*, 1989). Where appropriate, supplements were added to the medium at the following concentrations: oxytetracycline hydrochloride, 20 $\mu\text{g ml}^{-1}$ for *E. coli* and 12.5 $\mu\text{g ml}^{-1}$ for *M. extorquens* AM1; kanamycin sulphate, 50 $\mu\text{g ml}^{-1}$ for *E. coli*; ampicillin, 50 $\mu\text{g ml}^{-1}$; X-Gal, 20 $\mu\text{g ml}^{-1}$; isopropyl β -D-thiogalactoside, 0.1 mM.

Isolation of mutants. A culture of wild-type *M. extorquens* AM1 growing on methanol medium was harvested at the mid-exponential phase of growth and resuspended at a density of approximately 10^9 cells ml^{-1} in 0.2 M sodium acetate buffer (pH 6.4) containing 1 mg sodium nitrite ml^{-1} and 2 μg chloramphenicol ml^{-1} , which has been reported to enhance mutagenesis (Mishra & Tiwari, 1985). After incubation with shaking for 1 h at 30 °C, the cells were harvested and washed in succinate medium. Expression and penicillin enrichment were done as described by Tatra & Goodwin (1985). Mutants able to grow on succinate but not on methanol or ethanol were isolated by replica plating.

Preparation of cell extracts and enzyme assays. Cell extracts were prepared as described by Tatra & Goodwin (1985). Methanol dehydrogenase (EC 1.1.99.8), methylamine dehydrogenase (EC 1.4.99.3) and formaldehyde dehydrogenase (EC 1.2.99.3) were assayed polarographically (Tatra & Goodwin, 1985; Dawson *et al.*, 1990; Ford *et al.*, 1985). The following enzymes were assayed spectrophotometrically (Shimadzu UV 260 dual beam spectrophotometer) using published methods: formate dehydrogenase (EC 1.2.1.2; Johnson & Quayle, 1964); hydroxypyruvate reductase (EC 1.1.1.81; Goodwin, 1990); glycerate kinase (EC 2.7.1.31; Goodwin, 1990); serine:glyoxylate aminotransferase (EC 2.6.1.45; Goodwin, 1990); acetyl-CoA independent phosphoenolpyruvate carboxylase (Goodwin, 1990). Maly-CoA lyase (EC 4.1.3.24) could not be assayed directly because the substrate, maly-CoA, is not commercially available. The presence of this enzyme was deduced by measuring the apparent malate synthase activity, which is due to the concerted action of maly-CoA lyase and

Table 1. Bacterial strains and plasmids

Strain/plasmid	Relevant properties	Source/reference
Bacteria		
<i>M. extorquens</i> AM1		
Wild-type		Peel & Quayle (1961)
PCT48	Mutants defective in the conversion of acetyl-CoA to glyoxylate	Dunstan <i>et al.</i> (1972b)
PT1005		Stone & Goodwin (1989)
LS1		This study
<i>E. coli</i>		
DSM1607	F ⁻ <i>proA2 recA13 rpsL20</i> (Str ^r) <i>hsdS20</i> (<i>hsdR hsdM</i>)	NCIMB 11865
S17-1	<i>thi pro hsdR recA</i> ; contains the <i>tra</i> region of RP4 integrated in the chromosome	Simon <i>et al.</i> (1983)
CSR603	F ⁻ <i>proA2 recA1</i>	NCIMB 11867
XL1-Blue	<i>endA1 hsdR17</i> ($r_K^+ m_K^+$) <i>supE44 thi-1 recA1</i> (<i>lac</i>) F' <i>proAB lacI^qΔM15</i> Tn10 (Tc ^r)	Bullock <i>et al.</i> (1987)
P678-54	F <i>thr1 ara13 leu-6 azi8 tonA2 lacY1 minA1 ginU44 gal6 minB2 rpsL135 malA1 xyl7 mtl2 thi1</i>	Adler <i>et al.</i> (1967)
DH5α	<i>supE44 lacU169</i> (Φ80 <i>lacΔM15</i>) <i>hsdR17 recA1 endA1 gyrA96 thi1 relA1</i>	Bethesda Research Laboratories
Plasmid/cosmid		
pVK100	IncP1, Tc ^r , Km ^r , <i>cos</i> ; broad-host-range cloning vector	Knauf & Nester (1982)
pLA2917	IncP1, Tc ^r , Km ^r , <i>cos</i> ; broad-host-range cloning vector	Allen & Hanson (1985)
pRK415	IncP1, Tc ^r , <i>lacZ'</i> ; broad-host-range cloning vector	Keen <i>et al.</i> (1988)
pRK2013	Km ^r mobilizing plasmid; ColE1 replicon	Figurski & Helinski (1979)
pBluescriptII KS(+)	Amp ^r phagemid with ColE1 replicon and f1 phage origins of replication	Stratagene Cloning Systems
pTZ18U	Amp ^r <i>lacZ'</i> cloning vector	Bio-Rad
pSS48-1, pSS48-2	Cosmids isolated from a HindIII clone bank of <i>M. extorquens</i> AM1 DNA in pVK100	Stone & Goodwin (1989)
pLS27C	4.0 kb <i>EcoRI</i> fragment of pSS48-1 in pRK415	This study
pLS271C	1.0 kb <i>PstI</i> fragment of pLS27C in pLA2917	
pLS273C	2.2 kb <i>PstI</i> - <i>EcoRI</i> fragment of pLS27C in pRK415	
pLS274C	1.0 kb <i>EcoRI</i> - <i>PstI</i> fragment of pLS27C in pRK415	
pLS27	4.0 kb <i>EcoRI</i> fragment of pSS48-1 in pBluescript KS(+)	
pLS271	1.0 kb <i>PstI</i> fragment of pLS27 in pBluescript KS(+)	
pLS272	1.2 kb <i>EcoRI</i> - <i>BglII</i> fragment of pLS27 in pBluescript KS(+)	
pLS273	2.2 kb <i>PstI</i> - <i>EcoRI</i> fragment of pLS27 in pTZ18U	

malyl-CoA hydrolase (Stone & Goodwin, 1989). Methylmalonyl-CoA mutase (EC 5.4.99.2) was measured by the method of Zagalak *et al.* (1974). Enzyme assays were repeated using at least two independent cultures. Protein concentrations were determined by the method of Lowry using BSA fraction V as the standard.

Complementation analysis. Recombinant cosmids were introduced into *M. extorquens* AM1 mutants using either biparental matings (where *E. coli* S17-1 carrying a recombinant cosmid was the donor) or triparental matings (where other *E. coli* hosts carrying a recombinant cosmid were the donors and the mobilizing plasmid pRK2013 was supplied by *E. coli* DSM1607). Matings were done using a modification of the method of Fulton *et al.* (1984). Culture of the recipient (0.1 ml; 10⁸ cells) was spread onto a nutrient agar plate. The plate was dried and then 20 µl culture containing the donor (2 × 10⁷ cells), mixed with an equal number of mobilizer cells if appropriate, was dropped onto the lawn of recipient. After incubation overnight at 30 °C, the plates were replicated onto medium containing (i) succinate plus tetracycline plus carbenicillin (to select for cosmid transfer) and (ii) methanol plus tetracycline (to select for complementation). Carbenicillin (100 µg ml⁻¹) was

added to the succinate medium to inhibit the growth of *E. coli* which occurred due to carry-over of nutrients from the mating plates. To check that complementation, rather than recombinational rescue, was occurring, single colonies which had been selected for transfer of the cosmid were removed and tested for growth on methanol plus tetracycline. Demonstration that all of these colonies grew on methanol indicated that there was complementation *in trans*.

Plasmid isolation. A modification of the alkaline lysis method of Birnboim & Doly (1979) was used for small-scale preparation of plasmid and cosmid DNA (Sambrook *et al.*, 1989). Large-scale isolation of plasmid DNA for sequencing was done as described in the Promega Protocols and Applications Guide.

Subcloning. Subcloning and agarose gel electrophoresis were done as described by Sambrook *et al.* (1989). Enzymes for restriction digestion and DNA manipulations were obtained from Sigma or Northumbria Biologicals and used according to the manufacturers' instructions. A Prep-A-Gene DNA purification kit (Bio-Rad) was used to purify DNA fragments isolated from agarose gels. Recombinant plasmids were introduced into competent *E. coli* cells by transformation as described by Sambrook *et al.* (1989).

DNA-DNA hybridization. Chromosomal DNA was isolated from *M. extorquens* AM1 using the method of Fulton *et al.* (1984). The DNA was digested with an appropriate restriction enzyme and the resulting fragments were separated by agarose gel electrophoresis and then blotted onto a Hybond-N nylon membrane (Amersham), using the Hybaid Vacu-aid according to the manufacturer's instructions.

DNA probes were prepared by random incorporation of digoxigenin-labelled deoxyuridine triphosphate (Boehringer), according to the manufacturer's instructions. Hybridization was done at 68 °C in a Hybaid incubator as described by Sambrook *et al.* (1989), using stringent washing conditions, and the hybridized probe was detected using the DIG nucleic acid detection kit (Boehringer).

Nucleotide sequencing. Plasmid pLS27 was digested with *Pst*I and the resulting 1.0 kb fragment was cloned into pBluescript KS(+), generating plasmid pLS271. The 1.1 kb *Eco*RI-*Bgl*II fragment of pLS27 was ligated into pBluescript KS(+) which had been digested with *Eco*RI and *Bam*HI, giving plasmid pLS272. The 2.2 kb *Pst*I-*Eco*RI fragment of pLS27 was subcloned into pTZ18U which had been digested with the same enzymes and a nested set of unidirectional deletions of the resulting plasmid, pLS273, was made. This was done as described by Henikoff (1984), except that mung bean nuclease was used instead of S1 nuclease. DNA fragments were sequenced by the dideoxy chain-termination method (Sanger *et al.*, 1977). Primers were either obtained commercially or were custom synthesized.

Sequencing was either done with T7 DNA polymerase and [³⁵S]dATPαS using the Sequenase kit (US Biochemical) or by using a Pharmacia Automated Laser Fluorescence DNA sequencer.

Sequence data were compiled and analysed using the GCG programs (Devereux *et al.*, 1984) and the programs supplied in the PC/GENE software packages (Intelligenetics).

Preparation and labelling of *E. coli* minicells and analysis of labelled peptides. The *E. coli* minicell-producing strain P687-54 was transformed with pLS27 and with pBluescript KS(+). Minicells were isolated using a modified version of the method of Clark-Curtiss & Curtiss (1983) as described by Eggink *et al.* (1988). The protein products were labelled with [³⁵S]methionine, analysed by SDS-PAGE and visualized by fluorography (Eggink *et al.*, 1988).

RESULTS AND DISCUSSION

Isolation and characterization of mutants

Following nitrous acid mutagenesis, four mutants were isolated which had growth properties typical of mutants unable to convert acetyl-CoA to glyoxylate, i.e. were unable to grow on methanol or ethanol unless the medium was supplemented with glyoxylate. We have designated this phenotype Mea (methanol and ethanol assimilation deficient). Revertants were obtained at a frequency of approximately 10⁻⁹. As expected, the mutants were also unable to utilize methylamine, formate, malonate or β-hydroxybutyrate as sole carbon source, but could grow on oxalate, which is assimilated by metabolism to glyoxylate and formate and then converted to phosphoglycerate by the appropriate enzymes of the serine pathway (Blackmore & Quayle, 1970). Surprisingly, pyruvate, lactate and 1,2-propanediol, which are also thought to be assimilated by

a route involving metabolism to acetyl-CoA and its subsequent conversion to glyoxylate (Salem *et al.*, 1973b; Bolbot & Anthony, 1980a, b), supported growth of the new isolates and also of two previously described mutants, PCT48 and PT1005. However, all of the mutants grew much more slowly than the wild-type on 1,2-propanediol and this may account for the failure of Bolbot & Anthony (1980b) to observe growth of PCT48 on this substrate. These results suggest that, if acetate is an intermediate in the assimilation of pyruvate, lactate and 1,2-propanediol, it must be metabolized by a route which does not involve conversion to glyoxylate.

Biochemical and complementation analysis indicated that the four new mutants were identical so the results for only one – LS1 – are given. It was grown on medium containing succinate plus methanol plus methylamine, harvested, and incubated overnight in medium containing methanol plus methylamine to induce the C₁-metabolizing enzymes. Crude extracts were then assayed for the known key enzymes of the serine pathway. Serine:glyoxylate aminotransferase, hydroxypyruvate reductase, glyceralate kinase and the acetyl-CoA-independent phosphoenolpyruvate carboxylase were all detected, as were the C₁-oxidizing enzymes methanol dehydrogenase, methylamine dehydrogenase, formaldehyde dehydrogenase and formate dehydrogenase. The presence of malyl-CoA lyase was deduced by demonstrating an apparent malate synthase activity, which is due to the concerted action of malyl-CoA lyase and malyl-CoA hydrolase. We did not assay for serine hydroxymethyltransferase, but this enzyme must be present since LS1 grew on oxalate and on methanol in the presence of glyoxylate.

Complementation analysis

Stone & Goodwin (1989) isolated two overlapping cosmids (pSS48-1 and pSS48-2) from a *Hind*III genomic library of *M. extorquens* AM1 DNA constructed in the broad-host-range mobilizable cosmid pVK100, and demonstrated that they complemented two mutants, PCT48 and PT1005, which were unable to convert acetyl-CoA to glyoxylate; a third mutant with the same phenotype (Cou-4) was not complemented. pSS48-1 and pSS48-2 also complemented mutant LS1.

pLS27C, a subclone of pSS48-1 which contained a 4.0 kb *Eco*RI fragment of pSS48-1, also complemented all of the mutants but subclones containing smaller DNA fragments derived from pLS27C did not (Fig. 2). However, when one of these (pLS274C) was introduced into PT1005, approximately 1 % of the transconjugants which had received the cosmid grew on methanol, indicating that recombinational rescue had occurred. Recombinational rescue also occurred when pLS273C was introduced into LS1.

Mutant PCT48 contains a chromosomal deletion

In common with Dunstan *et al.* (1972b), we were unable to isolate revertants of this mutant and we therefore investigated their suggestion that PCT48 is a deletion

Fig. 2. (a) Restriction map of the 26 kb *Hind*III insert of cosmid pSS48-1. The hatched box indicates the position of the deletion in PCT48. (b) Complementation analysis of the Mea mutants using subclones of pSS48-1. C, Complementation; R, recombination; —, no complementation or recombination. The positions and directions of transcription of *meaA*, *orfB* and *orfC* are indicated by arrows.

Fig. 3. DNA-DNA hybridization of genomic DNA from wild-type *M. extorquens* AM1 and mutant PCT48. (a) Probed with the 4.0 kb *Eco*RI fragment from pLS27. Lanes: 1, *Eco*RI digest of genomic DNA from PCT48; 2, *Eco*RI digest of genomic DNA from wild-type; 3, *Pst*I digest of lambda DNA; 4, *Eco*RI-*Hind*III digest of lambda DNA. (b) Probed with the 1.0 kb *Pst*I fragment from pLS271C. Lanes: 1, *Pst*I digest of genomic DNA from PCT48; 2, *Pst*I digest of genomic DNA from wild-type; 3, *Pst*I digest of pLS271C; 4, *Pst*I digest of lambda DNA; 5, *Eco*RI-*Hind*III digest of lambda DNA. The *Pst*I fragment bound to pLA2917 (data not shown) and this accounts for the bands of 21 kb (vector DNA) and 6.0 kb (degraded vector DNA).

mutant. Chromosomal DNA isolated from the wild-type strain and from PCT48 was digested with *Eco*RI. The resulting fragments were separated by gel electrophoresis

The predicted amino acid sequences of the ORFs were compared with entries in the protein database at the National Center for Biotechnology Information (NCBI) using the program BLAST (Altschul *et al.*, 1990). The *meaA* gene product had a high degree of similarity with the methylmalonyl-CoA mutases, which are adenosylcobalamin-dependent enzymes catalysing the interconversion of methylmalonyl-CoA and succinyl-CoA (Fig. 5). The enzymes from *Propionibacterium shermanii*, *Streptomyces cinnamonensis* and *Porphyromonas gingivalis* are heterodimers consisting of a large subunit (approximately 79 kDa) and a small subunit (approximately 65 kDa) (Marsh *et al.*, 1989; Birch *et al.*, 1993). In contrast, the mouse and human enzymes comprise two identical subunits of 82 kDa (Ledley *et al.*, 1988; Jansen *et al.*, 1989; Wilkemeyer *et al.*, 1990). The *sbm* gene of *E. coli* encodes

MeaA	MSAQAQSAEVAEKRD-----KPWIIRTYAGHSTA	27
Mouse	MLRAKNQLFLLSPHYLKQNLIPSA--RWKRLHQQQPLHPEWAVLAKKQKLG--KNPEDLIWHTPEGISIKPLYSRAD--TLDLPEELPGVKPPTRGPPYPTMYTTPRWITRQYAGFSTV	114
P. sh	MSTLPRF--DSVDLGNAP-----VPADAARRFEELAAGAG-----TGEAWETARQIPVGTLLFNEEDVYKMDWLDYAGIPFPVHGPAATMYAFRRPWTIRQYAGFSTA	95
S. ci	MRIPF--DDIELGAGG-----GFSGSAEQWRAAVKESVG--KSESDLLWETPEGIAVKPLTYGADVEGLDPLETYPGVAPFLRGPPYPTMYTTPRWITRQYAGFSTA	116
Human	MLRAKNQLFLLSPHYLKQNLIPSA--RWKRLHQQQPLHPEWAVLAKKQKLG--KNPEDLIWHTPEGISIKPLYSRAD--TLDLPEELPGVKPPTRGPPYPTMYTTPRWITRQYAGFSTV	90
P. gi	MKNPY--KDIDIKSAG-----FVAKDATHWAEKIVAD-----WRTEQIMVKPLTKDDLEGMHLDVYSGLPPLRGPYSYGMYPMRPWTIRQYAGFSTA	84
E. co	MSNVQEQWLANKLSRREKTVDSLVHQTAEGIAIKPLYTEADLDNLEVTGTLPGLPFVVRGPRATMYTAPQWTIRQYAGFSTA	147
MeaA	AESNKLYRGNLAKGQGTGLSVAFDLPTQGTGYDPDELARGEVGVKGVSAHLGDMRALFDQIPLAQNMSTMTINATAPWLLSLYLVAEEQGAFLAALQGTQNDIIKEYLSRGTYVFPFA	234
Mouse	EESNKFYKDNKAGQGLSVAFDLATHRGVSDNPRVRGVDVGMAGVAIDTVEDTKILPDGIPLEKMSVMTMNGAVIPVLATFIVTGEQGVKPEKLTGTIQNDILKEFVMTYIYFPPE	215
P. sh	KESNAFYRRNLAAQKGLSVAFDLPTHRGVSDNPRVRGVDVGMAGVAIDSIYDMRQLFDGIPLDKMTVMTMNGAVIPVLALYVVAEEQGVKPEKLTGTIQNDILKEFVMTYIYFPPE	218
S. ci	EESNAFYRRNLAAQKGLSVAFDLPTHRGVSDNPRVRGVDVGMAGVAIDSIYDMRQLFDGIPLDKMTVMTMNGAVIPVLALYVVAEEQGVKPEKLTGTIQNDILKEFVMTYIYFPPE	210
Human	EESNAFYRRNLAAQKGLSVAFDLPTHRGVSDNPRVRGVDVGMAGVAIDSIYDMRQLFDGIPLDKMTVMTMNGAVIPVLALYVVAEEQGVKPEKLTGTIQNDILKEFVMTYIYFPPE	204
P. gi	EESNAFYRRNLAAQKGLSVAFDLPTHRGVSDNPRVRGVDVGMAGVAIDSIYDMRQLFDGIPLDKMTVMTMNGAVIPVLALYVVAEEQGVKPEKLTGTIQNDILKEFVMTYIYFPPE	204
E. co	KESNAFYRRNLAAQKGLSVAFDLATHRGVSDNPRVRGVDVGMAGVAIDTVEDTKILPDGIPLEKMSVMTMNGAVIPVLATFIVTGEQGVKPEKLTGTIQNDILKEFVMTYIYFPPE	204
MeaA	PSLRITKDVILFTTKNVPKNMNVCSYHLQEGATPVQELSYALAIAlAVLDTVRDPDFDEASFSDFSRISFFVAGMRFTVEICKMRAFAELWDEIAQERYGITDAKKRIFRYGVQ	267
Mouse	PSMKIADIIFYTAQAMPKFNPSISISGYHMQEAGTDAILLAYTADGLEYSRTGLQAG--LTIDEFAPRLSFFWIGMNFYMEIAKMRAGRRLWAHLIEKMPQPKNSKSLLRHACQ	350
P. sh	PSMKIISIEIPAYTSAMMPKFNPSISISGYHMQEAGTDAILLAYTADGLEYSRTGLQAG--LTIDEFAPRLSFFWIGMNFYMEIAKMRAGRRLWAHLIEKMPQPKNSKSLLRHACQ	330
S. ci	PSMKIISIEIPAYTSAMMPKFNPSISISGYHMQEAGTDAILLAYTADGLEYSRTGLQAG--LTIDEFAPRLSFFWIGMNFYMEIAKMRAGRRLWAHLIEKMPQPKNSKSLLRHACQ	333
Human	PSMKIISIEIPAYTSAMMPKFNPSISISGYHMQEAGTDAILLAYTADGLEYSRTGLQAG--LTIDEFAPRLSFFWIGMNFYMEIAKMRAGRRLWAHLIEKMPQPKNSKSLLRHACQ	352
P. gi	PSMKIADIIFYTAQAMPKFNPSISISGYHMQEAGTDAILLAYTADGLEYSRTGLQAG--LTIDEFAPRLSFFWIGMNFYMEIAKMRAGRRLWAHLIEKMPQPKNSKSLLRHACQ	325
E. co	PSMKIADIIFYTAQAMPKFNPSISISGYHMQEAGTDAILLAYTADGLEYSRTGLQAG--LTIDEFAPRLSFFWIGMNFYMEIAKMRAGRRLWAHLIEKMPQPKNSKSLLRHACQ	319
MeaA	VNSLGLTEQPENNVHRIILEMLAVTLTKRARARVOLPAWNEALGLRPWPDQWMMRQOILAFETDLEYYDDIFDGSFTVIEARVEALKEOTRAELTRIAEIGGAVTAVEAGEKRALV	387
Mouse	TSGWSLTDQPPYNNVIRTAIEAAMAVFG--GTQSLHNSFDEALGLPTFVSARITARIITIIQESGIPKVDADPWGSSYMECLTNDVYEAALKLIYEVEEMGGMAKAEVGPKEKRIE	467
P. sh	TSGWSLTDQPPYNNVIRTAIEAAMAVFG--GTQSLHNSFDEALGLPTFVSARITARIITIIQESGIPKVDADPWGSSYMECLTNDVYEAALKLIYEVEEMGGMAKAEVGPKEKRIE	447
S. ci	TSGWSLTDQPPYNNVIRTAIEAAMAVFG--GTQSLHNSFDEALGLPTFVSARITARIITIIQESGIPKVDADPWGSSYMECLTNDVYEAALKLIYEVEEMGGMAKAEVGPKEKRIE	450
Human	TSGWSLTDQPPYNNVIRTAIEAAMAVFG--GTQSLHNSFDEALGLPTFVSARITARIITIIQESGIPKVDADPWGSSYMECLTNDVYEAALKLIYEVEEMGGMAKAEVGPKEKRIE	469
P. gi	TSGWSLTDQPPYNNVIRTAIEAAMAVFG--GTQSLHNSFDEALGLPTFVSARITARIITIIQESGIPKVDADPWGSSYMECLTNDVYEAALKLIYEVEEMGGMAKAEVGPKEKRIE	442
E. co	TSGWSLTDQPPYNNVIRTAIEAALGL--GTQSLHNSFDEALGLPTFVSARITARIITIIQESGIPKVDADPWGSSYMECLTNDVYEAALKLIYEVEEMGGMAKAEVGPKEKRIE	436
MeaA	ESNARRISAIKGEQIVGVNKKWQGG--EPSPLTAGDGAIFTVSETVMEAEATRIEWRKSRDEBAVGOALADLEQAARSQ-----ANIMPPSIAAAKAGVTIGEWQRLREVPGGEYRA	499
Mouse	EAARARQARIDSGSEVIVGVNKKYHLEKEDSVHLLAID-----IISLKKQIEKLIKISSRDOALABOCSALQCAASG-----DGNILALAVDASRARCTVPEITDAFKKVGHEKA	576
P. sh	EAARARQARIDSGSEVIVGVNKKYHLEKEDSVHLLAID-----IISLKKQIEKLIKISSRDOALABOCSALQCAASG-----DGNILALAVDASRARCTVPEITDAFKKVGHEKA	561
S. ci	EAARARQARIDSGSEVIVGVNKKYHLEKEDSVHLLAID-----IISLKKQIEKLIKISSRDOALABOCSALQCAASG-----DGNILALAVDASRARCTVPEITDAFKKVGHEKA	564
Human	EAARARQARIDSGSEVIVGVNKKYHLEKEDSVHLLAID-----IISLKKQIEKLIKISSRDOALABOCSALQCAASG-----DGNILALAVDASRARCTVPEITDAFKKVGHEKA	578
P. gi	EAARARQARIDSGSEVIVGVNKKYHLEKEDSVHLLAID-----IISLKKQIEKLIKISSRDOALABOCSALQCAASG-----DGNILALAVDASRARCTVPEITDAFKKVGHEKA	551
E. co	EASAREQSLIDQKQVIVGVNKKYHLEKEDSVHLLAID-----IISLKKQIEKLIKISSRDOALABOCSALQCAASG-----DGNILALAVDASRARCTVPEITDAFKKVGHEKA	544
MeaA	P---TGVTLTQTVTSGA-----AEDARLLIAD--LGERLGETPRLVVGKPGILDGHSNQAQIALRLARDVGFVDVYDGIQTPTPEIVAKAKERGAHVIGLVLGSGSHVPLVREVAKLREAGL	610
Mouse	NDRMVSGAYRQEGESK-----EITSAIKRVKFMERGRRLGLLVAKMGQGGEDRGAKVIATGFADLGFDDVIGPLFQTPREVAHDAVDADVHAGVGTSLAAGHKTLPVPELIELTALGR	692
P. sh	NDRMVSGAYRQEGESK-----EITSAIKRVKFMERGRRLGLLVAKMGQGGEDRGAKVIATGFADLGFDDVIGPLFQTPREVAHDAVDADVHAGVGTSLAAGHKTLPVPELIELTALGR	677
S. ci	NDRMVSGAYRQEGESK-----EITSAIKRVKFMERGRRLGLLVAKMGQGGEDRGAKVIATGFADLGFDDVIGPLFQTPREVAHDAVDADVHAGVGTSLAAGHKTLPVPELIELTALGR	680
Human	NDRMVSGAYRQEGESK-----EITSAIKRVKFMERGRRLGLLVAKMGQGGEDRGAKVIATGFADLGFDDVIGPLFQTPREVAHDAVDADVHAGVGTSLAAGHKTLPVPELIELTALGR	694
P. gi	NDRMVSGAYRQEGESK-----EITSAIKRVKFMERGRRLGLLVAKMGQGGEDRGAKVIATGFADLGFDDVIGPLFQTPREVAHDAVDADVHAGVGTSLAAGHKTLPVPELIELTALGR	667
E. co	PSQCVTVGIAQSYHQSEKSAEADTAVTQEQFLADNGRRPRILLAKMGQGGEDRGAKVIATGFADLGFDDVIGPLFQTPREVAHDAVDADVHAGVGTSLAAGHKTLPVPELIELTALGR	664
MeaA	DHVPVVVGGIISTEDELVLKMMQVAVTAYTPKDYELDKIMVGLAKVVERALDKRAADRADTEAGVPGAPKRNESGAQVF	688
Mouse	PDILVVCQGVIPQDYEFLYEVGVSNVFGPG--TRIPRAAQVQLDDIEKCLAEKQSV	748
P. sh	PDILVVCQGVIPQDYEFLYEVGVSNVFGPG--TRIPRAAQVQLDDIEKCLAEKQSV	728
S. ci	PDILVVCQGVIPQDYEFLYEVGVSNVFGPG--TRIPRAAQVQLDDIEKCLAEKQSV	733
Human	PDILVVCQGVIPQDYEFLYEVGVSNVFGPG--TRIPRAAQVQLDDIEKCLAEKQSV	750
P. gi	PDILVVCQGVIPQDYEFLYEVGVSNVFGPG--TRIPRAAQVQLDDIEKCLAEKQSV	715
E. co	EDICVVCQGVIPQDYEFLYEVGVSNVFGPG--TRIPRAAQVQLDDIEKCLAEKQSV	714

Fig. 5. Alignment of the deduced amino acid sequence of MeaA with sequences of members of the methylmalonyl-CoA mutase family from other sources. P. sh, *Prop. shermanii* large subunit; S. ci, *S. cinnamomensis* large subunit; P. gi, *Porp. gingivalis* large subunit; E. co; *E. coli sbm* gene product. Identical residues are indicated by an asterisk, and conserved substitutions, according to the scheme PAGST, QNED, ILVM, HKR, YFW, C, are indicated by a dot. Numbers refer to nucleotide residues. The amino acid residues shown in bold and underlined represent the proposed vitamin-B₁₂-binding site and those in italics and underlined represent the conserved sequence in all methylmalonyl-CoA mutases.

a polypeptide of 78 kDa which also belongs to this family of proteins (Roy & Leadlay, 1992). Individual sequence alignments indicated that MeaA has 56–57% similarity with the large subunits of the bacterial enzymes, the mouse and human enzymes and the *sbm* gene product. The identities were 37–37.5% in the case of the *S. cinnamomensis* and *Porp. gingivalis* large subunits and the *sbm* gene product, and 34% with the mouse, human and *Prop. shermanii* large subunits. In contrast, there was only about 46% similarity and 24% identity with the small subunits of the bacterial enzymes.

Three short, highly conserved regions are present in a number of cobalamin-dependent enzymes and it has been proposed that they are involved in cobalamin binding (Marsh & Holloway, 1992; Crane *et al.*, 1992; Drennan *et al.*, 1994). These sequences (DXHXXG, SXL and GX₈GGX₁₄G) also occur in MeaA (Fig. 5). There is another highly conserved sequence (RIARNT) in all of the methylmalonyl-CoA mutases sequenced thus far, but this is not present in *meaA* (Fig. 5).

The deduced amino acid sequence of the polypeptide encoded by *orfB* did not have significant sequence similarity with any known protein. The predicted amino acid sequence encoded by the partial ORF of *orfC* had homology with the 3-oxoacyl-[acyl-carrier-protein] reductases of *Brassica napus* (63% similarity, 41.5% identity), *Arabidopsis thaliana* (64% similarity, 39% identity) (Slabas *et al.*, 1992) and *E. coli* (62% similarity, 37% identity) (Cronan & Rawlings, 1992). This enzyme, which is a member of the short-chain alcohol dehydrogenase family, catalyses the first reduction step in fatty acid biosynthesis.

Translation products of *meaA*

The 4.0 kb fragment cloned in pLS27 was expressed in *E. coli* minicells and the resulting products were analysed using denaturing 8% and 12.5% (w/v) SDS-polyacrylamide gels, which together would have resolved both the polypeptides predicted to be expressed from this fragment. Three polypeptides, with apparent molecular

Fig. 6. SDS-polyacrylamide gel (8%) showing polypeptides synthesized by *E. coli* minicells containing a recombinant plasmid with the 4.0 kb *EcoRI* insert of pLS27 (lane 1) cloned in pBluescript. The molecular masses of marker polypeptides are shown on the right-hand side. The arrows on the left-hand side indicate polypeptides which are synthesized by the recombinant plasmid, but not the control, pBlue (lane 2).

masses of 87, 80 and 64 kDa, were present in minicells expressing this fragment, but not in controls (Fig. 6). The first is somewhat larger than the size of the predicted gene product of *meaA*, but estimates of polypeptide size from SDS-PAGE are often inaccurate. Degradation of large proteins often occurs in minicell expression systems (Eggink *et al.*, 1988) and this presumably accounts for the presence of the 80 and 64 kDa polypeptides. We did not observe any small polypeptides of the size expected of the *orfB* gene product, which is predicted to be 25.5 kDa.

Mutant PCT48 has methylmalonyl-CoA mutase activity

Methylmalonyl-CoA mutase was assayed in cell extracts of the wild-type and mutant PCT48 grown and induced as described in Methods. The specific activity of this enzyme was similar in the wild-type and in mutant PCT48 which contains a deletion in *meaA*, the value being 280 nmol min⁻¹ (mg protein)⁻¹ ± 15%. This does not preclude the possibility that *meaA* encodes an isoenzyme of methylmalonyl-CoA mutase which is required specifically for growth on methanol, although if this is the case it is surprising that the specific activities in the mutant and wild-type are similar. The role of the highly conserved sequence present in all of the methylmalonyl-CoA mutases but not in *MeaA* is unknown, but it presumably has functional significance. Thus the possibility that *meaA* codes for a novel cobalamin-binding protein needs to be explored.

Our results support the suggestion made by Shimizu *et al.* (1984) that vitamin-B₁₂-dependent enzymes are involved in the assimilation of methanol and ethanol. These authors proposed that two adenosylcobalamin-dependent enzymes, methylmalonyl-CoA mutase and glutamate mutase, were required for the conversion of acetyl-CoA to glyoxylate and that β -methylaspartate, mesaconyl-CoA, β -methylmalyl-CoA, propionyl-CoA and methylmalonyl-CoA were intermediates (Fig. 1). It is unlikely that this pathway is correct since serine hydroxymethyltransferase and an ORF encoding a polypeptide of unknown function are essential for the conversion of acetyl-CoA to glyoxylate (Chistoserdova & Lidstrom, 1994a, b), although the possibility that these proteins are required to generate an inducer of the genes involved in this part of the serine pathway cannot be excluded. We have now shown that the *MeaA* protein is also needed, and further work is underway to determine whether it is a mutase with an unusual substrate specificity or whether it has some other function.

ACKNOWLEDGEMENTS

We thank the SERC for a PhD studentship to L.S. and NESCOT for providing facilities and assistance for this work. We are also very grateful to Dr Peter Leadlay, Cambridge, for advice on the methylmalonyl-CoA mutase assay and for the gift of coupling enzymes for the assay; and to Dr Chris Thurston, King's College London, for providing advice and facilities for some of the sequencing work.

REFERENCES

- Adler, H. I., Fisher, W. D., Cohen, A. & Hardigree, A. A. (1967). Minutiae *Escherichia coli* cells deficient in DNA. *Proc Natl Acad Sci USA* **57**, 321–326.
- Allen, L. N. & Hanson, R. S. (1985). Construction of broad host range cosmid cloning vectors: identification of genes necessary for growth of *Methylobacterium organophilum* on methanol. *J Bacteriol* **161**, 955–962.
- Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. (1990). Basic local alignment search tool. *J Mol Biol* **215**, 403–410.
- Anthony, C. (1982). *The Biochemistry of Methylophilus*. London: Academic Press.
- Bellion, E. & Hersh, L. B. (1972). Methylamine metabolism in a *Pseudomonas* species. *Arch Biochem Biophys* **153**, 368–374.
- Birch, A., Leiser, A. & Robinson, J. A. (1993). Cloning, sequencing and expression of the gene encoding methylmalonyl CoA mutase from *Streptomyces cinnamonensis*. *J Bacteriol* **175**, 3511–3519.
- Birnboim, H. C. & Doly, J. (1979). A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res* **7**, 1513–1523.
- Blackmore, M. A. & Quayle, J. R. (1970). Microbial growth on oxalate by a route not involving glyoxylate carboxylase. *Biochem J* **118**, 53–59.
- Bolbot, J. A. & Anthony, C. (1980a). The metabolism of pyruvate by the facultative methylotroph *Pseudomonas* AM1. *J Gen Microbiol* **120**, 233–244.
- Bolbot, J. A. & Anthony, C. (1980b). The metabolism of 1,2-propanediol by the facultative methylotroph *Pseudomonas* AM1. *J Gen Microbiol* **120**, 245–254.

- Bullock, W. O., Fernandez, J. M. & Short, J. M. (1987). XL-1 Blue: a high efficiency plasmid transforming *recA* *Escherichia coli* strain with β -galactosidase selection. *BioTechniques* **5**, 376–379.
- Chistoserdova, L. V. & Lidstrom, M. E. (1994a). Genetics of the serine cycle in *Methylobacterium extorquens* AM1: cloning, sequence, mutation and physiological effect of *glyA*, the gene for serine hydroxymethyltransferase. *J Bacteriol* **176**, 6759–6763.
- Chistoserdova, L. V. & Lidstrom, M. E. (1994b). Genetics of the serine cycle in *Methylobacterium extorquens* AM1: identification, sequence and mutation of three new genes involved in C_1 assimilation, *orf4*, *mtkA* and *mtkB*. *J Bacteriol* **176**, 7398–7404.
- Clark-Curtiss, J. E. & Curtiss, E. I. (1983). Analysis of recombinant DNA using *Escherichia coli* minicells. *Methods Enzymol* **101**, 347–362.
- Crane, A. M., Jansen, R., Andrews, E. R. & Ledley, F. D. (1992). Cloning and expression of a mutant methylmalonyl CoA mutase with altered cobalamin affinity that causes mut-methylmalonic aciduria. *J Clin Invest* **89**, 385–391.
- Cronan, J. E. & Rawlings, M. (1992). The gene encoding *Escherichia coli* acyl carrier protein lies within a cluster of fatty acid biosynthesis genes. *J Biol Chem* **267**, 5751–5754.
- Dawson, A., Southgate, G. & Goodwin, P. M. (1990). Regulation of methanol and methylamine dehydrogenases in *Methylophilus methylotrophus*. *FEMS Microbiol Lett* **68**, 93–96.
- Devereux, J., Haeberli, P. & Smithies, O. (1984). A comprehensive set of sequence analysis programs for the VAX. *Nucleic Acids Res* **12**, 387–395.
- Drennan, C. L., Matthews, R. G. & Ludwig, M. L. (1994). Cobalamin-dependent methionine synthase: the structure of a methylcobalamin-binding fragment and implications for other B_{12} -dependent enzymes. *Curr Opin Struct Biol* **4**, 919–929.
- Dunstan, P. M. & Anthony, C. (1973). Microbial metabolism of C_1 and C_2 compounds. The role of acetate during growth of *Pseudomonas* AM1 on C_1 compounds, ethanol and β -hydroxybutyrate. *Biochem J* **132**, 797–801.
- Dunstan, P. M., Anthony, C. & Drabble, W. T. (1972a). Microbial metabolism of C_1 and C_2 compounds: the involvement of glycolate in the metabolism of ethanol and of acetate by *Pseudomonas* AM1. *Biochem J* **128**, 99–106.
- Dunstan, P. M., Anthony, C. & Drabble, W. T. (1972b). Microbial metabolism of C_1 and C_2 compounds: the role of glyoxylate, glycolate and acetate in the growth of *Pseudomonas* AM1 on ethanol and on C_1 compounds. *Biochem J* **128**, 107–115.
- Eggink, G., Engel, H., Meijer, W. G., Otten, J., Kingma, J. & Witholt, B. (1988). Alkane utilisation in *Pseudomonas oleovorans*. *J Biol Chem* **263**, 13400–13405.
- Figurski, D. H. & Helinski, D. R. (1979). Replication of an origin containing derivative of plasmid RK2 dependent on a plasmid function provided in trans. *Proc Natl Acad Sci USA* **76**, 1648–1652.
- Ford, S., Page, M. D. & Anthony, C. (1985). The role of a methanol dehydrogenase modifier protein and aldehyde dehydrogenase in the growth of *Pseudomonas* AM1 on 1,2-propanediol. *J Gen Microbiol* **131**, 2173–2182.
- Fulton, G. L., Nunn, D. N. & Lidstrom, M. E. (1984). Molecular cloning of a malyl CoA lyase gene from *Pseudomonas* sp. strain AM1, a facultative methylotroph. *J Bacteriol* **160**, 718–723.
- Goodwin, P. M. (1990). Assay of assimilatory enzymes in crude extracts of some serine pathway methylotrophs. *Methods Enzymol* **188**, 361–365.
- Henikoff, S. (1984). Unidirectional digestion with exonuclease III creates targeted breakpoints for DNA sequencing. *Gene* **28**, 351–359.
- Jansen, R., Kalousek, F., Fenton, W. A., Rosenberg, L. E. & Ledley, F. D. (1989). Cloning of full length methylmalonyl-CoA mutase from a cDNA library using the polymerase chain reaction. *Genomics* **4**, 198–205.
- Johnson, P. A. & Quayle, J. R. (1964). Microbial growth on C_1 compounds. 6. Oxidation of methanol, formaldehyde and formate by methanol grown *Pseudomonas* AM1. *Biochem J* **93**, 281–290.
- Keen, N. T., Tamaki, S., Kobayashi, D. & Trollinger, D. (1988). Improved broad-host-range plasmids for DNA cloning in Gram negative bacteria. *Gene* **70**, 191–197.
- Knauf, V. C. & Nester, E. W. (1982). Wide host range cloning vectors: a cosmid clone bank of an *Agrobacterium* Ti plasmid. *Plasmid* **8**, 45–54.
- Kortstee, G. J. J. (1981). The second part of the icl^- serine pathway. In *Microbial Growth on C_1 Compounds*, pp. 211–219. Edited by H. Dalton. London: Heyden & Son.
- Ledley, F. D., Lumetta, M., Nguyen, P. N., Kohlhouse, J. F. & Allen, R. H. (1988). Molecular cloning of L-methylmalonyl-CoA mutase: gene transfer and analysis of *mut* cell lines. *Proc Natl Acad Sci USA* **85**, 3518–3521.
- MacLennan, D. G., Ousby, J. C., Vasey, R. B. & Cotton, N. T. (1971). The influence of dissolved oxygen on *Pseudomonas* AM1 grown on methanol in continuous culture. *J Gen Microbiol* **69**, 395–404.
- Marsh, E. N. G. & Holloway, D. E. (1992). Cloning and sequencing of glutamate mutase component S from *Clostridium tetanomorphum*: homologies with other cobalamin-dependent enzymes. *FEBS Lett* **310**, 167–170.
- Marsh, E. N., Mckie, N., Davis, N. K. & Leadlay, P. F. (1989). Cloning and structural characterization of the genes coding for adenosylcobalamin-dependent methylmalonyl CoA mutase from *Propionibacterium shermanii*. *Biochem J* **260**, 345–352.
- Mishra, A. K. & Tiwari, D. N. (1985). Enhancement of NTG mutagenesis by chloramphenicol in *Gloeotrichia ghosei*. *Mol & Gen Genet* **201**, 351–352.
- Peel, D. & Quayle, J. R. (1961). Microbial growth on C_1 compounds. 1. Isolation and characterisation of *Pseudomonas* AM1. *Biochem J* **81**, 465–469.
- Platt, T. (1986). Transcriptional termination and the regulation of gene expression. *Annu Rev Biochem* **55**, 339–372.
- Roy, I. & Leadlay, P. F. (1992). Physical map location of the new *Escherichia coli* gene *sbm*. *J Bacteriol* **174**, 5763–5764.
- Salem, A. R., Hacking, A. J. & Quayle, J. R. (1973a). Cleavage of malyl-Coenzyme A into acetyl Coenzyme A and glyoxylate by *Pseudomonas* AM1 and other C_1 -unit-utilizing bacteria. *Biochem J* **136**, 89–96.
- Salem, A. R., Wagner, C., Hacking, A. J. & Quayle, J. R. (1973b). The metabolism of lactate and pyruvate by *Pseudomonas* AM1. *J Gen Microbiol* **76**, 375–388.
- Salem, A. R., Hacking, A. J. & Quayle, J. R. (1974). Lack of malyl-CoA lyase in a mutant of *Pseudomonas* AM1. *J Gen Microbiol* **81**, 525–527.
- Sambrook, J., Fritsch, E. F. & Maniatis, T. (1989). *Molecular Cloning: a Laboratory Manual*, 2nd edn. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory.
- Sanger, F., Nicklen, S. & Coulson, A. R. (1977). DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci USA* **74**, 5463–5467.
- Shimizu, S., Ueda, S. & Sato, K. (1984). Physiological role of vitamin B_{12} in a methanol utilising bacterium, *Protaminobacter ruber*. In *Microbial Growth on C_1 Compounds*, pp. 113–117. Edited by R. L.

Crawford & R. S. Hanson. Washington, DC: American Society for Microbiology.

Simon, R., Priefer, U. & Puhler, A. (1983). A broad host range mobilisation system for *in vivo* genetic engineering; transposon mutagenesis in Gram negative bacteria. *Biotechnology* **1**, 784–791.

Slabas, A. R., Chase, D., Nishida, I., Murata, N., Sidebottom, C., Safford, R., Sheldon, P. S., Kekwick, R. G., Hardie, D. G. & Mackintosh, R. W. (1992). Molecular cloning of higher plant 3-oxoacyl-(acyl carrier protein) reductase. Sequence identities with the *nodG* gene product of the nitrogen fixing soil bacterium *Rhizobium meliloti*. *Biochem J* **283**, 321–326.

Stone, S. & Goodwin, P. M. (1989). Characterization and complementation of mutants of *Methylobacterium* AM1 which are defective in C₁ assimilation. *J Gen Microbiol* **135**, 227–235.

Tatra, P. K. & Goodwin, P. M. (1985). Mapping of some genes involved in C₁ metabolism in the facultative methylotroph *Methylobacterium* sp. strain AM1 (*Pseudomonas* AM1). *Arch Microbiol* **143**, 169–177.

Wilkemeyer, M. F., Crane, A. M. & Ledley, F. D. (1990). Primary structure and activity of mouse methylmalonyl CoA mutase. *Biochem J* **271**, 449–455.

Zagalak, B., Retey, J. & Sund, H. (1974). Studies on methylmalonyl CoA mutase from *Propionibacterium shermanii*. *Eur J Biochem* **44**, 529–534.

Received 15 August 1995; revised 24 October 1995; accepted 7 November 1995.