aerotactic response via a methyl-accepting chemotaxis protein (3). This poses the question of whether there is a fundamental reason for diversity in the mechanisms of sensing oxygen. One possible explanation lies in the nature of the sensor domain of DcrA. This periplasmically located domain binds a *c*-type haem which may detect molecular oxygen or redox state (2). In *E. coli* it is highly unlikely that aerotaxis could be mediated by a *c*-type cytochrome since this bacterium represses all *c*-type cytochromes under aerobic growth conditions (4).

James Moir

Centre for Metalloprotein Spectroscopy and Biology, University of East Anglia, Norwich NR4 7TJ, UK.

Tel: +44 1603 592405. Fax: +44 1603 592250.

1. Lindbeck, J. C., Goulbourne, E. A., Johnson, M. S. & Taylor, B. L. (1995). Aerotaxis in *Halobacterium salinarum* is methylation-dependent. *Microbiology* 141, 2945–2953.
2. Fu, R., Wall, J. D. & Voordouw, G. (1994). DcrA, a *c*-type heme-containing methyl-accepting protein from *Desulfovibrio vulgaris* Hildenborough, senses the oxygen concentration or redox potential of the environment. *J Bacteriol* 176, 344–350.
3. Niwano, M. & Taylor, B. L. (1982). Novel sensory adaptation mechanism in bacterial chemotaxis to oxygen

and phospho-transferase substrates. *Proc Natl Acad Sci USA* 79, 11–15.

4. Iobbinivol, C., Crooke, H., Griffiths, L., Grove, J., Hussain, H., Pommier, J., Mejean, V. & Cole, J. A. (1994). A reassessment of the range of *c*-type cytochromes synthesised by *Escherichia coli* K-12. *FEMS Microbiol Lett* 119, 89–94.

Author's reply

Role of DcrA in aerotaxis

The *dcrA* gene product from *Desulfovibrio vulgaris* is a novel and intriguing protein. Early studies were consistent with a role of DcrA as a methylation-dependent sensor of oxygen concentration or of the environmental redox potential (1). Subsequent investigations have revealed that DcrA is a member of a large sensor protein family in *D. vulgaris* (2), but have not yet established a physiological role for DcrA. Methylation of DcrA is increased in response to dithionite and decreased in response to oxygen, as Moir correctly noted. However, ongoing investigations have failed, so far, to demonstrate a defective aerotaxis phenotype in a *dcrA* mutant (B. L. Taylor, I. B. Zhulin, S. Patel, R. Fu & G. Voordouw, unpublished observations).

Therefore, if the DcrA haem moiety binds oxygen, it likely signals to a pathway other than aerotaxis. The possibility that DcrA signals redox information remains a viable and attractive option. Redox taxis has been demonstrated in *Azospirillum brasilense* (3) and in *Escherichia coli* (V. A. Bessalov, I. B. Zhulin & B. L. Taylor, unpublished observations).

Barry L. Taylor

Department of Microbiology and Molecular Genetics, Loma Linda University, Loma Linda, CA 92350, USA.

Tel: +1 909 824 4480. Fax: +1 909 824 4035.
e-mail: bltaylor@ccmail.llu.edu

1. Fu, R., Wall, J. D. & Voordouw, G. (1994). DcrA, a *c*-type heme-containing methyl-accepting protein from *Desulfovibrio vulgaris* Hildenborough, senses the oxygen concentration or redox potential of the environment. *J Bacteriol* 176, 344–350.
2. Deckers, H. M. & Voordouw, G. (1994). Identification of a large family for putative chemoreceptor proteins in an ordered library of the *Desulfovibrio vulgaris* Hildenborough genome. *J Bacteriol* 176, 351–358.
3. Grishanin, R. N., Chalmers, I. I. & Zhulin, I. B. (1991). Behaviour of *Azospirillum brasilense* in a spatial gradient of oxygen and a 'redox' gradient of an artificial electron acceptor. *J Gen Microbiol* 137, 2781–2785.