Identification of staphylococcal and streptococcal causes of bovine mastitis using 16S–23S rRNA spacer regions

Päivi Forsman,¹ Anu Tilsala-Timisjärvi¹ and Tapani Alatossava^{1,2}

Author for correspondence: Päivi Forsman. Tel: +358 8 553 1790. Fax: +358 8 553 1061.
e-mail: forsman@cypress.csc.fi

¹ Department of Biology,
University of Oulu,
Linnanmaa, FIN-90570
Oulu, Finland

² Biotechnology Laboratory,
REDEC of Kajaani,
University of Oulu, FIN-
88600 Sotkamo, Finland

Bovine mastitis is caused mainly by certain *Staphylococcus* and *Streptococcus* species. The sequences of the 16S–23S rRNA spacer regions were determined for the nine species which cause mastitis: *Staphylococcus aureus*, *Staphylococcus chromogenes*, *Staphylococcus epidermidis*, *Staphylococcus hyicus*, *Staphylococcus simulans*, *Staphylococcus xylosus*, *Streptococcus agalactiae*, *Streptococcus dysgalactiae* and *Streptococcus uberis*. Significant variation was found between the spacer sequences of different species with the lengths of the spacers varying from 240 to 461 bp. Between genera the spacers shared only short conserved regions (8–9 bp) and within genera the sequence identities varied from 53 to 85%. This variation made it possible to construct specific primer pairs for these species and genera. The specificities of these primers were tested with 25 bacterial species and 51 isolates from cattle with clinical mastitis. The DNA-based identification of the mastitis species was mostly successful.

Keywords: *Staphylococcus*, *Streptococcus*, bovine mastitis, intergenic rRNA spacer

INTRODUCTION

Bovine mastitis is a multifactorial disease and is one of the most difficult to control. It can be caused by many different bacterial species, the most common of which are *Staphylococcus* and *Streptococcus* species. The prevalence of different species varies geographically, temporally and also due to control measures adopted in herds. In addition, different pathogens are typical of different types of mastitis (clinical, subclinical or heifer mastitis).

In many countries the most common bacterial species causing mastitis are *Staphylococcus aureus*, *Streptococcus dysgalactiae* and *Streptococcus uberis* (Aarestrup *et al.*, 1995; Ministry of Agriculture and Forestry Mastitis Committee, 1989; Jonsson *et al.*, 1991). *Streptococcus agalactiae* was previously the most common, but due to the use of antibiotic therapy its prevalence has decreased in many countries (Bramley & Dodd, 1984). The importance of coagulase-negative staphylococci (CNS) as pathogenic agents has been under discussion

(Kloos & Bannermann, 1994). CNS are part of normal skin flora and have previously been regarded only as minor pathogens. However, the proportion of CNS mastitis has increased markedly during the last decade (Myllys *et al.*, 1994). The most frequently isolated CNS species include *Staphylococcus hyicus*, *Staphylococcus simulans*, *Staphylococcus epidermidis*, *Staphylococcus chromogenes* and *Staphylococcus xylosus* (Harmon & Langlois, 1989; Jarp, 1991; Todhunter *et al.*, 1993; Aarestrup *et al.*, 1995; Honkanen-Buzalski *et al.*, 1994).

The identification of pathogens causing mastitis is important for disease control and epidemiological studies. In a clinical laboratory this is done using traditional microbiological methods (Klastrup & Schmidt Madsen, 1974). The species determination usually takes at least 24 h. There are no rapid and cheap methods available for the classification of CNS to the species level and therefore CNS species are usually determined only at the group level. There are commercial biochemical kits available for the determination of CNS species (e.g. API-Staph, Staph-Ident, Staph-Trac, Staph-Zym) but they have proved unreliable for the identification of veterinary pathogens (Watts & Yancey, 1994).

The development of molecular biological methods, such as nucleic acid analysis, protein patterns or fatty acid

Abbreviations: ATC-PCR, Air Thermo-Cycler PCR; CNS, coagulase-negative staphylococci.

The GenBank accession numbers for the sequences reported in this paper are U39765–U39773, U39813–U39814 and U90010–U90027.

profiles, has added possibilities for the rapid identification of bacteria (Busse *et al.*, 1996). Species-specific DNA sequences can be used for the identification of bacterial species. The 16S–23S rRNA intergenic spacer of the ribosomal RNA operon (*rrn*) has proven useful for identification of strains and species (Barry *et al.*, 1991; Jensen *et al.*, 1993; Gürtler & Stanisich, 1996). The spacer region is considered non-functional and is consequently argued to be under minimal selective pressure (Barry *et al.*, 1991). Its evolutionary rate is 10 times greater than that of 16S rDNA (Leblond-Bourget *et al.*, 1996), the sequence of which is routinely used for phylogenetic studies of bacteria (Ludwig & Schleifer, 1994). The high evolutionary rate makes it possible to distinguish closely related bacterial species.

The aims of this study were to determine the sequences of the 16S–23S rRNA intergenic spacers from the most common staphylococcal and streptococcal bovine mastitis pathogens and to investigate the possibility of designing species-specific oligonucleotide primers from the spacer sequences for the rapid identification of these bacterial species.

METHODS

Bacterial strains and culture conditions. All bacterial species and strains used in this study (Table 1) were grown aerobically in Tryptic Soy Broth (Difco) at 37 °C. Most of the specificity control species (Table 2) were obtained as cultures on agar plates or as DNA. The rest of the control species (e.g. all *Lactobacillus* species) were grown as recommended in the ATCC catalogue. Mastitic milk samples were obtained from the Food and Environmental Laboratory, Oulu, Finland and from the National Veterinary and Food Research Institute (EELA), Helsinki, Finland. Milk samples were cultivated overnight on M17 (Difco) agar plates at 37 °C. The total bacterial mass was harvested for DNA isolation. For PCR sensitivity tests, *Sta. aureus* DNA was isolated directly from mastitic milk according to the method described by Hynes *et al.* (1992).

Bacterial DNA isolation and amplification of spacer region.

Genomic DNA was isolated by the method of Anderson & McKay (1983) with modifications described by Steenson & Klaenhammer (1985). Oligonucleotide primers used for amplifying the 16S–23S rRNA intergenic spacer region were selected from the conserved regions at the 3' end of the 16S rRNA and the 5' end of the 23S rRNA genes. The sequences of the primers were 5' GTCGGAATCGCTAGTAATCG 3' for 16-1A (bases 1333–1352 in the *Escherichia coli* 16S rDNA sequence, A14565) and 5' GGGTTCCTCCCATTCGGA 3' for 23-1B (bases 130–114 in the *E. coli* 23S rDNA sequence, A14566). PCR was performed in a DNA thermal cycler 480 (Perkin-Elmer) with a DyNAzyme DNA Polymerase Kit (Finnzymes). A typical reaction mixture (50 µl) consisted of reaction buffer (10 mM Tris/HCl, pH 8.8, 1.5 mM MgCl₂, 50 mM KCl, 0.1% Triton X-100), 200 µM each dNTP, 1 µM 16-1A and 23-1B primers, 5 ng bacterial DNA and 0.5 U DyNAzyme DNA polymerase. The reaction mixtures were overlaid with mineral oil (Sigma), incubated at 92 °C for 2 min, then subjected to 30 cycles of 95 °C for 30 s, 55 °C for 30 s, 72 °C for 30 s. Amplification products were analysed on 1.5% (w/v) agarose gels.

Sequencing methods. PCR products from the spacer regions were sequenced directly by a cycle sequencing method using a CircumVent Thermal Cycle Dideoxy DNA Sequencing Kit (New England Biolabs). Sequencing reactions were performed using 15 cycles of 95 °C for 40 s, 55 °C for 30 s and 72 °C for 2 min. The oligonucleotide fragments produced were separated on polyacrylamide gels (Sequagel-6, National Diagnostics). Both strands of DNA were sequenced. If the sequencing was not performed using the total PCR products, the products were cloned into a TA-cloning vector (Original TA Cloning Kit, Invitrogen). Ten positive clones were selected from each species, and plasmid DNA was isolated by alkaline lysis–PEG precipitation (Sambrook *et al.*, 1989). Inserts were sequenced using a CircumVent Thermal Cycle Dideoxy DNA Sequencing Kit or DNA Sequencing Kit for the ABI Prism automatic sequencer (Perkin Elmer).

Analysis of sequence data. The sequences were analysed with DNASIS version 7.0 (Hitachi), and with GCG version 6.0 (Devereux *et al.*, 1984).

Table 1. Bacterial strains used for determination of 16S–23S rRNA spacer sequences

Species	Strain*	Length of spacer (bp)	tRNA in spacer	GenBank accession no.
<i>Str. agalactiae</i>	ATCC 27956	279	Ala	U39765
<i>Str. dysgalactiae</i>	ATCC 27957	285	Ala	U39767
<i>Str. uberis</i>	ATCC 27958	340	Ala	U39768
<i>Sta. aureus</i>	ATCC 25923	458	Ile	U39769
<i>Sta. chromogenes</i>	ATCC 43764	279	–	U39770
<i>Sta. epidermidis</i>	ATCC 12228	261–263	–	U39771, U90018–U90021
<i>Sta. hyicus</i>	KNS 264/92	375–380	–	U39772, U90010–U90015
		461	Ile	U90016
<i>Sta. simulans</i>	ATCC 11631	240–242	–	U39813, U90022–U90027
		333	Ile	U39814
<i>Sta. xylosus</i>	ATCC 12162	283	–	U39773, U90017

* The ATCC strains were obtained from the American Type Culture Collection, Rockville, MD, USA and the KNS strain was obtained from EELA, National Veterinary and Food Research Institute, Helsinki, Finland.

Table 2. Bacterial strains used for assessing the specificity of primers

Species	Strain	PCR products*
<i>Actinomyces pyogenes</i>	Isolated from mastitic milk‡	STAX (150 bp–2.8 kb)
<i>Bacillus cereus</i>	Isolated from mastitic milk‡	–
<i>Corynebacterium bovis</i>	EMO-M/18-I-17‡	STRA (1.5 kb); STAS (2.4 kb)
<i>Enterobacter cloacae</i>	Isolated from mastitic milk‡	–
<i>Enterococcus</i> sp.	Isolated from mastitic milk‡	STR
<i>Escherichia coli</i>	Isolated from mastitic milk‡	STAX (200 bp–1.6 kb)
<i>Klebsiella oxytoca</i>	Isolated from mastitic milk‡	–
<i>Klebsiella pneumoniae</i>	Isolated from mastitic milk‡	–
<i>Lactobacillus acidophilus</i>	ATCC 4356	–
<i>Lactobacillus casei</i>	ATCC 27092	–
<i>Lactobacillus delbrueckii</i>	ATCC 15808	–
<i>Lactobacillus helveticus</i>	ATCC 15009	–
<i>Lactococcus lactis</i>	F7/2 industrial starter strain	–
<i>Leptospira borgpetersenii</i> serovar hardjo subtype hardjobovist†	Clinical isolate	–
<i>Micrococcus varians</i>	Isolated from mastitic milk‡	STAX (500 bp–1 kb)
<i>Mycoplasma bovis</i>	Donetta	STAH (240 bp)
<i>Nocardia brasiliensis</i>	HAMBI 1960	–
<i>Peptococcus indolicus</i>	AHC 14990‡	–
<i>Propionibacterium freudenreichii</i>	ATCC 6207	–
<i>Pseudomonas aeruginosa</i>	Isolated from mastitic milk‡	–
<i>Str. bovis</i>	ATCC 27960	STR
<i>Str. thermophilus</i>	ATCC 19987	STR
<i>Str. zooepidemicus</i>	EMO-M/18-III-63‡	STR

* The DNA of these strains were tested with STAA, STAC, STAE, STAH, STAS, STAX, STRA, STRD, STRU, STA and STR primer pairs (Table 4). Primer pairs which yielded a PCR product are indicated.

† The subtype was determined by restriction endonuclease digestion of genomic DNA.

‡ Strains isolated from mastitic milk were obtained from EELA, National Veterinary and Food Research Institute, Helsinki, Finland where the species was determined.

PCR with species-specific primers. Based on the comparison of the nucleotide sequences of the spacers, specific primers were designed for each of the species. Genus-specific primers for the *Staphylococcus* and *Streptococcus* species included in this study were also constructed. The PCR reactions were carried out as above using optimized $MgCl_2$ concentrations for each primer pair (see Table 4). A rapid new generation thermal cycler, Air Thermo-Cycler (ATC, Idaho Technology), was also tested for amplification with the specific primer pairs. ATC-PCR uses glass capillary tubes instead of microcentrifuge tubes, which are incubated at 94 °C for 15 s, then 30 cycles of 94 °C for 0 s, 55 °C for 0 s and 72 °C for 15 s. This program takes a total of 15 min instead of 2 h using a conventional thermocycler. The reaction buffers included 0.25 µg BSA ml⁻¹. The $MgCl_2$ concentrations used in ATC-PCR are listed in Table 4.

RESULTS AND DISCUSSION

Amplification of the spacer region for sequencing

Primers complementary to conserved sequences near the 3' end of the 16S and the 5' end of the 23S rRNA genes were used to amplify the 16S–23S spacer from nine

species which commonly cause mastitis. In staphylococcal species more than one PCR product was detected, although in all cases one band predominated. It is known that the bacterial genome can contain several *rrn* operons, e.g. *Sta. aureus* has nine operons (Gürtler & Barrie, 1995). The size of the main band differed between species. The weaker bands may also represent heteroduplex molecules resulting from cross-hybridization of amplification products from different kinds of operons (Jensen & Straus, 1993).

Sequencing the spacer regions from total PCR products

The 16S–23S spacer region from streptococcal species and *Sta. chromogenes* could be sequenced directly from the PCR products. This indicated that there was a low level of sequence variation in the *rrn* operons within these species. Only the spacer sequence of *Sta. chromogenes* contained a variable nucleotide site (marked S at position 331 in Fig. 1a), with both G and C detected at this position. The previously published spacer sequence

of *Str. agalactiae* (GenBank accession no. L31412; Hall *et al.*, 1995) was compared to our *Str. agalactiae* sequence and only one insertion/deletion and one substitution were found (positions 98 and 351 in Fig. 1a). Sequencing of the spacer region from *Sta. aureus*, *Sta. epidermidis*, *Sta. hyicus*, *Sta. simulans* and *Sta. xyloso* gave multiple sequences that could not be read, suggesting the presence of insertions and/or deletions in the 16S–23S spacer regions of different *rrn* operons. The purification of the main PCR product from agarose gels for sequencing did not improve the readability of sequences, possibly because the insertions and/or deletions were too small to cause sufficient length variation in the spacers. The aligned nucleotide sequences of the spacer regions of streptococcal species are shown in Fig. 1(b) and the spacer region of *Sta. chromogenes* is shown in Fig. 1(a).

Sequencing the spacer regions from plasmid clones

The PCR products from the spacer regions of *Sta. aureus*, *Sta. epidermidis*, *Sta. hyicus*, *Sta. simulans* and *Sta. xyloso* were cloned into TA-cloning vector. Ten clones from different PCR amplifications were sequenced from *Sta. epidermidis*, *Sta. hyicus*, *Sta. simulans* and *Sta. xyloso*. Several different spacer sequences were obtained: five from *Sta. epidermidis*, seven from *Sta. hyicus*, eight from *Sta. simulans* and two from *Sta. xyloso* (Fig. 1a, clone types A–H). Most of the differences between the sequences from a single strain were single base insertions/deletions and substitutions. Given that the rate of artefactual misincorporation during PCR is relatively low and random, it is likely that these differences represent mutations present in the genome. The first sequence shown for each species (Fig. 1a) was obtained from clones from two independent amplifications. Some substitutions and insertions/deletions of tRNA genes were also found. All the streptococcal species had a tRNA-Ala gene in the spacer and *Sta. aureus*, *Sta. hyicus* and *Sta. simulans* had a tRNA-Ile gene in some of their *rrn*-operons (Fig. 1, Table 1). In general, however, the sequence variation within species was very low. From *Sta. aureus* only one clone was sequenced, since the sequences of all *Sta. aureus* *rrn* operons have been published previously by Gürtler & Barrie (1995). Our cloned *Sta. aureus* spacer sequence was 99% homologous to the allele *rrnF* (GenBank accession no. U11779; Gürtler & Barrie, 1995), with only three single base insertions/deletions and four mismatches.

Sequence variation between species

The pairwise sequence identities between 16S–23S spacer sequences of different species were calculated using the DNA Maximum Homology program of DNASIS (Table 3). This program favours short deletions over long ones and may overestimate the influence of a long deletion on the degree of DNA identity. The

sequence identity between species within a genus varied from 53 to 85%. The calculated sequence identities between species of different genera were about 50%, but because gaps were used freely, this is an overestimate. The sequences were aligned manually in Fig. 1(a) and (b) to minimize the number of sequence differences. The spacer sequences for *Str. bovis* (GenBank accession no. U39766; this study) and *Str. thermophilus* (U32965; Tilsala-Timisjärvi & Alatosava, 1997) were included in the alignment (Fig. 1b) to obtain more information about the variation of the spacer regions within the *Streptococcus* genus. This knowledge was important for the design of species-specific primers. The amount of sequence difference between staphylococci and streptococci prevented their reliable alignment. The only regions of sequence similarity (≥ 8 bp) between these two genera have been indicated in the alignments (Fig. 1a and b). Both conserved regions are important for RNA processing (Chiaruttini & Milet, 1993). When comparing the species within genera, the 3' terminal third of the spacer region was found to be highly variable and difficult to align, especially in staphylococcal species. The 5' terminus was more conserved (Fig. 1).

Design of primer pairs from the species-specific sequence regions

The variation between the 16S–23S spacer sequences of different mastitis pathogens made it possible to design specific primer pairs for each of the nine species considered in this study. The oligonucleotides were selected near the ends of the spacer region (Fig. 1, Table 4). In addition, primer pairs for *Staphylococcus* and *Streptococcus* were designed (Fig. 1, Table 4). The primer pairs were selected so that the same PCR cycling conditions could be employed for all of them. The optimum $MgCl_2$ concentrations were adjusted for both conventional PCR (Perkin-Elmer 480) and ATC-PCR (Table 4). There was some sequence variation within species at the primer regions. In those cases the primers were selected from the most common spacer type. Species-specific PCR primers have been published previously for some of the species (although not for strains isolated from cases of bovine mastitis) from the 16S rRNA gene, e.g. *Sta. aureus* and *Sta. epidermidis* (Saruta *et al.*, 1995; Gribaldo *et al.*, 1997).

Sensitivity of the primers

Products were detected using as little as 50 pg purified bacterial DNA, with 35 cycles in PCR. The DNA concentrations were measured with a spectrophotometer (Gene Quant II, Pharmacia Biotech) and with a fluorometer (TKO100 Mini-Fluorometer, Hoefer Scientific). When the sensitivity of the primers was estimated by isolating DNA directly without cultivation from mastitic milk samples containing *Sta. aureus*, a 1 μ l milk sample containing 40 c.f.u. yielded a PCR product (results not shown).

(a)

	1	10	20	30	40	50	60	70	80	90	100	
U11779	AAGGATA-TA	TTCGGAACAT	CTTCTTCAGA	AGATG----	CGAATAACG-	-TGACATATT	GTATTCAGTT	TTGAATGTTT	ATTT-----	AA	CATTCAAA--	196
au	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	87
cr	*****A*	*A*****T**	*GCT**T*AG	C**AA----	*****G	-A*****	*****	*****C*C	*****TCGAGG	*****C--	*****	93
ep-A	*****	*****	*****A*-**	*****A----	*G*****	-*****	*****	*****	*****	*****	*****	82
ep-B	*****	*****	*****A*-**	*****A----	*G*****	-*****	*****	*****	*****	*****	*****	82
ep-C	*****	*****	*****A*-**	*****A----	*G*****	-*****	*****	*****	*****	*****	*****	84
ep-D	*****	*****	*****A*-**	*****A----	*G*****	-*****	*****	*****	*****	*****	*****	84
ep-E	*****	*****	*****A*-**	*****A----	*G*****	-*****	*****	*****	*****	*****	*****	83
hy-A	*****A*	*A*****T**	*GC**AG*C	G*TA-----	*****AA	*****	*****	*****C*C	*****T-GAGG	*****C--	*****	90
hy-B	*****A*	*A*****T**	*GC**AG*C	G*TA-----	*****AA	*****	*****	*****C*C	*****T-GAGG	*****C--	*****	90
hy-C	*****A*	*A*****T**	*GC**AG*C	G*TA-----	*****G	*****	*****	*****C*C	*****T-GAGG	*****C--	*****	91
hy-D	*****A*	*A*****T**	*GC**AG*C	G*TA-----	*****AA	*****	*****	*****C*C	*****T-GAGG	*****C--	*****	90
hy-E	*****A*	*A*****T**	*G***AG*C	G*TA-----	*****G	-A*****	*****	*****C*C	*****T-GAGG	*****C--	*****	90
hy-F	*****A*	*A*****T**	*GC**AG*C	G*TA-----	*****AA	*****	*****	*****C*C	*****T-GAGG	*****C--	*****	90
hy-G	*****A*	*A*****T**	*G***AG*C	G*TA-----	*****G	-A*****	*****	*****C*C	***ATTG*G	-***M*TT	*****	93
si-A	*****	*****G	T***GCAG*	*AC*AAA-	*****	-*****	*****	*****	*****GA*	*****G-	*****	93
si-B	*****T**	*****G	T***GCAG*	*AC*AAA-	*****	-*****	*****	*****	*****GA*	*****G-	*****	94
si-C	*****	*****G	T***GCAG*	*AT*AAA-	*****	-*****	*****	*****	*****GA*	*****G-	*****	93
si-D	*****	*****A	T***GCAG*	*AT*AAA-	*****	-*****	*****	*****	*****GA*	*****G-	*****	93
si-E	*****	*****G	T***GCAG*	*AC*AAA-T	*****	-*****	*****	*****	*****GA*	*****G-	*****	94
si-F	*****T**	*****G	T***GCAG*	*AC*AAA-T	*****	-*****	*****	*****	*****GA*	*****G-	*****	93
si-G	*****T**	*****G	T***GCAG*	*AC*AAA-	*****	-*****	*****	*****	*****GA*	*****G-	*****	94
si-H	*****	*****G	T***GCAG*	*AC*AAA-	*****	-*****	*****	*****	*****GA*	*****CGT	*****	94
xy-A	*****	*****	*****T**G	*****ACAGA	*****AT	-*****	*****	*****C*C	***GG*G	T*****GT	*****	93
xy-B	*****	*****	*****T**G	*****ACAGA	*****AT	-*****	*****	*****C*C	***GG*G	T*****GT	*****	93
tRNA ¹¹⁰												
U11779	AAAATGGGCC	TATAGCTCAG	-CTGGTTAGA	GCG-ACGC-T	GATAAGCGTG	AGGTGGG-TG	GTTCGAGTCC	ACTTAGGCC	ACCA-----	TTATTGTGAC	286	
au	*****	*****	*****	*****C*****	*****	*****	*****	*****	*****	*****	178	
cr	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	100	
ep-A	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	89	
ep-B	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	89	
ep-C	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	91	
ep-D	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	91	
ep-E	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	90	
hy-A	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	97	
hy-B	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	97	
hy-C	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	98	
hy-D	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	97	
hy-E	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	97	
hy-F	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	97	
hy-G	*****	*****	GC*****	***C***TC	*****	*****C*	*****W*	*****	*****	*****W*	184	
si-A	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	100	
si-B	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	101	
si-C	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	100	
si-D	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	100	
si-E	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	100	
si-F	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	101	
si-G	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	101	
si-H	G*G*****	*****	*****	***C***C*	*****	*****	*****	*****	*****TTTGA	*T*****	192	
xy-A	GC-----	*****	*****	***C***C*	*****	*****	*****	*****	*****A	*A*****	106	
xy-B	GC-----	*****	*****	***C***C*	*****	*****	*****	*****	*****A	*A*****	106	
tRNA ¹¹⁰												
U11779	ATTGAAACT	AGATAAGTAA	GTAAAATA--	TAGATTTTAC	CAAGCAAAAC	CGAGTGAATA	AAGAGTTTTA	AATAAGCTTG	AATTCATAA-	GAAATAATCG	383	
au	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	275	
cr	*****	*****	*****	*****	*****	*****C-	***C*----	***-----	***A*----	***T*----	180	
ep-A	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	175	
ep-B	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	175	
ep-C	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	177	
ep-D	*****	*****	*****C*	***-----	*****	*****	*****	*****	*****G*--	*****	177	
ep-E	*****	*****	*****C*	***-----	*****	*****	*****	*****	*****	*****	176	
hy-A	*****	*****	*****	*****	*****	*****C-	***C*----	***-----	***A**A*-	***T*----	178	
hy-B	*****	*****	***C*----	*****	*****	*****C-	***C*----	***-----	***A**A*-	***T*----	178	
hy-C	*****	*****	*****	*****	*****	*****C-	***C*----	***-----	***A**A*-	***T*----	179	
hy-D	*****	*****	*****	*****	*****	*****C-	***C*----	***-----	***A**A*-	***T*----	178	
hy-E	*****	*****	*****	*****	*****	*****C-	***C*----	***-----	***A**A*-	***T*----	178	
hy-F	*****	*****	*****	*****	*****	*****C-	***C*----	***-----	***A**A*-	***T*----	178	
hy-G	*****	*****	*****	*****	*****	*****C-	***C*----	***-----	***A**A*-	***T*----	265	
si-A	*****	*****	*****A*--	*****	*****	*****	T*****T	*****T	T*****T	***TG****	193	
si-B	*****	*****	*****A*--	*****	*****	*****	T*****T	*****T	T*****T	***TG****	194	
si-C	*****	*****	*****A*--	*****	*****	*****	T*****T	*****T	T*****T	***TG****	193	
si-D	*****	*****	*****A*--	*****	*****	*****	T*****T	*****T	T*****T	***TG****	192	
si-E	*****	*****	*****A*--	*****	*****	*****	T*****T	*****T	T*****T	***TG****	193	
si-F	*****	*****	*****A*--	*****	*****	*****	T*****T	*****T	T*****T	***TG****	194	
si-G	*****	*****	*****A*--	*****	*****	*****	T*****T	*****T	T*****T	***TG****	194	
si-H	*****	*****	*****A*--	*****	*****	*****	T*****T	*****T	T*****T	***TG****	285	
xy-A	*****	*****	*****TA	*****	*****	*****A	T*****T	*****T	T*****T	***TG****	205	
xy-B	*****	*****	*****TA	*****	*****	*****A	T*****T	*****T	T*****T	***TG****	205	

Fig. 1. For legend see page 3497.

	310	320	330	340	350	360	370	380	390	400	
U11779	C-TAGTGTTC	-----	---GAAAGAA	CACTCACAAAG	A-TTAATAAC	GTGTTTAAAT	C-----TTT	TTATAAAA--	-----	-----G	441
au	*-----*	-----	*****	*****	*-----*	*****	*-----*	*****	-----	-----T	333
cr	*-----CG**	-----	---C***C	S*****T	*A*****	TG**GG*TG*	TGTTAT*G*	***ATTCGA-	-----	-----	245
ep-A	*C*****	-----	*****	*****	*-----*	TA***T*GC	TATTTA-***	***AT**CA-	-----	-----	239
ep-B	*-----*	-----	*****	*****C*	*-----*	TA***T*GC	TATTTA-***	***AT**CA-	-----	-----	238
ep-C	*-----*	-----	*****	*****	*-----*	TA***T*GC	TATTTA-***	***AT**CA-	-----	-----	240
ep-D	*-----*	-----	*****	*****	*-----*	TA***T*GC	TATTTA-***	***AT**CA-	-----	-----	240
ep-E	*-----*	-----	*****	*****	*-----*	TA***T*GC	TATTTA-***	***AT**CA-	-----	-----	239
hy-A	*-----CG**	-----	---C***C	G*****T	*A*****	TG**GG*TG*	TGTTAA*G*	***CTTCG*A	TGACAGATGT	T-----TT*	258
hy-B	*-----CG**	-----	---C***C	G*****T	*A*****	TG**GG*TG*	TGTTAA*G*	***CTTCG*A	TGACAGATGT	T-----TT*	258
hy-C	*-----CG**	-----	---C**NC	G*****T	*A*****	TG**GG*TG*	TGTTAA*G*	***CTTCG*A	TGACAGATGT	T-----TT*	259
hy-D	*-----CG**	-----	---C**NC	G*****T	*A*****	TG**GG*TG*	TGTTAA*G*	***CTTCG*A	TGACAGATGT	T-----TT*	258
hy-E	*-----CG**	-----	---G***C	G*****T	*A*****	TG**GG*TG*	TGTTAA*G*	***CTTCG*A	TGACAGATGT	T-----TT*	264
hy-F	*-----CG**	-----	---C***C	G*****T	*A*****	TG**GG*TG*	TGTTAA*G*	***CTTCG*A	TGACAGATGT	T-----TT*	264
hy-G	*-----CG**	-----	---C***C	G*****T	*A*****	TG**GG*TG*	TGTTAA*G*	***CTTCG*A	TGACAGATGT	T-----TT*	345
si-A	*-----AA**	AATTGCGAC	GGCA**C**T	T*****T	*-----*	-----	-----	-----	-----	-----	241
si-B	*-----AA**	AATTGCGAC	GGCA**C**T	T*****T	*-----*	-----	-----	-----	-----	-----	242
si-C	*-----AA**	AATTGCGAC	GGCA**C**T	T*****T	*-----*	-----	-----	-----	-----	-----	241
si-D	*-----AA**	AATTGCGAC	GGCA**C**T	T*****T	*-----*	-----	-----	-----	-----	-----	240
si-E	*-----AA**	AATTGCGAC	GGCA**C**T	T*****T	*-----*	-----	-----	-----	-----	-----	241
si-F	*-----AA**	AATTGCGAC	GGCA**C**T	T*****T	*-----*	-----	-----	-----	-----	-----	242
si-G	*-----AA**	AATTGCGAC	GGCA**C**T	T*****T	*-----*	-----	-----	-----	-----	-----	242
si-H	*-----AA**	AATTGCGAC	GGCA**C**T	T*****T	*-----*	-----	-----	-----	-----	-----	333
xy-A	*-----*	-----	*****	*****	*-----*	---A***-*	GGG-----*	***A-----	-----	-----	254
xy-B	*-----*	-----	*****	*****	*-----*	---A***-*	GGG-----*	***A-----	-----	-----	254
	410	420	430	440	450	460	470	480	490	500	
U11779	AAAACGTTTA	GCAGACAATG	AGTTAAATTA	TTTAAAGCA	GAGTTTACTT	ATGTAAATGA	GCATTTAAAA	TAATGAAAA-	CGAAGCGTA	TGTGAGGTT	540
au	*****	*****	*****	*****	*****	*****	*T*****	*****	*****	***TAA***	432
cr	-----	-----	-----	-----	-----	-----	-----	-----	-----	***AA***A*	263
ep-A	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	239
ep-B	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	238
ep-C	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	240
ep-D	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	240
ep-E	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	239
hy-A	*****G	T***T*T**	*A*CGC*AAC	AAGAGCGAAG	*CG*****	CC***GCA*	CTGAG*G*TT	*GTGCGCGA	T*****A*	**CA*A**A*	358
hy-B	*****G	T***T*T**	*A*CGC*AAC	AAGAGCGAAG	*C*C*****	CC***GCA*	CTGAG*G*TT	*GTGCGCGA	T*****A*	**CA*A**A*	358
hy-C	*****G	T***T*T**	*A*CGC*AAC	AAGAGCGAAG	*C*C*****	CC***GCA*	CTGAG*G*TT	*GTGCGCGA	T*****A*	**CA*A**A*	359
hy-D	*****G	T***T*T**	*A*CGC*AAC	AAGAGCGAAG	*C*C*****	T***G*A*	CTGAG*G*TT	*GTGCGCGA	TN*****A*	**CA*A**A*	358
hy-E	***G*****	T***T*T**	*A*CGC*AAC	AAGAGCGAAG	*C*C*****	T***G*A*	CTGAG*G*TT	*GTGCGCGA	T*****A*	**CA*A**A*	364
hy-F	*****G	T***T*T**	*A*CGC*AAC	AAGAGCGAAG	*CS-*****	CC***GCA*	CTGAG*G*TT	*GTGCGCGA	T*****A*	**CA*A**A*	363
hy-G	*****G	T***T*T**	*A*CGC*AAC	AAGAGCGAAG	NC-*****	CC***RCA*	CTGAG*G*TT	*GTGCGCGA	TN***N**A*	**CA*A**A*	444
si-A	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	241
si-B	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	242
si-C	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	241
si-D	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	240
si-E	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	241
si-F	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	242
si-G	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	242
si-H	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	333
xy-A	-----	-----	-----	-----	-----	-----	-----	-----	***AC	*-CGT***G	268
xy-B	-----	-----	-----	-----	-----	-----	-----	-----	***AC	*-CGT***G	268
	510	520									
U11779	TGACTTATAA	AAATGGTGA	AACATA	566							
au	*****	*****	*****	458							
chr	**C*---A**	*C*---C*	-A*-	279							
ep-A	--*---CA**	*T*****	*****	262							
ep-B	--*---CA**	*T*****	*****	261							
ep-C	--*---CA**	*T*****	*****	263							
ep-D	--*---CA**	*T*****	*****	263							
ep-E	--*---CA**	*T*****	*****	262							
hy-A	**---C*A**	*C*---C*	T*A*-	375							
hy-B	**---C*A**	*C*---C*	T*A*-	375							
hy-C	**---C*A**	*C*---C*	T*A*-	376							
hy-D	**---C*A**	*C*---C*	T*A*-	375							
hy-E	*AT--C*A-C	*CG*TTA		378							
hy-F	**C*---A**	*C*---C*	T*A*-	380							
hy-G	**C*---A**	*C*---C*	T*A*-	461							
si-A	-----	-----	-----	241							
si-B	-----	-----	-----	242							
si-C	-----	-----	-----	241							
si-D	-----	-----	-----	240							
si-E	-----	-----	-----	241							
si-F	-----	-----	-----	242							
si-G	-----	-----	-----	242							
si-H	-----	-----	-----	333							
xy-A	*-----*	*-----*	**A*-	283							
xy-B	*-----*	*-----*	**A*-	283							

Fig. 1. For legend see facing page.

(b)

	1	10	20	30	40	50	60	70	80	90	100	
L31412	AAGGATAA--	GGAAACCTGC	CATTTCGRTC	TTGTTTAGTT	TTGAGAGGTC	TTG	-----	-----	---TGGGGGCC	TTAGCTCAGC	TGGGAGA--CN	97
ag	*****--	*****G*	*****G*	*****G*	*****G*	*****G*	-----	-----	*****G*	*****G*	*****G*	78
dy	*****--T	*****AC*T	T*GGGT*G*	A*****	*****G*	*****G*	-----	-----	*****G*	*****G*	*****G*	77
ub	*****--	*****AC*T	TGG*AA*	A*****	*****G*	*****G*	CAAGACG	CAGAGACAAA	CTG*****	*****G*	*****G*	97
th	*****AAC	*****TGTA	*T*GA*TT*	A*****	*****G*	*****G*	-----	-----	*****G*	*****G*	*****G*	78
bo	*****AC	*****GCA	*G**GG*A	A*****	*****G*	*****G*	-----	-----	*****G*	*****G*	*****G*	77
----- tRNA ^{Ala}												
	110	120	130	140	150	160	170	180	190	200		
L31412	OCTGCTTTGC	ACGCAGGAGG	TCAGCGNNTC	GATCNCGCTA	GGCTCCATT-	GAATCGAAAG	GTTC-AA--	-----	-----	-----	162	
ag	*****	*****G**	*****C*****	*****C*****	*****A	*G**A**G*	A*C*T**A--	-----	-----	-----	143	
dy	*****	*****G**	*****C*****	*****C*****	*****A	*G**A**G*	A*C*T**A--	-----	-----	-----	145	
ub	*****	*****G**	*****C*****	*****C*****	*****A	*G**AC*G--	---A*CTGA	ACTTAATAGA	AGTGAAGTTT	CATTGTATCT	193	
th	*****	*****G**	*****C*****	*****C*****	*****A	*****A	*****A	-----	-----	-----	143	
bo	*****	*****G**	*****C*****	*****C*****	*****A	*****A	*****A	-----	-----	-----	143	
	210	220	230	240	250	260	270	280	290	300		
L31412	---ATTGTT	CATTGAAAT	TGAATATCTA	TATCAAAATC	CACGATCTA-	GAAATAGAT-	TGTA-GAAA-	GTAACAAGAA	AATAAACCGA	AAA-----	247	
ag	-----	*****	*****	*****	*****A-	*****A-	*****A-	*****	*****	*****	228	
dy	---A*C**C	*****	*****	*****	*****A*	*****T**	***C**TA	*****	*****	*****	233	
ub	TAGT**A**C	*****	*****	*****	*****AT-	*****AT**	*****A	*****	*****	***AAACGA	289	
th	--GT*****	*****	*****	*****	**T--**G**A	*T***TACAT	A-***T*	*****	*****	**-----	229	
bo	--G*****C	*****	*****	*****	*****TC*A	*****T*AT	***--T*	*****	*****	**G-----	230	
	310	320	330	340	350	360						
L31412	---CGCTGT	GAA--TATTT	AATGAGTTT	-CTAGTTT	AAGAACTAG	TTTAA--TAA	298					
ag	-----	*****	*****	*****	*****G*	*****G*	279					
dy	-----	*****A*C	*****T*****	T*****-A*	G*T*T*****	*AA*GA**	285					
ub	TAAA*---	***CA***A	**AA*AA*CA	AGA**G*C**	*G*---*	G-A*---	340					
th	-----	*****	*****	---*G*CG*	***GCCAA*	-----	272					
bo	-----	**-----*	*****	--A*G*CA*	C****-C*	--A*-----	272					

Fig. 1. Alignments of the 16S–23S intergenic spacer sequences from (a) staphylococcal species and (b) streptococcal species. Symbols: au, *Sta. aureus*; cr, *Sta. chromogenes*; ep, *Sta. epidermidis*; hy, *Sta. hyicus*; si, *Sta. simulans*; xy, *Sta. xylosus*; ag, *Str. agalactiae*; dy, *Str. dysgalactiae*; ub, *Str. uberis*; th, *Str. thermophilus* (GenBank accession no. U32965; Tilsala-Timisjärvi & Alatossava, 1997); bo, *Str. bovis* (U39766; this study). The clone types are indicated after the species in capital letters (A–H). U11779 and L31412 are the spacer sequences of *Sta. aureus* and *Str. agalactiae*, respectively (Gürtler & Barrie, 1995; Hall et al., 1995). The species-specific primers are presented in bold, and the genus-specific primers are boxed. The regions (minimum length 8 bp) identical in all staphylococcal and streptococcal species are indicated with double lines. Identical nucleotides are indicated with asterisks, deletions with dashed lines, and transitions and transversions compared to the uppermost sequence of the alignment are indicated with capital letters. The locations of tRNA genes are indicated above the aligned sequences.

Specificity of the primers

Each primer pair was tested at least twice with DNA from 31 different bacterial species (Tables 1 and 2) using 5 ng template DNA in a 20 µl PCR reaction with conventional PCR. Appropriate controls were included in PCR; a positive control for DNA and primers and a negative control for PCR mix. In addition, the primer pairs were tested after raising the concentration of DNA from staphylococcal and streptococcal species up to 100 ng in a 20 µl PCR reaction. The performance of ATC-PCR was tested with nine DNA samples (Table 1). The 'species-specific' primer pairs (Table 4) amplified their targets mainly as expected, amplifying only the corresponding DNA (Table 2, Fig. 2). The unexpected products with STAX, STRA, STAH and STAS primer pairs differed in size from the expected products, and no amplification with the 'genus-specific' primers was observed. If these cases represented true *Streptococcus* or *Staphylococcus* species, a product with the genus-specific primers would have been expected also. The genus-specific primers amplified as expected: the *Staphylococcus* specific primers amplified only staphylococcal DNA, and the *Streptococcus* specific primers amplified streptococcal, lactococcal and enterococcal

DNA (Table 2). Lactococci and enterococci are very closely related to streptococci and have similar sequences in the primer regions (Chiaruttini & Milet, 1993; Sechi & Daneo-Moore, 1993; Tilsala-Timisjärvi & Alatossava, 1997; Wood & Holzapel, 1995).

Validity of the primers with mastitis samples

The original staphylococcal and streptococcal strains from which the spacer regions were sequenced were mostly ATCC strains (Table 1), not isolated from bovine mastitis. To investigate if there was sequence variation between strains within a species, which could affect the utility of the assays, we performed PCR on 51 isolates from cases of bovine mastitis. The conventional species determination was made in a veterinary clinical microbiology laboratory (Food and Environmental Laboratory, Oulu, Finland or National Veterinary and Food Research Institute Helsinki, Finland). PCR reactions used purified DNA from each cultivated isolate (Table 5). The staphylococcal genus-specific primer pair distinguished the genus with 100% sensitivity and specificity. The streptococcal genus-specific primer pair detected one *Sta. aureus* isolate and all streptococcal isolates.

Table 3. Sequence identity between 16S–23S rDNA spacer sequences of different *Staphylococcus* and *Streptococcus* species

au, *Sta. aureus* clone; cr, *Sta. chromogenes*; ep-A, *Sta. epidermidis* clone type A; hy-A, *Sta. hyicus* clone type A; si-A, *Sta. simulans* clone type A; xy-A, *Sta. xylosus* clone type A; ag, *Str. agalactiae*; dy, *Str. dysgalactiae*; ub, *Str. uberis*.

DNA identity (%)*								
Spacer	cr	ep-A	hy-A	si-A	xy-A	ag	dy	ub
au	63	68	69	58	65	50	52	53
cr		74	71	66	70	52	54	52
ep-A			59	71	77	52	52	52
hy-A				53	57	51	52	51
si-A					71	51	51	50
xy-A						53	53	52
ag							85	70
dy								70

* The tRNA gene region was excluded from the *Sta. aureus* (au) spacer sequence for analysis. The calculations were made by the DNA Maximum Homology program of DNASIS.

Ninety percent of the isolates identified conventionally at the species level (36/40) were recognized with the species-specific primers. The reason for the failure of

Fig. 2. The PCR products from different streptococcal and staphylococcal species amplified with corresponding species-specific primer pairs. Lanes: 1, *Sta. aureus* ATCC 25923; 2, *Sta. chromogenes* ATCC 43764; 3, *Sta. epidermidis* ATCC 12228; 4, *Sta. hyicus* KNS 264/92; 5, *Sta. simulans* ATCC 11631; 6, *Sta. xylosus* ATCC 12162; 7, 1 kb ladder (Gibco BRL); 8, *Str. agalactiae* ATCC 27956; 9, *Str. dysgalactiae* ATCC 27957; 10, *Str. uberis* ATCC 27958.

identification for four isolates may be sequence variation between strains in the primer regions. In addition, three isolates gave a PCR product with a second species-specific primer pair (Table 5). These products may also be due to sequence variation. Alternatively, the milk samples may have contained minor amounts of these other bacterial species. Only five of the CNS isolates were identified to the species level in the clinical

Table 4. Species- and genus-specific oligonucleotide primers from the 16S–23S rRNA intergenic spacer region for bovine mastitis pathogens

Species	Oligonucleotide*	Sequence (5'–3')	Length (nt)	MgCl ₂ in PCR (mM)†	MgCl ₂ in ATC-PCR (mM)†	Size of the main PCR product (bp)
<i>Str. agalactiae</i>	STRA-AgI	GGAAACCTGCCATTGCG	18	3.0	5.0	280
	STRA-AgII	TAACCTTAACCTTATTAACCTAG	22			
<i>Str. dysgalactiae</i>	STRD-DyI	TGGAACACGTTAGGGTCG	18	3.0	3.0	270
	STRD-DyII	CTTTTACTAGTATATCTTAACTA	23			
<i>Str. uberis</i>	STRU-UbI	TAAGGAACACGTTGGTTAAG	20	1.5	3.0	330
	STRU-UbII	TCCAGTCCTTAGACCTTCT	19			
<i>Sta. aureus</i>	STAA-AuI	TCTTCAGAAAGATGCGGAATA	20	2.0	3.0	420
	STAA-AuII	TAAGTCAAACGTTAACATACG	21			
<i>Sta. chromogenes</i>	STAC-ChrI	ACGGAATATCGCTTTTAAGC	20	1.5	3.0	250
	STAC-ChrII	CGTTTACATTTCGGCTTTTCG	19			
<i>Sta. epidermidis</i>	STAE-EpI	TCTACGAAGATGAGGGATA	19	3.0	3.0	240
	STAE-EpII	TTTCCACCATATTTTGAATTGT	22			
<i>Sta. hyicus</i>	STAH-HyI	TACGGAATATCGCCTTAGG	19	1.5	4.0	250
	STAH-HyII	AAAACATCTGTCATCCGAAG	20			
<i>Sta. simulans</i>	STAS-SiI	ATTTCGGAACAGTTTCGCAG	19	1.5	4.0	220
	STAS-SiII	ATTGTGAGTAATCGTTTGCC	20			
<i>Sta. xylosus</i>	STAX-XyI	TCTTTAGAAAGATGACAGAGG	20	1.5	3.0	260
	STAX-XyII	TGACTTTTAAACACGACGAAG	20			
<i>Streptococcus</i> genus	STR I	TGTTTAGTTTTGAGAGGTCTTG	22	1.5	5.0	150–210
	STR II	CGTGGAATTTGATATAGATATTC	23			
<i>Staphylococcus</i> genus	STA I	GGAATAACGTGACATATTGTA	21	1.5	3.0	100–200
	STA II	TTCACCTCGTTTGTGCTTGG	19			

* Primer I in each primer pair is from the 5' terminus of the spacer and primer II is from the 3' terminus of the spacer.

† Optimal MgCl₂ concentrations in PCR reactions.

Table 5. Specificity of the primer pairs with DNA of 51 isolates from mastitic milk

Bacterial sample	No. of samples	Identification with PCR
<i>Sta. aureus</i>	19	<i>Staphylococcus</i> , <i>Sta. aureus</i>
<i>Sta. aureus</i>	2	<i>Staphylococcus</i>
<i>Sta. aureus</i>	1	<i>Staphylococcus</i> , <i>Sta. aureus</i> , <i>Streptococcus</i>
<i>Sta. aureus</i>	2	<i>Staphylococcus</i> , <i>Sta. aureus</i> , <i>Str. agalactiae</i>
<i>Sta. aureus</i>	1	<i>Staphylococcus</i> , <i>Sta. aureus</i> , <i>Sta. hyicus</i>
<i>Sta. chromogenes</i> *	1	<i>Staphylococcus</i> , <i>Sta. chromogenes</i>
<i>Sta. epidermidis</i> *	1	<i>Staphylococcus</i> , <i>Sta. epidermidis</i>
<i>Sta. hyicus</i> *	1	<i>Staphylococcus</i> , <i>Sta. hyicus</i>
<i>Sta. simulans</i> *	1	<i>Staphylococcus</i> , <i>Sta. simulans</i>
<i>Sta. xylosus</i> *	1	<i>Staphylococcus</i>
CNS	1	<i>Staphylococcus</i> , <i>Sta. chromogenes</i>
CNS	3	<i>Staphylococcus</i> , <i>Sta. simulans</i>
CNS	4	<i>Staphylococcus</i> , <i>Sta. xylosus</i>
CNS	3	<i>Staphylococcus</i>
<i>Str. uberis</i>	5	<i>Streptococcus</i> , <i>Str. uberis</i>
<i>Str. uberis</i>	1	<i>Streptococcus</i> , <i>Str. uberis</i> , <i>Str. agalactiae</i>
<i>Str. agalactiae</i>	2	<i>Streptococcus</i> , <i>Str. agalactiae</i>
<i>Str. dysgalactiae</i>	1	<i>Streptococcus</i> , <i>Str. dysgalactiae</i>
<i>Str. dysgalactiae</i>	1	<i>Streptococcus</i>

* CNS samples identified at the species level.

laboratories (Table 5). Eight out of eleven CNS samples (73 %) were detected by PCR with the species-specific primers. With the API-Staph system we identified one of the remaining CNS samples as *Staphylococcus warneri*, which could not be detected with our PCR primers. Two other CNS samples remain unidentified.

Applicability of the results

The important aims of this study were to investigate the level of sequence variation in the 16S–23S spacer region between different staphylococcal and streptococcal mastitis pathogens and the potential use of this variation for the rapid identification of these pathogens to species level. The conventional microbiological and biochemical methods for identifying mastitis pathogens are time-consuming. In addition, CNS species are still difficult to identify, and thus species identification is not done routinely, in spite of their increased prevalence in mastitis cases (White *et al.*, 1989; Myllys *et al.*, 1994). The advantages of PCR compared to other diagnostic methods are rapidity and facility. Furthermore, screening for antibiotic resistance could be combined with PCR-based species identification. In practice, the PCR identification could be done in two stages. First, performing the tests for the genus (STA, STR), *Sta. aureus*, the streptococcal species and antibiotic resistance (e.g. *blaZ* for β -lactamase gene). The second PCR would be the identification of CNS species if necessary. The main disadvantage of a PCR identification method might be excessive sensitivity, with minor contaminants in samples resulting in misdiagnosis. The multiple copy number of *rrn* operons in a bacterial genome is an

advantage, increasing the sensitivity of PCR, but the potential for variation within a species may reduce the sensitivity for all strains. The current primer pairs work reasonably well, but not yet with 100 % of isolates. Improvement in both primer design and PCR conditions is still needed. The primer pairs designed in this study proved to be specific and appropriately sensitive, such that the normal bacterial flora in milk did not give PCR products. However, a more extensive study of sequence variation in strains of these species isolated from cases of bovine mastitis is still needed to critically assess specificity. The traditional species identification would have to be confirmed by sequencing part(s) of the 16S rRNA gene, since even biochemical identification systems are not 100 % reliable. Our future work will focus on detecting mastitic pathogens by PCR directly from milk.

ACKNOWLEDGEMENTS

We thank Professor Tuula Honkanen-Buzalski from the National Veterinary and Food Research Institute, Helsinki, Finland and Taito Palo and Eija Eskola from the Food and Environmental Laboratory, Oulu, Finland for collaboration and mastitic milk samples, Associate Professor Satu Pyörälä, University of Helsinki, Helsinki, Finland and the Anaerobic Laboratory in the National Public Health Institute, Helsinki, Finland for *Peptococcus indolicus* strains, Professor Karl-Erik Johansson from the National Veterinary Institute, Uppsala, Sweden for providing *Mycoplasma bovis* DNA and Dr Roger Marshall, Massey University, Palmerston North, New Zealand for providing *Leptospira borgpetersenii* serovar *hardjo* subtype *hardjobovis* DNA.

REFERENCES

- Aarestrup, F. M., Wegener, H. C., Rosdahl, V. T. & Jensen, N. E. (1995). Staphylococcal and other bacterial species associated with intramammary infections in Danish dairy herds. *Acta Vet Scand* 36, 475–487.
- Anderson, D. G. & McKay, L. L. (1983). Simple and rapid method for isolating large plasmid DNA from lactic streptococci. *Appl Environ Microbiol* 46, 549–552.
- Barry, T., Colleran, G., Glennon, M., Dunican, L. K. & Gannon, F. (1991). The 16S/23S ribosomal spacer region as a target for DNA probes to identify Eubacteria. *PCR Methods Appl* 1, 51–56.
- Bramley, A. J. & Dodd, F. H. (1984). Reviews of the progress of dairy science: mastitis control – progress and prospects. *J Dairy Sci* 51, 481–512.
- Busse, H.-J., Denner, E. B. M. & Lubitz, W. (1996). Classification and identification of bacteria: current approaches to an old problem. Overview of methods used in bacterial systematics. *J Biotechnol* 47, 3–38.
- Chiaruttini, C. & Milet, M. (1993). Gene organization, primary structure and RNA processing analyses of a ribosomal RNA operon in *Lactococcus lactis*. *J Mol Biol* 230, 57–76.
- Devereux, J., Haeberli, P. & Smithies, O. (1984). A comprehensive set of sequence analysis programs for VAX and Convex systems. *Nucleic Acids Res* 12, 387–395.
- Gribaldo, S., Cookson, B., Saunders, N., Marples, R. & Stanley, J. (1997). Rapid identification by specific PCR of coagulase-negative staphylococcal species important in hospital infection. *J Med Microbiol* 46, 45–53.
- Gürtler, V. & Barrie, H. D. (1995). Typing of *Staphylococcus aureus* strains by PCR-amplification of variable length 16S–23S rDNA spacer regions: characterization of spacer sequences. *Microbiology* 141, 1255–1265.
- Gürtler, V. & Stanisich, V. A. (1996). New approaches to typing and identification of bacteria using the 16S–23S rDNA spacer region. *Microbiology* 142, 3–16.
- Hall, L. M. C., Duke, B. & Urwin, G. (1995). An approach to the identification of the pathogens of bacterial meningitis by the polymerase chain reaction. *Eur J Clin Microbiol Infect Dis* 14, 1090–1094.
- Harmon, R. J. & Langlois, B. E. (1989). Mastitis due to coagulase-negative *Staphylococcus* species. *Agri-Practice* 10, 29–34.
- Honkanen-Buzalski, T., Myllys, V. & Pyörälä, S. (1994). Bovine clinical mastitis due to coagulase-negative staphylococci and their susceptibility to antimicrobials. *J Vet Med B* 41, 344–350.
- Hynes, W. L., Ferretti, J. J., Gilmore, M. S. & Segarra, R. A. (1992). PCR amplification of streptococcal DNA using crude cell lysates. *FEMS Microbiol Lett* 94, 139–142.
- Jarp, J. (1991). Classification of coagulase-negative staphylococci isolated from bovine clinical and subclinical mastitis. *Vet Microbiol* 27, 151–158.
- Jensen, M. A. & Straus, N. (1993). Effect of PCR-conditions on the formation of heteroduplex and single-stranded DNA products in the amplification of bacterial ribosomal DNA spacer regions. *PCR Methods Appl* 3, 186–194.
- Jensen, M. A., Webster, J. A. & Straus, N. (1993). Rapid identification of bacteria on the basis of polymerase chain reaction-amplified ribosomal DNA spacer polymorphisms. *Appl Environ Microbiol* 59, 945–952.
- Jonsson, P., Olsson, S.-O., Olofson, A.-S., Fälth, C., Holmberg, O. & Funke, H. (1991). Bacteriological investigations of clinical mastitis in heifers in Sweden. *J Dairy Res* 58, 179–185.
- Klastrup, O. & Schmidt Madsen, P. (1974). Scandinavian recommendations on examination of quarter milk samples. *Nordisk Veterinaermed* 26, 197–204.
- Kloos, W. E. & Bannermann, T. L. (1994). Update on significance of coagulase-negative staphylococci. *Clin Microbiol Rev* 7, 117–140.
- Leblond-Bourget, N., Philippe, H., Mangin, I. & Decaris, B. (1996). 16S rRNA and 16S to 23S internal transcribed spacer sequence analyses reveal inter- and intraspecific *Bifidobacterium* phylogeny. *Int J Syst Bacteriol* 46, 102–111.
- Ludwig, W. & Schleifer, K. H. (1994). Bacterial phylogeny based on 16S and 23S rRNA sequence analysis. *FEMS Microbiol Rev* 15, 155–173.
- Ministry of Agriculture & Forestry Mastitis Committee (1989). *Prevention of Mastitis*. Memo no. 19/1989, Ministry of Agriculture and Forestry, Helsinki, Finland.
- Myllys, V., Honkanen Buzalski, T., Huovinen, P., Sandholm, M. & Nurmi, E. (1994). Association of changes in the bacterial ecology of bovine mastitis with changes in the use of milking machines and antibacterial drugs. *Acta Vet Scand* 35, 329–335.
- Sambrook, J., Fritsch, E. F. & Maniatis, T. (1989). *Molecular Cloning: a Laboratory Manual*, 2nd edn. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory.
- Saruta, K., Hoshina, S. & Machida, K. (1995). Genetic identification of *Staphylococcus aureus* by polymerase chain reaction using single-base-pair mismatch in 16S ribosomal RNA gene. *Microbiol Immunol* 39, 839–844.
- Sechi, L. & Daneo-Moore, L. (1993). Characterization of intergenic spacers in two *rrn* operons of *Enterococcus hirae* ATCC 9790. *J Bacteriol* 175, 3213–3219.
- Steenison, L. R. & Klaenhammer, T. R. (1985). *Streptococcus cremoris* M12R transconjugants carrying the conjugal plasmid pTR2030 are insensitive to attack by lytic bacteriophages. *Appl Environ Microbiol* 50, 851–858.
- Tilsala-Timisjärvi, A. & Alatossava, T. (1997). Development of oligonucleotide primers from the 16S–23S rRNA intergenic sequences for identifying different dairy and probiotic lactic acid bacteria by PCR. *Int J Food Microbiol* 35, 49–56.
- Todhunter, D. A., Cantwell, L. L., Smith, K. L., Hoblet, K. H. & Hogan, J. S. (1993). Characteristics of coagulase-negative staphylococci isolated from bovine intramammary infections. *Vet Microbiol* 34, 373–380.
- Watts, J. L. & Yancey, R. J., Jr (1994). Identification of veterinary pathogens by use of commercial identification systems and new trends in antimicrobial susceptibility testing of veterinary pathogens. *Clin Microbiol Rev* 7, 346–356.
- White, D. G., Harmon, R. J., Matos, J. E. S. & Langlois, B. E. (1989). Isolation and identification of coagulase-negative *Staphylococcus* species from bovine body sites and streak canals of nulliparous heifers. *J Dairy Sci* 72, 1886–1892.
- Wood, B. J. B. & Holzapfel, W. H. N. (1995). *Genera of Lactic Acid Bacteria*. Glasgow: Blackie Academic & Professional.

Received 3 March 1997; revised 23 June 1997; accepted 21 July 1997.