CORRIGENDUM

NMR analyses of compatible solutes in a halotolerant *Brevibacterium* sp.

Shinichi Nagata, Kyoko Adachi and Hiroshi Sano

Microbiology (1996), **142**, 3355–3362

In Table 2, p. 3360, the values given for 'Amino acids' and 'Total' in the top and bottom rows were incorrect, and the words 'and hydroxyproline' were missing from the footnote. The correct version of the table is given below.

Table 2. Internal solute and ion concentrations (mM) in *Brevibacterium* sp. grown in C medium with different NaCl concentrations

NaCl concn (M) ...	0	0.1	0.5	1.0	1.5	2.0
Amino acids*	89.4	97.2	124.6	140.0	198.2	149.7
Glutamate	134.7 ± 13.6	193.7 ± 3.3	241.4 ± 29.3	209.5 ± 13.3	193.3 ± 9.8	156.9 ± 0.5
Glycine betaine	99.2 ± 5.1	94.9 ± 10.4	136.3 ± 28.1	126.0 ± 24.1	135.0 ± 1.2	140.5 ± 4.8
Ectoine	0.0	0.0	108.5 ± 1.3	252.1 ± 18.1	385.3 ± 8.9	480.3 ± 5.2
Hydroxyectoine	0.0	0.0	0.0	0.0	4.8 ± 0.1	63.9 ± 3.7
Hydroxyproline	44.5 ± 3.9	51.4 ± 5.8	59.6 ± 15.3	97.9 ± 17.6	155.5 ± 0.9	65.6 ± 11.7
TMAH	66.7 ± 2.8	94.7 ± 4.5	123.6 ± 5.1	110.9 ± 12.9	111.5 ± 10.1	116.8 ± 10.6
Na ⁺	0.0	16.2 ± 7.2	35.5 ± 4.7	43.4 ± 7.7	77.2 ± 1.1	117.0 ± 18.8
K ⁺	288.1	297.8	364.4	372.9	350.1	327.9
Total	722.6	845.9	1193.9	1352.7	1610.9	1618.6

* Total amounts of amino acids, excluding glutamate and hydroxyproline, determined by amino acid analysis (Mimura *et al.*, 1994).