etiologic agent of Chagas disease. *Science* **309**, 409–415.

Feil, H., Feil, W. S., Chain, P. & 17 other authors (2005). Comparison of the complete genome sequences of *Pseudomonas syringae* pv. *syringae* B728a and pv. *tomato* DC3000. *Proc Natl Acad Sci U S A* **102**, 11064–11069.

Ivens, A. C., Peacock, C. S., Worthey, E. A. & 98 other authors (2005). The genome of the kinetoplastid parasite, *Leishmania major*. *Science* **309**, 436–442.

Kill, K., Binnewies, T. T., Sicheritz-Pontén, T., Willenbrock, H., Hallin, P. F., Wassenaar, T. M. & Ussery, D. W. (2005). Genome update: sigma factors in 240 bacterial genomes. *Microbiology* **151**, 3147–3150.

Margulies, M., Egholm, M., Altman, W. E. & 53 other authors (2005). Genome sequencing in microfabricated high-density picolitre reactors. *Nature* **437**, 376–380.

Méthé, B. A., Nelson, K. E., Deming, J. W. & 24 other authors (2005). The psychrophilic lifestyle as revealed by the genome sequence of *Colwellia psychrerythraea* 34H through genomic and proteomic analyses. *Proc Natl Acad Sci U S A* **102**, 10913–10918.

Pennisi, E. (2005). Biochemistry. Cut-rate genomes on the horizon? *Science* **309**, 862.

Shendure, J., Porreca, G. J., Reppas, N. B. & 7 other authors (2005). Accurate multiplex polony sequencing of an evolved bacterial genome. *Science* **309**, 1728–1732.

Van Domselaar, G. H., Stothard, P., Shrivastava, S. & 7 other authors (2005). BASys: a web server for automated bacterial genome annotation. *Nucleic Acids Res* **33**, W455–W459.

Vasconcelos, A. T., Ferreira, H. B., Bizarro, C. V. & 83 other authors (2005). Swine and poultry pathogens: the complete genome sequences of two strains of *Mycoplasma hyopneumoniae*

and a strain of *Mycoplasma synoviae*. *J Bacteriol* **187**, 5568–5577.

DOI 10.1099/mic.0.28423-0

Clinical significance of seeding dispersal in biofilms

We read with interest the recent paper by Purevdorj-Gage *et al.* (2005) on seeding dispersal in *Pseudomonas aeruginosa* biofilms. The authors compared a mucoid cystic fibrosis (CF) *P. aeruginosa* strain (strain FRD1) with strain PAO1 and found it to be seeding-dispersal-negative. They concluded that seeding dispersal might not be utilized by mucoid variants of CF strains, but rather be a transmission mechanism utilized by environmental strains of *P. aeruginosa*. We have been investigating mucoid CF isolates in a flow-through biofilm model (Webb *et al.*, 2003) and have some observations that are relevant to this conclusion. Our studies have shown that CF isolates can exhibit biofilm developmental processes and seeding dispersal similar to strain PAO1 in this experimental model. We found that CF *P. aeruginosa* isolates ($n=6$) each exhibited a characteristic pattern of biofilm development and microcolony formation that was reproducible and 'true to strain' in replicated biofilm experiments. Some CF strains exhibited a developmental pattern similar to that reported for strain FRD1, without obvious seeding dispersal within

the time-frame of our experiments (7 days). However, other strains did form hollow structures with highly motile cells in the centre and seeding dispersal events, as described for strain PAO1, after 4–5 days of culture (Fig. 1a, b). Clearly, much still remains to be understood about the mechanisms underlying seeding-dispersal and 'hollow-colony' formation. Previous studies using *P. aeruginosa* strain PAO1 linked this behaviour with bacteriophage-mediated lysis of a subpopulation of cells inside microcolony structures. We also observed dispersal-associated death in our CF isolates. BacLight LIVE/DEAD staining (Molecular Probes) of CF biofilms after day 6 of culture showed that all six strains tested exhibited regions of cell death within microcolonies. For at least one strain the pattern of cell death was identical to that seen with strain PAO1 (e.g. Fig. 1c). Coincident with this microcolony death, bacteriophage titres reached levels of $>10^7$ p.f.u. ml⁻¹ in flow-cell effluents. Our evidence to date thus suggests that death-associated dispersal mechanisms, as have been described for strain PAO1 (Webb *et al.*, 2003, 2004), are also central to CF strain transmission.

S. M. Kirov,¹ J. S. Webb²
and S. Kjelleberg²

¹School of Medicine, University of Tasmania, Private Bag 29, Hobart, Tasmania 7001, Australia

Fig. 1. Microcolonies of mucoid CF strains seen in a flow-cell model. (a) Sites of evacuation 'blebs' (arrows) from a hollow microcolony (strain U3A) after 4 days of culture. Bar, 46 µm. (b) A hollow microcolony (strain U3A) after 4 days of culture showing highly motile cells which appear blurred in the centre. Bar, 19 µm. (c) Cell death in the central region of a microcolony (strain 75) at ~7 days of culture (x - y plane view; BacLight LIVE/DEAD viability stain). Live cells fluoresce green; dead cells appear red in this confocal micrograph. Bar, 15 µm.

²School of Biotechnology and Biomolecular Sciences and Centre for Marine Biofouling and Bio-innovation, University of New South Wales, Sydney, NSW 2052, Australia

Correspondence: Sylvia M. Kirov (S.M.Kirov@utas.edu.au)

Purevdorj-Gage, B., Costerton, W. J. & Stoodley, P. (2005). Phenotypic differentiation and seeding dispersal in non-mucoid and mucoid *Pseudomonas aeruginosa* biofilms. *Microbiology* **145**, 1569–1576.

Webb, J. S., Thompson, L. S., James, S., Charlton, T., Tolker-Nielsen, T., Koch, B., Givskov, M. & Kjelleberg, S. (2003). Cell death in *Pseudomonas aeruginosa* biofilm development. *J Bacteriol* **185**, 4585–4592.

Webb, J. S., Lau, M. & Kjelleberg, S. (2004). Bacteriophage and phenotypic variation in *Pseudomonas aeruginosa* biofilm development. *J Bacteriol* **186**, 8066–8073.

DOI 10.1099/mic.0.28295-0

Clinical significance of seeding dispersal in biofilms: a response

We welcome the dialogue concerning the potential clinical significance of seeding dispersal (Purevdorj-Gage *et al.*, 2005) in the life cycle of mucoid *Pseudomonas aeruginosa* biofilms. We had based our hypothesis that the seeding dispersal phenomenon may be more relevant for non-mucoid, environmental strains on (1) the propensity of diseases associated with reduced mucociliary clearance in the lung, such as cystic fibrosis (CF) or chronic obstructive pulmonary disease (COPD), to select for mucoid *P. aeruginosa* phenotypes, and (2) that conversion to mucoidy is usually associated with a concomitant down-regulation of flagella production and loss of swimming motility (see Garrett *et al.*, 1999). However, as Kirov *et al.* discuss above, there may be greater diversity in mucoid CF isolates than generally acknowledged, and the perceived dichotomy between mucoidy and swimming phenotypes should be a topic for debate. More recently it has been shown that expression of flagellum genes in

response to oxygen limitation precedes loss of mucoidy and is reversible during this state of transition (Wyckoff *et al.*, 2002). Further, in an ongoing screen of mucoid CF isolates it was found that 6 out of 20 were motile (D. J. Wozniak, personal communication). However, it was suspected that growth medium may also play a role in the outcome of the motility assay and exact proportions may vary depending on culture conditions. The growth-condition-dependent transient switching between mucoid and swimming phenotypes is problematic when relating a particular biofilm behaviour with phenotype and stresses the importance of attempting to characterize the phenotypic state at various time points during biofilm development. Broadly though, the finding of Dr Wozniak is in agreement with that of Kirov *et al.* We agree that to fully assess the role of seeding dispersal (and other yet unidentified behavioural developmental phenotypes) in the context of lung infections, biofilm studies should include greater diversity in strains, growth conditions and be conducted over longer time scales. The interesting observation that seeding motility occurred in a mucoid CF isolate clearly demonstrates that the phenomenon may have clinical relevance.

Another point we would like to briefly address is the difference in terminology used by our two groups in reference to the hollowing out of clusters. Kirov *et al.* refer to hollowing resulting from ‘bacteriophage-mediated lysis’ as ‘death-associated dispersal’. We use the term ‘seeding dispersal’ in a more general sense to describe sequential steps from the formation of microcolonies culminating in the release of live cells from the centres of the colonies. Our working hypothesis for seeding dispersal which integrates observations from our two groups is thus: (1) the formation of cell clusters; (2) the differentiation of the cell cluster into two or possibly three distinct phenotypes consisting of the cells in the stationary cluster ‘wall’, the highly motile seeding subpopulation in the interior and the lysing interior subpopulation (Webb *et al.*, 2003) which allows the interior to liquefy; (3) highly agitated ‘seeding motility’ in the interior; and finally (4) the swimming, or swarming, of cells out of the microcolony. In the case of PAO1 the release appears to

occur through directed motility, but we recognize that hollowing in non-motile species such as *Staphylococcus epidermidis* (P. Stewart & B. Pitts, personal communication) and *Actinobacillus actinomycetemcomitans* (Kaplan *et al.*, 2003) may be driven by water currents. It is interesting to speculate that the recent increase in reports of hollow clusters occurring in a diversity of bacterial biofilms is evidence of convergent evolution in biofilm development.

P. Stoodley,¹ B. Purevdorj-Gage^{2†} and J. W. Costerton³

¹Center for Genomic Sciences, Allegheny-Singer Research Institute, Pittsburgh, USA

²Center for Biofilm Engineering and Dept of Microbiology, Montana State University-Bozeman, MT 59717, USA

³University of Southern California, Center For Biofilms, Los Angeles, CA, USA

†Present address: Division of Health Sciences, Montana State University-Bozeman, MT 59717, USA

Correspondence: B. Purevdorj-Gage (laura_p@erc.montana.edu)

Garrett, E. S., Perlegas, D. & Wozniak, D. J. (1999). Negative control of flagellum synthesis in *Pseudomonas aeruginosa* is modulated by the alternative sigma factor AlgT (AlgU). *J Bacteriol* **181**, 7401–7404.

Kaplan, J. B., Meyenhofer, M. F. & Fine, D. H. (2003). Biofilm growth and detachment of *Actinobacillus actinomycetemcomitans*. *J Bacteriol* **185**, 1399–1404.

Purevdorj-Gage, B., Costerton, W. J. & Stoodley, P. (2005). Phenotypic differentiation and seeding dispersal in non-mucoid and mucoid *Pseudomonas aeruginosa* biofilms. *Microbiology* **145**, 1569–1576.

Webb, J. S., Thompson, L. S., James, S., Charlton, T., Tolker-Nielsen, T., Koch, B., Givskov, M. & Kjelleberg, S. (2003). Cell death in *Pseudomonas aeruginosa* biofilm development. *J Bacteriol* **185**, 4585–4592.

Wyckoff, T. J. O., Thomas, B., Hassett, D. J. & Wozniak, D. J. (2002). Static growth of mucoid *Pseudomonas aeruginosa* selects for non-mucoid variants that have acquired flagellum-dependent motility. *Microbiology* **148**, 3423–3430.

DOI 10.1099/mic.0.28456-0