Mini-Review

Correspondence

B. C. Fries

fries@aecom.yu.edu

Phenotypic switching in *Cryptococcus neoformans*A. Guerrero,¹ N. Jain,² D. L. Goldman³ and B. C. Fries⁴^{1,2}Departments of Microbiology and Immunology¹ and Department of Microbiology² at All India Institute of Medical Sciences, New Delhi, India^{3,4}Department of Pediatrics³ and Medicine⁴, Albert Einstein College of Medicine, Bronx, NY 10461, USA

Phenotypic switching has been described in serotype A and D strains of *Cryptococcus neoformans*. It occurs *in vivo* during chronic infection and is associated with differential gene expression and changes in virulence. The switch involves changes in the polysaccharide capsule and cell wall that affect the yeast's ability to resist phagocytosis. In addition, the phenotypic switch variants elicit qualitatively different inflammatory responses in the host. In animal models of chronic cryptococcosis, the immune response of the host ultimately determines which of the switch variants are selected and maintained. The importance of phenotypic switching is further underscored by several findings that are relevant in the setting of human disease. These include the ability of the mucoid colony variant of RC-2 (RC-2 MC) but not the smooth variant (RC-2 SM) to promote increased intracerebral pressure in a rat model of cryptococcal meningitis. Furthermore, chemotherapeutic and immunological antifungal interventions can promote the selection of the RC-2 MC variant during chronic murine infection.

Introduction

Phenotypic switching enables micro-organisms to undergo rapid microevolution and to adapt to different micro-environments (Hammerschmidt *et al.*, 1996; Lysnynansky *et al.*, 1996; Schwan & Hinnebush, 1998; Silverman *et al.*, 1979; Slutsky *et al.*, 1985). Phenotypic switching has been observed in prokaryotic and eukaryotic human pathogens, including pathogenic fungi. For human pathogens the host represents a microenvironment of particular interest and phenotypic switching in this setting can affect the host–pathogen relationship with consequences that can translate into changes in virulence. An example of how phenotypic switching can alter the host–pathogen relationship is provided by *Trypanosoma cruzi*, where phenotypic switching generates antigenically different variants that can escape recognition by a specific antibody (Myler *et al.*, 1984). Other examples come from encapsulated bacteria, where phenotypic phase variation controls the expression of the capsule. This process is pivotal in determining whether the micro-organism is a commensal or an invasive pathogen (Swartley *et al.*, 1997). Because it is difficult to examine a pathogen population *in vivo*, studies addressing the phenotypic diversity of an isogenic population are difficult to undertake (Ebert, 1998). In this review, we will outline our findings and demonstrate why *Cryptococcus neoformans* represents an ideal model to examine the effects of phenotypic switching on host–pathogen interaction.

For fungi, phenotypic switching is an *in vitro* phenomenon defined as the spontaneous emergence of colonies with altered colony morphology at rates higher than the somatic

mutation rates (Soll, 1992). Macroscopic phenotypic switching involving changes in colony morphology is associated with a diverse array of changes on the cellular level. In contrast to mass conversion, phenotypic switching occurs only in a small proportion of the pathogen population and it represents one mechanism by which ‘phenotype heterogeneity’ can be achieved. Phenotypic switching is reversible and thus a controlled process, which is different from a random mutation. In fungi, switching frequencies range between 10^{-2} and 10^{-5} and are higher than the expected frequency of spontaneous mutations, which in eukaryotes is between 10^{-7} and 10^{-8} mutations per generation. Phenotypic switching was first described in *Candida albicans* 20 years ago (Slutsky *et al.*, 1985, 1987) and the underlying molecular mechanisms have been extensively studied (Miller & Johnson, 2002; Perez-Martin *et al.*, 1999). In more recent years phenotypic switching has also been demonstrated in other fungi including *Cryptococcus neoformans* (Eissenberg *et al.*, 1996; Fries *et al.*, 1999, 2001; Goldman *et al.*, 1998; Kugler *et al.*, 2000; Lachke *et al.*, 2000; Sinha *et al.*, 2000).

Phenotype variability in chronic cryptococcosis

Cryptococcus neoformans is a encapsulated yeast that causes life-threatening infections in patients with AIDS. Following the introduction of effective retroviral therapy in the United States, the annual incidence of cryptococcosis per 1000 persons with AIDS decreased approximately 10-fold, from 24–66 cases in 1992 to 2–7 cases in 2000 (Mirza *et al.*, 2003). In less fortunate parts of the world, however, chronic

cryptococcosis still constitutes a major opportunistic infection. In Africa, chronic *C. neoformans* affects up to 40 % of AIDS patients (Perfect & Casadevall, 2002). Chronic meningoencephalitis is the most common clinical manifestation of *C. neoformans* infection, and remains difficult to treat, despite antifungal therapy (Powderly, 2000). Lifelong suppressive antifungal therapy is recommended to reduce the likelihood of recurrent cryptococcosis in AIDS patients in whom the underlying HIV disease cannot be successfully suppressed. Treatment failures in patients with chronic cryptococcosis are generally not the result of acquired drug resistance but represent persistence of the initial strain (Brandt *et al.*, 1996, 2001; Pfaller *et al.*, 1999; Spitzer *et al.*, 1993).

The propensity of *C. neoformans* to undergo rapid changes is referred to as ‘microevolution’. Microevolution produces phenotypic variability that can alter the virulence of the pathogen. For the *C. neoformans* strain ATCC 24067 analysis of different isolates maintained in different laboratories revealed striking differences in phenotypic characteristics and virulence (Chen & Casadevall, 1999; Franzot *et al.*, 1998). Several observations support the concept that microevolution of *C. neoformans* occurs during the course of human infection and contributes to persistence of infection by challenging the already compromised host with variants that escape ongoing immune responses. Serial *C. neoformans* isolates from chronically infected HIV patients demonstrated differences in murine infection (Fries & Casadevall, 1998) as well as changes in the polysaccharide capsule (Cherniak *et al.*, 1995). In a similar fashion, investigations in chronic murine infection models have documented the emergence of both giant cells and poorly encapsulated forms (Feldmesser *et al.*, 2001). In addition to the emergence of variants that exhibit changes of the sterol content in the cell membrane (Currie *et al.*, 1995), the emergence of variants

with changes in colony morphology and karyotype pattern has been described (Fries *et al.*, 1996). Furthermore, analysis of the colony morphology in primary clinical specimens has shown that phenotypic variation in colony morphology can be observed in isolates from the spinal fluid (Fries *et al.*, 2005a). The exact frequency of phenotypic switching in clinical isolates is not known. In a recent analysis of *C. neoformans* strains from 39 patients in India we detected phenotypic switching from a smooth to a mucoid colony morphology in 2 out of the 39 patient isolates. There is no obvious association with mating type or serotype (Jain *et al.*, 2005). In summary, many studies demonstrate that phenotypic changes occur rapidly during chronic infection (Currie *et al.*, 1995; Fries *et al.*, 1996; Sukroongreung *et al.*, 2001). Phenotypic switching is one mechanism that can facilitate microevolution.

Phenotypic switching of *C. neoformans* strains in vitro

For *C. neoformans*, phenotypic switching has so far been described in serotype A (SB4, J32) and serotype D strains (24067a, RC-2) (summarized in Table 1). In these strains colonies with altered morphology arise spontaneously at a frequency of about 1 in 10⁻⁴ to 10⁻⁵ and reversion to the parent smooth colony type occurs at a comparable frequency (approx. 1 in 10⁻³ to 10⁻⁵). The smooth colonies (S and SM) of both SB4 and 24067 are round with a smooth dome surface and smooth edges. The mucoid (M and MC) colonies are round with smooth edges and a shiny and mucoid-appearing colony surface. Wrinkled (WR), serrated (C) and pseudohyphal (PH) colonies exhibit an irregular dome surface with or without serrated margins. Both smooth and mucoid colonies represent standard colony types common to many *C. neoformans* strains, whereas the WR, C and PH colony types of SB4 and 24067a are rarely observed in

Table 1. Characteristics of *C. neoformans* switching strains

Strain	Serotype	Phenotype	Switching frequency	Virulence	Inflammatory response
SB4	A	Smooth (S) Serrated (C) Wrinkled (WR)	2·2 × 10 ⁻³ –9 × 10 ⁻⁶	WR>S>C in mice	C>S>WR in rats*
J32	A	Smooth (NM) Muroid (M)	3·4 × 10 ⁻³ –4·1 × 10 ⁻⁵	ND	ND
24067a	D	Smooth (SM) Pseudohyphal (PH) Wrinkled (WR)	2·2 × 10 ⁻³ –<6 × 10 ⁻⁶	WR persists, SM and PH are cleared in rats	WR>PH>SM in rats†
RC-2	D	Smooth (SM) Muroid (MC)	10 ⁻⁴ –0·5 × 10 ⁻⁴	MC>SM in mice and rats	MC>SM in mice‡

*Minimal inflammation in WR-infected rats in contrast to intense caseous necrosis in C-infected rats.
†No inflammation in SM-infected rats in contrast to extensive granuloma formation in WR-infected rats.
‡Organized granuloma formation in SM-infected mice in contrast to extensive and destructive inflammatory response in MC-infected mice.

clinical isolates. This review will focus on RC-2 because to this date it is the most thoroughly examined switching system.

The MC colonies of RC-2 exhibit an indistinguishable colony micro-architecture when compared to the parent SM colony types. This colony phenotype is the result of excessive production of a viscous exopolysaccharide which is secreted on the colony surface, leading to the shiny appearance of these colonies (Fig. 1). In contrast, the micro-architecture of PH colonies exhibits changes due to altered cell morphology (e.g. pseudohyphal cells), and the microarchitecture of WR colonies is changed due to a change in the packing of cells. Similar changes are observed in the switching *Candida* strain 3153A (Radford *et al.*, 1994).

More detailed analysis of newly generated MC switch variants demonstrates that beyond the colony morphology other phenotypic characteristics are changed (Table 2). The MC switch variant grows more slowly and exhibits a larger polysaccharide capsule compared to the SM parent strain. In older MC colonies the capsular exopolysaccharide is abundantly shed. In addition, the MC switch variant of RC-2 exhibited increased sensitivity to lysing enzyme when compared to the SM parent.

Changes of the capsular polysaccharide associated with phenotypic switching

C. neoformans is a facultative intracellular pathogen and one unique characteristic is its thick polysaccharide capsule (Bulmer & Sans, 1968; Feldmesser *et al.*, 2000; Levitz *et al.*, 1999; Steenbergen *et al.*, 2001). This capsule prevents phagocytosis and rapid destruction within the intracellular milieu of macrophages. Glucuronoxylomannan (GXM) constitutes the major exopolysaccharide of the capsule; it is composed of (1→3)-linked linear α -D-mannopyranan with

Table 2. Phenotypic characteristics that are affected by phenotypic switching

	RC-2 SM	RC-2 MC
Doubling time at 30 °C	2.5 h	2.8 h
Doubling time at 37 °C	2.6 h	2.7 h
Cell size at 37 °C	5.9 ± 3	7.1 ± 0.79
Capsule size at 37 °C	1.7 ± 0.6	2.9 ± 0.54
Capsule induction in 5% CO ₂	6.2 ± 0.9	6.9 ± 0.5
Phagocytosis index by alveolar macrophages	39.16 ± 8.6	8.9 ± 2.0
Melanization	Yes	Yes
MIC amphotericin B	2	4
MIC fluconazole	2.5	1.25
GXM triad structure as determined by NMR	M1	M1
Elementary composition (C, O, H and N)	No difference	No difference
Viscosity of GXM	MC ≫ SM	MC ≫ SM
Cell charge	27.68 ± 2.44 mV	30.28 ± 3.6 mV
Concentration of lysing enzyme required for complete lysis	48 µg ml ⁻¹	12 µg ml ⁻¹

β -D-xylopyranosyl (Xylp) and β -D-glucopyranosyluronic acid (Glc pA) residues added to the mannose at various positions. Cherniak *et al.* (1998) defined six structural reporter groups (SRG) based on the amount of 2-O-linked, 4-O-linked Xylp residues and 2-O-linked Glc pA residues. The type and prevalence of the GXM SRG usually correlates with differences in serological reactivity of *C. neoformans* strains. Antigenic testing has been used to classify strains according to one of five serotypes (A, D, C, B, AD).

Analogous to other encapsulated pathogens, phenotypic switching in *C. neoformans* can alter the exopolysaccharide. We have demonstrated that phenotypic switching results in significant changes of the biochemical composition of GXM of C colonies of SB4, as well as WR and PH colonies of 24067a (Fries *et al.*, 1999). The GXMs of the C colony type are composed of mixtures of SRGs (M2 and M3 for C) whereas SB4 SM exhibits predominately SRG M2. In a similar fashion the PH and WR colonies of 24067a exhibit a mix of SRGs (M1 and M5) whereas the SM parents are predominately M1 and M2. The addition of a Xylp group at the 4-O position in M3 and M5 most likely requires a different enzyme than linkage to the 2-O position. Interestingly, M3 SRGs are traditionally thought to be present only in the GXM of *C. neoformans* var. *gattii* isolates (serotypes B and C) and not in GXM of *C. neoformans* var. *neoformans* isolates (serotypes A and D) (Fig. 2a).

Phenotypic switching can also alter the biophysical properties of GXM (Fries *et al.*, 2001). The MC GXM of the RC-2 strain is more viscous than the GXM of the SM parent.

Fig. 1. SM and MC colony types of RC-2.

Fig. 2. GXM differences associated with phenotypic switching. (a) The core repeating triad of GXM polymer. The SRG M1, M2, M3 and M5 differ in their Xylp residue content. M3 and M5 have Xylp residues also linked in the 4-O position. (b) Size-exclusion chromatography shows earlier elution of MC GXM (RC-2) compared with SM GXM (RC-2), and confirms differences of the two GXMs (reproduced with permission from Fries *et al.* 2001).

MC GXM spins down faster during ultracentrifugation and is eluted off a size-exclusion column earlier than SM GXM (Fig. 2b). By NMR analysis, however, the SM and MC GXM exhibit the same average biochemical repeat structure (Table 2). Thus at this point it is not known what mediates the apparent difference between SM and MC GXM. Most importantly, biological assays have established that the MC GXM is more potent in inhibiting phagocytosis by macrophages *in vitro* and *in vivo* and thus contributes to the enhanced virulence of the MC switch variant in the RC-2 strain (Fries *et al.*, 2001).

Differences in virulence of phenotypic switch variants and mechanism of enhanced virulence

In all the 'switching' *C. neoformans* strains studied so far, phenotypic switching results in variants that exhibit augmented virulence in murine or rat infection models as measured by organ fungal burden, inflammatory response and/or survival (Fig. 3). In the RC-2 switching system RC-2 MC elicits a strikingly different immune response when compared to RC-2 SM. In a pulmonary infection model in the mouse this strain elicits a more vigorous, macrophage- and neutrophil-dominated inflammatory response associated with altered cytokine and chemokine expression profiles. Specifically the chemokines MCP and MIP-1 α were upregulated in MC-infected mice whereas the cytokines IL-4, IL-10, IL-2 and TNF α were downregulated when compared to RC-2-SM-infected mice (Fries *et al.*, 2001). In contrast, RC-2 SM elicits an effective lymphocyte-dominated immune response that promotes clearance of the fungal infection. The damage-driven inflammatory response in MC-infected mice leads to lung damage and rapid demise (Fig. 3). It is of note that differences in cytokine response were also observed in human peripheral blood cells that were infected with MC and SM yeast cells (Pietrella *et al.*, 2003).

Phenotypic switching of *C. neoformans* *in vivo*

Despite an association with virulence, it is a challenge to demonstrate that switching actually occurs in the host during the infection because most experimental infections require infectious doses that are higher than the switching rate (10^{-2} to 10^{-5}). As a result it is difficult to exclude that the emergence of new phenotypes *in vivo* is not the result of *in vivo* selection of altered phenotypes, derived from a mixed inoculum (Ebert, 1998). However, the characteristics of *C. neoformans* as a pathogen include two features that allowed us to prove that switching occurred *in vivo*: (1) the ability to induce progressive infection with a relatively small inoculum; and (2) the fact that infections are often chronic and are lethal to the host only after a prolonged period of

Fig. 3. (a) Mice infected with SM survive longer than mice infected intratracheally with MC (left panel). (b) Histological analysis of lung tissue with a macrophage-specific antibody (Mac-3) at day 14 shows a macrophage-dominated inflammatory response in MC-compared to SM-infected lung tissue (reproduced with permission from Fries *et al.*, 2001). Sections stained with haematoxylin and eosin; bar, 500 μ m.

infection. By using a small inoculum (10^3 SM cells), infecting many mice simultaneously, and applying the Poisson formula, we were able to confidently exclude the possibility ($P=1.1 \times 10^{-12}$) of a mixed infection as the cause for recovery of MC colony types after infection with SM (Fries *et al.*, 2001). This result provided the first unequivocal proof that phenotypic switching occurs during the course of chronic cryptococcosis. In addition, careful analysis of primary *C. neoformans* isolates demonstrated that mucoid in addition to smooth colonies can be detected in primary spinal fluid specimens from humans (Fries *et al.*, 2005b).

The biological relevance of phenotypic switching during chronic infection

The biological relevance of phenotypic switching from SM to MC in chronic infection was established in three independent studies. First, we demonstrated that the occurrence of phenotypic switching is associated with lethal outcome in murine infection. For this experiment BALB/c mice ($n=108$) were infected with 10^4 SM cells, an inoculum that is generally cleared within 3 months by BALB/c mice. Mice autopsied before the onset of symptomatic disease (before day 36) and long-term survivors had significantly ($P<0.001$, ANOVA) fewer MC colonies than those autopsied at the time of death between day 42 and day 140 (Fries *et al.*, 2005a).

Second, we showed that antifungal treatment of the chronically infected host can promote the selection of the MC switch variants *in vivo*. Amphotericin B treatment reduced fungal burden less effectively in MC-infected than in SM-infected mice, and consequently resulted in a more pronounced prolongation of survival in SM-infected compared to MC-infected mice (20 versus 42 days, $P<0.05$). Administration of anti-capsular monoclonal antibody mediated better protection in SM-infected than MC-infected mice, although a protective effect was not consistently observed at all doses. Most interestingly, both antifungal drug therapy and administration of anti-capsular monoclonal antibody promoted the selection of MC variants in SM-infected mice, a phenomenon manifested by a statistically higher percentage of mucoid colonies in SM-infected mice compared to non-treated control mice (Fries *et al.*, 2005a). This finding suggests that both chemotherapeutic and immunological antifungal interventions may promote the selection of the more virulent mucoid variant, which could affect the outcome of infection in chronically infected hosts.

Third, the RC-2 MC but not the RC-2 SM variant is able to promote increased intracerebral pressure (ICP) in a rat model of cryptococcal meningitis. This finding is important because in human infection increased ICP is the leading cause of high morbidity and mortality. This complication is only seen in a subpopulation of patients, and is rather difficult to treat. Our studies suggest that *C. neoformans* strain characteristics that are altered by phenotypic switching can determine whether ICP develops during chronic infection (Fries *et al.*, 2005b). Thus, similar to *Candida* (Soll

et al., 1989; Soll, 2002; Vargas *et al.*, 2000), phenotypic switching in *C. neoformans* can contribute to virulence during chronic human infection.

Molecular mechanism of phenotypic switching

The molecular mechanisms mediating phenotypic switching in *C. neoformans* are currently not understood. Phenotypic switching in some microbes can be achieved by genetic changes that follow a Mendelian inheritance pattern, whereas for other microbes the mechanisms are epigenetic and inherited in a non-Mendelian fashion (e.g. silencing). In *Candida*, white–opaque switching and sexual mating are controlled by mating type locus homeodomain proteins. In other microbes a variety of mechanisms have been implicated in generating phenotypic variability within a population, including transposition of mobile sequences, silencing of gene expression, activation of mutator genes and infection with prions, which modify the fidelity of translation termination (Chao *et al.*, 1983; Deitsch *et al.*, 1997; Klar *et al.*, 2001; Lachke *et al.*, 2000; Martin *et al.*, 1993; Perez-Martin *et al.*, 1999; Srikantha *et al.*, 2001; True & Lindquist, 2000).

Although karyotype instability was observed in strains 24067a and SB4 (Fries *et al.*, 1999; Goldman *et al.*, 1998), similar to the switching *C. albicans* strain 3153A, it could not consistently be correlated with phenotypic variability and was not reversible (Perez-Martin *et al.*, 1999; Rustchenko-Bulgac, 1991; Soll, 1992). Investigations involving differential display of RC-2 SM and RC-2 MC mRNA demonstrated that phenotypic switching was associated with downregulation of genes in the MC switch variant relative to the SM switch variant. The function of the majority of these genes is unknown; however, some may represent immunogenic epitopes (Guerrero *et al.*, 2003). Future studies will be directed at determining the specific function of the individual genes.

In conclusion

The phenomenon of phenotypic switching has been described for many different types of pathogens, and it enables pathogens to adapt to a changing environment. Such environments include different hosts that will employ sophisticated immunological tools to rid themselves of the pathogen, and variable environmental niches where microbes can be subjected to many types of selection pressures. Virulence of a pathogen is dependent on the pathogen's interaction with the host; thus it may not be a fixed characteristic of the pathogen but the product of a complex interaction between the host and the pathogen (Casadevall & Pirofski, 1999, 2001). Changes of the host's immune competence as well as changes of the pathogen's virulence factors affect this complex relationship.

C. neoformans is an excellent model organism to study phenotypic switching and its impact on progression of chronic infection. Data from our laboratory demonstrated for the first time phenotypic switching in a fungus during

experimental infection (Fries *et al.*, 2001). The importance of this finding is underscored by the demonstration that the phenotypic switch changes the outcome of infection (D'Souza & Heitman, 2001) and that selection of switch variants is promoted in the setting of antifungal therapy. *C. neoformans* is a haploid fungus that exhibits an asexual reproduction mode during human infection. Molecular studies suggest that *C. neoformans* populations have a largely clonal population structure that may have limited options for rapid change in response to environmental stress. Similarly, the limited virulence traits of *C. neoformans* in the setting of chronic infection may require constant microevolution to evade the immune response. We propose that phenotypic switching allows this pathogen rapid change without the disadvantage of an increased mutation rate, thus avoiding the accumulation of disadvantageous mutations. Future studies will concentrate on elucidating the underlying mechanism for phenotypic switching, which may differ for individual strains.

Acknowledgements

This work was supported by R01 AI 59681 to B.C.F.; N.J. was supported by the AIDS International Training and Research Program (NID D43-TW01403) of the Albert Einstein College of Medicine.

References

- Brandt, M. E., Pfaller, M. A., Hajjeh, R. A., Graviss, E. A., Rees, J., Spitzer, E. D., Pinner, R. W. & Mayer, L. W. (1996). Molecular subtypes and antifungal susceptibilities of serial *Cryptococcus neoformans* isolates in human immunodeficiency virus-associated cryptococcosis. Cryptococcal Disease Active Surveillance Group. *J Infect Dis* **174**, 812–820.
- Brandt, M. E., Pfaller, M. A., Hajjeh, R. A., Hamill, R. J., Pappas, P. G., Reingold, A. L., Rimland, D. & Warnock, D. W. (2001). Trends in antifungal drug susceptibility of *Cryptococcus neoformans* isolates in the United States: 1992 to 1994 and 1996 to 1998. *Antimicrob Agents Chemother* **45**, 3065–3069.
- Bulmer, G. S. & Sans, M. D. (1968). *Cryptococcus neoformans*. III. Inhibition of phagocytosis. *J Bacteriol* **95**, 5–8.
- Casadevall, A. & Pirofski, L. A. (1999). Host-pathogen interactions: redefining the basic concepts of virulence and pathogenicity. *Infect Immun* **67**, 3703–3713.
- Casadevall, A. & Pirofski, L. (2001). Host-pathogen interactions: the attributes of virulence. *J Infect Dis* **184**, 337–344.
- Chao, L., Vargas, C., Spear, B. & Cox, E. (1983). Transposable elements as mutator genes in evolution. *Nature* **303**, 633–635.
- Chen, L. C. & Casadevall, A. (1999). Variants of a *Cryptococcus neoformans* strain elicit different inflammatory responses in mice. *Clin Diagn Lab Immunol* **6**, 266–268.
- Cherniak, R., Morris, L. C., Belay, T., Spitzer, E. D. & Casadevall, A. (1995). Variation in the structure of glucuronoxylomannan in isolates from patients with recurrent cryptococcal meningitis. *Infect Immun* **63**, 1899–1905.
- Cherniak, R., Valafar, H., Morris, L. & Valafar, F. (1998). *Cryptococcus neoformans* chemotyping by quantitative analysis of ^1H nuclear magnetic resonance spectra of glucuronoxylomannans with a computer-simulated artificial neural network. *Clin Diagn Lab Immunol* **5**, 146–159.
- Currie, B., Sanati, H., Ibrahim, A. S., Edwards, J. E., Jr, Casadevall, A. & Ghannoum, M. A. (1995). Sterol compositions and susceptibilities to amphotericin B of environmental *Cryptococcus neoformans* isolates are changed by murine passage. *Antimicrob Agents Chemother* **39**, 1934–1937.
- Deitsch, K. W., Moxon, E. R. & Wellems, T. E. (1997). Shared themes of antigenic variation and virulence in bacterial, protozoal, and fungal infections. *Microbiol Mol Biol Rev* **61**, 281–293.
- D'Souza, C. A. & Heitman, J. (2001). It infects me, it infects me not: phenotypic switching in the fungal pathogen *Cryptococcus neoformans*. *J Clin Invest* **108**, 1577–1578.
- Ebert, D. (1998). Experimental evolution of parasites. *Science* **282**, 1432–1435.
- Eissenberg, L. G., Poirier, S. & Goldman, W. E. (1996). Phenotypic variation and persistence of *Histoplasma capsulatum* yeasts in host cells. *Infect Immun* **64**, 5310–5314.
- Feldmesser, M., Kress, Y., Novikoff, P. & Casadevall, A. (2000). *Cryptococcus neoformans* is a facultative intracellular pathogen in murine pulmonary infection. *Infect Immun* **68**, 4225–4237.
- Feldmesser, M., Kress, Y. & Casadevall, A. (2001). Dynamic changes in the morphology of *Cryptococcus neoformans* during murine pulmonary infection. *Microbiology* **147**, 2355–2365.
- Franzot, S., Mukherjee, J., Cherniak, R., Chen, L., Hamdan, J. & Casadevall, A. (1998). Microevolution of a standard strain of *Cryptococcus neoformans* resulting in differences in virulence and other phenotypes. *Infect Immun* **66**, 89–97.
- Fries, B. C. & Casadevall, A. (1998). Serial isolates of *Cryptococcus neoformans* from patients with AIDS differ in virulence for mice. *J Infect Dis* **178**, 1761–1766.
- Fries, B. C., Chen, F., Currie, B. P. & Casadevall, A. (1996). Karyotype instability in *Cryptococcus neoformans* infection. *J Clin Microbiol* **34**, 1531–1534.
- Fries, B. C., Goldman, D. L., Cherniak, R., Ju, R. & Casadevall, A. (1999). Phenotypic switching in *Cryptococcus neoformans* results in changes in cellular morphology and glucuronoxylomannan structure. *Infect Immun* **67**, 6076–6083.
- Fries, B. C., Taborda, C. P., Serfass, E. & Casadevall, A. (2001). Phenotypic switching of *Cryptococcus neoformans* occurs in vivo and influences the outcome of infection. *J Clin Invest* **108**, 1639–1648.
- Fries, B. C., Cook, E., Wang, X. & Casadevall, A. (2005a). Effects of antifungal interventions on the outcome of experimental infections with phenotypic switch variants of *Cryptococcus neoformans*. *Antimicrob Agents Chemother* **49**, 350–357.
- Fries, B. C., Lee, S. C., Kennan, R., Zhao, W., Casadevall, A. & Goldman, D. L. (2005b). Phenotypic switching of *Cryptococcus neoformans* can produce variants that elicit increased intracranial pressure in a rat model of cryptococcal meningoencephalitis. *Infect Immun* **73**, 1779–1787.
- Goldman, D., Fries, B., Franzot, S., Montella, L. & Casadevall, A. (1998). Phenotypic switching in the human pathogenic fungus *Cryptococcus neoformans* is associated with changes in virulence and pulmonary inflammatory response in rodents. *Proc Natl Acad Sci U S A* **95**, 14967–14972.
- Guerrero, A., Jain, N., Cook, E., Casadevall, A. & Fries, B. (2003). Phenotypic switch variants of *Cryptococcus neoformans* differ in gene expression profile. In *103rd General Meeting of the American Society for Microbiology*, Washington DC, Abstract F39.
- Hammerschmidt, S., Hilse, R., van Putten, J., Gerady-Schahn, R., Unkmair, A. & Frosch, M. (1996). Modulation of cell surface sialic acid expression in *Neisseria meningitidis* via a transposable genetic element. *EMBO J* **15**, 192–198.

- Jain, N., Wickes, B. L., Keller, S. M., Fu, J., Casadevall, A., Jain, P., Ragan, M. A., Banerjee, U. & Fries, B. (1993). *J Clin Microbiol* **43**, 5733–5742.
- Klar, A. J., Srikantha, T. & Soll, D. R. (2001). A histone deacetylation inhibitor and mutant promote colony-type switching of the human pathogen *Candida albicans*. *Genetics* **158**, 919–924.
- Kugler, S., Schurtz Sebghati, T., Groppe Eissenberg, L. & Goldman, W. E. (2000). Phenotypic variation and intracellular parasitism by *Histoplasma capsulatum*. *Proc Natl Acad Sci U S A* **97**, 8794–8798.
- Lachke, S. A., Srikantha, T., Tsai, L. K., Daniels, K. & Soll, D. R. (2000). Phenotypic switching in *Candida glabrata* involves phase-specific regulation of the metallothionein gene MT-II and the newly discovered hemolysin gene HLP. *Infect Immun* **68**, 884–895.
- Levitz, S. M., Nong, S. H., Seetoo, K. F., Harrison, T. S., Speizer, R. A. & Simons, E. R. (1999). *Cryptococcus neoformans* resides in an acidic phagolysosome of human macrophages. *Infect Immun* **67**, 885–890.
- Lysnynansky, I., Rosengarten, R. & Yogev, D. (1996). Phenotypic switching of variable surface lipoproteins in *Mycoplasma bovis* involves high-frequency chromosomal rearrangement. *J Bacteriol* **178**, 5395–5401.
- Martin, D. W., Schurr, M. J., Mudd, M. H., Govan, J. R., Holloway, B. W. & Deretic, V. (1993). Mechanism of conversion to mucoidy in *Pseudomonas aeruginosa* infecting cystic fibrosis patients. *Proc Natl Acad Sci U S A* **90**, 8377–8381.
- Miller, M. G. & Johnson, A. D. (2002). White-opaque switching in *Candida albicans* is controlled by mating-type locus homeodomain proteins and allows efficient mating. *Cell* **110**, 293–302.
- Mirza, S. A., Phelan, M., Rimland, D. & 8 other authors (2003). The changing epidemiology of cryptococcosis: an update from population-based active surveillance in 2 large metropolitan areas, 1992–2000. *Clin Infect Dis* **36**, 789–794.
- Myler, P., Allison, J., Agabian, N. & Stuart, K. (1984). Antigenic variation in African trypanosomes by gene replacement or activation of alternate telomeres. *Cell* **39**, 203–211.
- Perez-Martin, J., Uria, J. A. & Johnson, A. D. (1999). Phenotypic switching in *Candida albicans* is controlled by a SIR2 gene. *EMBO J* **18**, 2580–2592.
- Perfect, J. R. & Casadevall, A. (2002). Cryptococcosis. *Infect Dis Clin North Am* **16**, 837–874, v–vi.
- Pfaller, M. A., Zhang, J., Messer, S. A., Brandt, M. E., Hajjeh, R. A., Jessup, C. J., Tumberland, M., Mbidde, E. K. & Ghannoum, M. A. (1999). In vitro activities of voriconazole, fluconazole, and itraconazole against 566 clinical isolates of *Cryptococcus neoformans* from the United States and Africa. *Antimicrob Agents Chemother* **43**, 169–171.
- Pietrella, D., Fries, B., Lupo, P., Bistoni, F., Casadevall, A. & Vecchiarelli, A. (2003). Phenotypic switching of *Cryptococcus neoformans* can influence the outcome of the human immune response. *Cell Microbiol* **5**, 513–522.
- Powderly, W. G. (2000). Current approach to the acute management of cryptococcal infections. *J Infect* **41**, 18–22.
- Radford, D., Challacombe, S. & Walter, J. (1994). A scanning electron microscopy investigation of the structure of colonies of different morphologies produced by phenotypic switching in *Candida albicans*. *J Med Microbiol* **40**, 416–423.
- Rustchenko-Bulgac, E. (1991). Variations of *Candida albicans* electrophoretic karyotypes. *J Bacteriology* **191**, 6586–6596.
- Schwan, T. G. & Hinnebusch, B. J. (1998). Bloodstream- versus tick-associated variants of a relapsing fever bacterium. *Science* **280**, 1938–1940.
- Silverman, M., Zieg, J., Hilmen, M. & Simon, M. (1979). Phase variation in *Salmonella*: genetic analysis of a recombinational switch. *Proc Natl Acad Sci U S A* **76**, 391–395.
- Sinha, H., Pain, A. & Johnstone, K. (2000). Analysis of the role of *recA* in phenotypic switching of *Pseudomonas tolaasii*. *J Bacteriol* **182**, 6532–6535.
- Slutsky, B., Buffo, J. & Soll, D. R. (1985). High-frequency switching of colony morphology in *Candida albicans*. *Science* **230**, 666–669.
- Slutsky, B., Staebell, M., Anderson, J., Risen, L., Pfaller, M. & Soll, D. R. (1987). “White-opaque transition”: a second high-frequency switching system in *Candida albicans*. *J Bacteriol* **169**, 189–197.
- Soll, D. R. (1992). High-frequency switching in *Candida albicans*. *Clin Microbiol Rev* **5**, 183–203.
- Soll, D. R. (2002). *Candida* commensalism and virulence: the evolution of phenotypic plasticity. *Acta Trop* **81**, 101–110.
- Soll, D., Galask, R., Isley, S., Rao, T., Stone, D., Hicks, J., Schmid, J., Mac, K. & Hanna, C. (1989). Switching of *Candida albicans* during successive episodes of recurrent vaginitis. *J Clin Microbiol* **27**, 681–690.
- Spitzer, E. D., Spitzer, S. G., Freundlich, L. F. & Casadevall, A. (1993). Persistence of initial infection in recurrent *Cryptococcus neoformans* meningitis. *Lancet* **341**, 595–596.
- Srikantha, T., Tsai, L., Daniels, K., Klar, A. J. & Soll, D. R. (2001). The histone deacetylase genes *HDA1* and *RPD3* play distinct roles in regulation of high-frequency phenotypic switching in *Candida albicans*. *J Bacteriol* **183**, 4614–4625.
- Steenbergen, J. N., Shuman, H. A. & Casadevall, A. (2001). *Cryptococcus neoformans* interactions with amoebae suggest an explanation for its virulence and intracellular pathogenic strategy in macrophages. *Proc Natl Acad Sci U S A* **98**, 15245–15250.
- Sukroongreung, S., Lim, S., Tantimavanich, S., Eampokalap, B., Carter, D., Nilakul, C., Kulkeratiyut, S. & Tansuphaswadikul, S. (2001). Phenotypic switching and genetic diversity of *Cryptococcus neoformans*. *J Clin Microbiol* **39**, 2060–2064.
- Swartley, L., Marfin, A., Edupuganti, S., Liu, L., Cieslak, P., Perkins, B. & Wenger, J. (1997). Capsule switching of *Neisseria meningitidis*. *Proc Natl Acad Sci U S A* **94**, 271–276.
- True, H. L. & Lindquist, S. L. (2000). A yeast prion provides a mechanism for genetic variation and phenotypic diversity. *Nature* **407**, 477–483.
- Vargas, K., Messer, S. A., Pfaller, M., Lockhart, S. R., Stapleton, J. T., Hellstein, J. & Soll, D. R. (2000). Elevated phenotypic switching and drug resistance of *Candida albicans* from human immunodeficiency virus-positive individuals prior to first thrush episode. *J Clin Microbiol* **38**, 3595–3607.