Phenotypic characterization of OmpX, an Ail homologue of *Yersinia pestis* KIM

Anna M. Kolodziejek, Dylan J. Sinclair, Keun S. Seo, Darren R. Schnider, Claudia F. Deobald, Harold N. Rohde, Austin K. Viall, Scott S. Minnich, Carolyn J. Hovde, Scott A. Minnich† and Gregory A. Bohach†

Correspondence
Gregory A. Bohach
gbohach@uidaho.edu

Department of Microbiology, Molecular Biology, and Biochemistry, University of Idaho, Moscow, ID 83844-3052, USA

The goal of this study was to characterize the *Yersinia pestis* KIM OmpX protein. *Yersinia* spp. provide a model for studying several virulence processes including attachment to, and internalization by, host cells. For *Yersinia enterocolitica* and *Yersinia pseudotuberculosis*, Ail, YadA and Inv, have been implicated in these processes. In *Y. pestis*, YadA and Inv are inactivated. Genomic analysis of two *Y. pestis* strains revealed four loci with sequence homology to Ail. One of these genes, designated y1324 in the *Y. pestis* KIM database, encodes a protein designated OmpX. The mature protein has a predicted molecular mass of 17.47 kDa, shares approximately 70% sequence identity with *Y. enterocolitica* Ail, and has an identical homologue, designated Ail, in the *Y. pestis* CO92 database. The present study compared the *Y. pestis* KIM6⁺ parental strain with a mutant derivative having an engineered disruption of the OmpX structural gene. The parental strain (and a merodiploid control strain) expressed OmpX at 28 and 37 °C, and the protein was detectable throughout all phases of growth. OmpX was required for efficient adherence to, and internalization by, cultured HEp-2 cell monolayers and conferred resistance to the bactericidal effect of human serum. Deletion of *ompX* resulted in a significantly reduced autoaggregation phenotype and loss of pellicle formation *in vitro*. These results suggest that *Y. pestis* OmpX shares functional homology with *Y. enterocolitica* Ail in adherence, internalization into epithelial cells and serum resistance.

Received 31 December 2006
Revised 18 May 2007
Accepted 21 May 2007

INTRODUCTION

Yersinia pestis, the agent of plague, probably evolved from *Yersinia pseudotuberculosis* relatively recently, resulting in three pandemics (Wren, 2003). Today, treatment and prevention are not yet 100% effective and a better understanding of pathogenesis is needed to enhance protection (McEvedy, 1988; Titball & Williamson, 2004; Zietz & Dunkelberg, 2004). *Y. pestis* carries a number of pathogenesis genes on the chromosome, but the pCD1 plasmid is essential for virulence. pCD1 encodes a type III secretion system and effector proteins with several functions, including circumventing host innate immunity (Leigh *et al.*, 2005; Lindler *et al.*, 1990).

Y. pestis enters epithelial cells, another mechanism to circumvent the immune response (Cowan *et al.*, 2000). In airborne disease, penetration of epithelial cells may promote development of pulmonary lesions (Liu *et al.*, 2006). Although Pla protease enhances interaction with epithelial cells (Cowan *et al.*, 2000; Lahteenmaki *et al.*,

1998, 2001a), Psa fimbriae have also been implicated. Neither is sufficient to confer complete internalization. Thus, *Y. pestis* adherence and internalization mechanisms are not yet completely elucidated (Liu *et al.*, 2006).

Outer-membrane proteins (Omps) have β -strand structures with membrane-spanning domains, and participate in channelling, antibiotic resistance and signal transduction (Bockmann & Caflisch, 2005). One Omp, OmpX, was first described for *Enterobacter cloacae* (Stoorvogel *et al.*, 1991), but homologues, including PagC, Lom, Rck and Ail (the attachment–invasion locus protein of *Yersinia* spp.), were identified in other Gram-negative bacteria (Dupont *et al.*, 2004; Heffernan *et al.*, 1992a; Meccas *et al.*, 1995). Proteins in this family promote invasion, resistance to complement-mediated killing, survival in macrophages, and internalization in epithelial cells (Cirillo *et al.*, 1996).

In the *Y. pestis* KIM genomic database, four ORFs for OmpX or Ail variants have been identified: y1324, y1682, y2034 (Caspi *et al.*, 2006; Karp *et al.*, 2005) and y2446 (UniProtKB/TrEMBL database). The y1324 gene encodes a protein which has a predicted molecular mass of 21 569 Da (including its signal sequence), has high sequence identity (99 and 68.5%) to Ail in *Y. pseudotuberculosis* and *Yersinia*

†These authors contributed equally to this work.

Abbreviations: C_T, threshold cycle; HIS, heat-inactivated serum; normal human serum.

enterocolitica, respectively, and is designated OmpX. The other three genes, y1682, y2034 and y2446 encode predicted proteins with 37.5, 46.4 and 45.9%, amino acid sequence identity, respectively, with *Y. enterocolitica* Ail.

Although *Y. enterocolitica* Ail has been extensively studied (Miller *et al.*, 1990, 2001), similar studies have not been reported for its *Y. pestis* homologues. Two previous studies showed that loss of an unidentified protein in the *Y. pestis* outer membrane affects autoaggregation (Podladchikova & Rykova, 2006) and adhesion to phagocytes (Kukleva *et al.*, 2000). The present study was conducted to characterize *Y. pestis* OmpX by engineering an OmpX deletion mutation in *Y. pestis* KIM6⁺ and comparing the properties of this mutant strain to the parental and complemented strains.

METHODS

Media, strains, and plasmids. Bacteria were cultured in Luria-Bertani (LB) low salt medium (EMD). Antibiotics were used at the following concentrations: nalidixic acid (Nal), 50 µg ml⁻¹; chloramphenicol (Cm), 30 µg ml⁻¹; and kanamycin (Kn), 50 µg ml⁻¹. LB agar with 5% sucrose and lacking NaCl was used to select for double-crossover recombinants, employing the *sacBR* locus. Congo red agar was used to confirm the pigmentation phenotype of the *Y. pestis* KIM6⁺ Nal^r strain (Surgalla & Beesley, 1969).

Strains and plasmids used in the study are listed in Table 1. A spontaneous Nal^r mutant was obtained by plating *Y. pestis* KIM6⁺ on LB containing Nal. The pMS20 suicide plasmid, with *sacBR* encoding levansucrase, and Cm^r, was used as described previously (Gay *et al.*, 1983). In some experiments, *Y. pestis* strains constitutively expressing green fluorescent protein (GFP) were used. These were constructed by electroporating pFVP25.1 (a gift from G. Mallo; *Caenorhabditis* Genetics Center, Minneapolis, MN; <http://www.cbs.umn.edu/CGC>), which encodes Amp^r and GFP.

HEp-2 cells (ATCC CCL-23) (Dziewanowska *et al.*, 2000) were grown in 6% CO₂ (37 °C) in growth medium (GM) [low glucose Dulbecco's modified Eagle's medium (Gibco) supplemented with 10% (v/v) fetal bovine serum (FBS) (HighClone) and 1% (v/v) penicillin/streptomycin solution (Gibco)].

Construction of a *Y. pestis* KIM6⁺ Nal^r *ompX* deletion mutant.

The *ompX* gene was amplified by PCR using primers complementary to the predicted promoter and terminator regions and containing engineered *Xba*I and *Eco*RV sites, respectively. The resulting 750 bp PCR product was cloned into pMS20 (Smith, 2000) and the construct (pMHZ1) was transformed into *Escherichia coli* CC118 λ pir. A representative clone was digested with *M*feI and *N*deI (sites in *ompX*), generating a 426 bp deletion. A gene conferring Kn^r (neomycin phosphotransferase; *npt*), with flanking FRT (flippase recognition target) sites, was amplified by PCR using pKD4 (Datsenko & Wanner, 2000) as a template and, and cloned into pMHZ1. The resulting construct, pMHZ2 carrying Kn^r, was transformed into *E. coli* S17-1 λ pir. A transformant harbouring pMHZ2 was mated with *Y. pestis* KIM6⁺ Nal^r as described previously (Smith, 2000) and counter-selected on LB agar with Nal and Cm. This merodiploid strain (*ompX*⁺/*ompX*::*npt*) was generated by a homologous single-crossover recombination. It was maintained and served as (i) a single copy *ompX* complementation control and (ii) an isogenic precursor for selecting the *ompX*::*npt* disruption. The latter was isolated on LB agar containing sucrose to select for a second crossover event while maintaining selection for the *ompX*::*npt* disruption. Sucrose-resistant, Cm-sensitive, colonies were tested by PCR for the

ompX::*npt* disruption (generation of a 1828 bp product). DNA sequencing was used to confirm all constructs.

PCR. Primers (Table 1) were purchased from Integrated DNA Technologies and reagents were from Invitrogen. Primers used to amplify Tn5 *npt* contained engineered *M*feI and *N*deI restriction sites. In the reverse primer, a mutation was introduced to exclude the *N*deI restriction site naturally present in the FRT region. Amplification of this region, including the FRT sequences, enables deletion of the antibiotic resistance gene using the flippase-bearing pCP20 plasmid (Datsenko & Wanner, 2000). PCR products were visualized on agarose gels and sizes were determined using the 1 kb Plus DNA ladder (Fisher Scientific).

Expression of *ompX* and the downstream (y1325) *hcaT* gene were measured by real-time PCR. Total RNA was extracted as described previously (Rebeil *et al.*, 2006). Reverse transcription with hexanucleotides (Roche), real-time PCR using the SYBR green I dye master mix and an ABI 7000 thermocycler (Perkin-Elmer Applied Biosystems), and data analysis were as described previously (Seo *et al.*, 2007). Relative quantities of mRNA were normalized to the amount of *proS* mRNA in the samples. RNA isolation for each strain was done in duplicate on two different days and real-time PCR was performed in triplicate for each RNA sample.

Culture growth measurements. Growth rate comparisons were done in LB broth (50 ml) at 28 and 37 °C on an orbital shaker (200 r.p.m.). A Beckman Coulter DU530 spectrophotometer (Beckman Instruments) was used to measure OD₆₀₀ of the cultures hourly, with vigorous vortexing before each reading.

Internalization assays. *Y. pestis* cells from ~12 h (OD₆₀₀ 0.55–0.75) cultures grown at 28 °C (200 r.p.m.) in LB medium were washed in PBS (0.01 M sodium phosphate, 0.8% NaCl, pH 7.2) and resuspended in internalization medium (IM; GM lacking FBS and antibiotics). Dilutions were made in PBS containing 0.1% Triton X-100 and 0.2% glycerol to determine cell numbers by plate counts. HEp-2 cells (1.5 × 10⁵ per well) in GM were incubated in 24-well plates (6% CO₂, 37 °C). After 42 h, the cell monolayers were washed three times with IM and 1 ml adjusted *Y. pestis* suspension was added to each well to produce co-cultures with m.o.i. values between 10 and 20. The plates were centrifuged (5 min, 200 g, 18 °C) and incubated for 1 h as described above. Each well was washed three times with IM and the co-cultures were incubated for 1.5 h in IM with 500 µg gentamicin per well (Gibco) to kill extracellular *Y. pestis*. The wells were washed three times and trypsin-like enzyme (Tryp-Le Express, Gibco) was added and incubated for 7 min to detach the HEp-2 cells. Triton X-100 (0.025%) was added to release intracellular bacteria and bacterial numbers were determined from plate counts. Initial experiments, using standard techniques (Shaffer & Goldin, 1974), showed that inactivating *ompX* did not significantly alter the gentamicin minimal bactericidal concentration (MBC; 3.9–7.8 µg ml⁻¹).

Cell-association assays. Procedures were conducted as described above except that, after 1 h incubation, the HEp-2/*Y. pestis* co-cultures were washed nine times to remove unbound bacteria and the gentamicin exposure step was omitted. In some experiments, microscopy was used to assess cell association of *Y. pestis* strains expressing GFP from plasmid pFVP25.1. HEp-2 cells were incubated for 48 h in a four-chamber coverglass (Nalge Nunc International) starting with an initial inoculum of 5.0 × 10⁴ bacteria ml⁻¹. The assays were performed as described above except that after the ninth wash the cells were fixed with 3.7% formaldehyde and stained with 16.5 nM phalloidin conjugated with Alexa Fluor 546 as described by the manufacturer (Invitrogen). Images of nine 0.7 µm-thick slices collected by LSM5 Pa laser scanning microscope (Carl Zeiss MicroImaging) with a Plan-Apochromat 63 ×/1.4 Oil DIC lens were processed by Zeiss LSM Image Examiner version 3.2.0.70.

Table 1. Strains, plasmids and PCR primers

Strain or plasmid	Relevant genotype	Reference or source
Strains		
<i>E. coli</i> CC118 λ pir	R ⁻ M ⁺ λ pir ⁺	de Lorenzo <i>et al.</i> (1990)
<i>E. coli</i> S17-1 λ pir	Δ recA RP4 2-Tc::Mu-Kn::Tn7 λ pir ⁺ tra ⁺ ; Tp ^r Str ^r	de Lorenzo <i>et al.</i> (1990)
<i>Y. pestis</i> KIM6 ⁺	pgm ⁺ pYV ⁻ pMT1	S. C. Straley, U. Kentucky
<i>Y. pestis</i> KIM6 ⁻	pgm ⁻ hmsT ⁺ pYV ⁻ pMT1	S. C. Straley, U. Kentucky
<i>Y. pestis</i> KIM6 ⁺ Nal ^r	pgm ⁺ pYV ⁻ pMT1 Nal ^r from <i>Y. pestis</i> KIM6 ⁺	This study
<i>Y. pestis</i> KIM6 ⁺ Nal ^r Δ ompX::npt	pgm ⁺ pYV ⁻ Nal ^r Δ ompX::npt	This study
<i>Y. pestis</i> KIM6 ⁺ Nal ^r ompX ⁺ /ompX::npt	pgm ⁺ pYV ⁻ Nal ^r ompX ⁺ with integrated pMHZ2; merodiploid for ompX	This study
<i>Y. pestis</i> KIM6 ⁺ Nal ^r pFVP25.1	Continuously expresses GFP	This study
<i>Y. pestis</i> KIM6 ⁺ Nal ^r Δ ompX::npt pFVP25.1	Continuously expresses GFP	This study
<i>Y. pestis</i> KIM6 ⁺ Nal ^r ompX ⁺ /ompX::npt pFVP25.1	Continuously expresses GFP	This study
Plasmids		
pKD4	Amp ^r ; containing Kn ^r gene flanked by FRT sites	Datsenko & Wanner (2000)
pMS20	mob ⁺ , pir-dependent oriR6K, sacBR Cm ^R	Smith (2000)
pMHZ1	pMS20 containing y1324 ompX gene from <i>Y. pestis</i>	This study
pMHZ2	pMS20 containing Δ ompX::npt flanked by FRT sites	This study
pFVP25.1	Amp ^r ; encodes GFP	G. Mallo <i>Caenorhabditis</i> Genetics Center
PCR primers: application and characteristics		
Primer sequence (F, forward; R, reverse)		
Amplification of y1324 ompX gene	5'-CGCGTCTAGATCATGTGTGATATTTG-3' (F) 5'-GCGCGATATCTAGCCTACCCCTATTA-3' (R)	This study
Disruption of ompX by insertion of npt cassette	5'-ACACGCCAATTGTTGTGTAGGCTGGAG-3' (F) 5'-TATGTGCATATGATGGGAATTAGCCATGGTCCTCAATATCC-3' (R)	This study
Confirmation of the mutation: flank ompX 200 bp 5' and 3' from the predicted ORF	5'-GCGCGATATCAACATTATCTGGTCCG-3' (F) 5'-GCGCGGATCCTATTCATTGGTGACTTG-3' (R)	This study
Detection of hmsT gene: predicted product length 639 bp	5'-TGATTGCGGTTATAAGCTTGAGC-3' (F) 5'-GGCCATAGTTGTCGTTATAGGCTT-3' (R)	This study
Detection of hmsT gene: predicted product length 1134 bp	5'-GTCTACTGACAGCACGATATTATGCAG-3' (F) 5'-ATTTCTCTTGGCCCGGTAGAG-3' (R)	This study
Real-time PCR; ompX gene	5'-CATTTGCTCAGACTCGCCG-3' (F) 5'-AGATGTTTTACTTCTACTCTCTGATTGACC-3' (R)	This study
Real-time PCR; hcaT gene	5'-TTTACTCTCCGCTTCCTTGG-3' (F) 5'-CCAAAGCATCCGTCAATGGT-3' (R)	This study
Real-time PCR; proS gene	5'-GTGGTCATAAGTTGGTTGCGC-3' (F) 5'-GCACCGAAATCGCTCATCA-3' (R)	This study

Analysis of OmpX expression. To examine OmpX levels at various growth phases and temperatures, bacteria were grown for 12 h as described above, washed in PBS, and pelleted by centrifugation at 12 000 g at 4 °C for 5 min. Proteins were extracted with a buffer containing 8 M urea (Sigma), 2 % CHAPS (EMD Chemicals), 20 mM DTT (Bio-Rad), bromophenol blue, Protease Inhibitors (Amersham Biosciences), and 100 × Nuclease (Amersham Biosciences). Protein samples and the Benchmark protein ladder (Invitrogen) were resolved by SDS-PAGE (Laemmli, 1970) on 12.5 % polyacrylamide gels and visualized by Coomassie blue staining.

Tandem mass spectrometry (MS/MS) analysis. Protein bands observed in SDS-PAGE gels were excised manually, destained, and subjected to standard protease digestion procedures (Shevchenko *et al.*, 1996). Samples were processed in high-recovery tubes from Axygen. Proteins were digested overnight at 37 °C using trypsin (Worthington) and 0.1 % n-octyl β -glucoside (Fluka) (Katayama *et al.*, 2001). Peptides were recovered with extraction buffer containing 50 % acetonitrile and 5 % trifluoroacetic acid and the resulting sample was concentrated under vacuum and resuspended in 5 % acetonitrile/0.1 % formic acid.

ProteinLynx Global Server 2.2 and Protein Expression Informatics System software Version 1.0 were used for MS/MS spectra analysis, peptide sequencing and protein identification. MS/MS data were compared to protein sequence databases from the University of Wisconsin (<http://www.genome.wisc.edu/sequencing/pestis.htm>) and the Sanger Institute (http://www.sanger.ac.uk/Projects/Y_pestis/) and results were analysed using Mascot software (Matrix Science).

Statistical analysis. Data were analysed using either the Mann–Whitney or Student’s *t* tests. These analyses were conducted with SigmaPlot 9.0 (Systat Software) software.

Bioinformatic analysis and generation of a *Y. pestis* KIM6⁺ OmpX deletion mutant

(*Y. pestis* KIM) beginning with GTG, and with a predicted 38 residue signal peptide (Fig. 1). Ail (Sanger Institute; *Y. pestis* CO92) begins with ATG and encodes a putative 26-residue signal peptide. ORF Finder (NCBI Tools for Bioinformatics Research) predicted the signal peptide sequence in the *Y. pestis* CO92 database for both genes.

Real-time PCR data indicated that *ompX* transcription was not significantly different in the parental and complemented strains (mean change in threshold cycle (ΔC_T): -6.550 and -6.285 ; $P=0.182$), and expression by the *Y. pestis* *ompX::npt* deletion mutant was undetectable. Furthermore, expression of the downstream *y1325* (*hcaT*) gene was not significantly different in any strain [mean ΔC_T :

8.16 (parental), 8.82 (deletion), 8.22 (merodiploid); $P=0.279$ (parental vs deletion), and $P=0.907$ (parental vs merodiploid)].

Proteomic analysis

SDS-PAGE analysis of *Y. pestis* whole-cell lysates (Fig. 2) indicated that the *Y. pestis* KIM6⁺ Nal^r $\Delta ompX::npt$ strain lacked a protein band at approx. 19 kDa which was present in parental *Y. pestis* KIM6⁺ Nal^r, control merodiploid *Y. pestis* KIM6⁺ Nal^r $ompX^+/ompX::npt$, and *Y. pestis* KIM6⁻ strains. This protein was expressed at both 28 and 37 °C and during exponential and stationary phases. This protein band from gels containing the parental strain, and corresponding gel slices from the mutants, was excised, digested with trypsin, and sequenced. MS/MS data were used to search protein databases for *Y. pestis* KIM and *Y. pestis* CO92. This revealed that the missing protein was identical to the mature form of OmpX (gene tag y1324) and Ail (gene tag ypo2905) in these two databases, respectively. In this report, we used the nomenclature in the *Y. pestis* KIM database (OmpX rather than Ail) to be

consistent with the gene designation used in the database for the strain most closely related to the strain used in our study.

Sequence coverage by ion mapping started at a putative signal peptide cleavage site predicted with SignalP 3.0 Server (Bendtsen *et al.*, 2004; Nielsen & Krogh, 1998) (Fig. 1). This suggests that the signal peptide is cleaved between Ala and Glu (Fig. 1); *Y. enterocolitica* Ail is cleaved at an analogous site (Miller *et al.*, 1990). The predicted molecular mass of mature OmpX, calculated with ProtParameters (Gasteiger *et al.*, 2005), is 17.47 kDa, slightly smaller than the ~19 kDa size estimated by SDS-PAGE.

Disruption of *ompX* does not impair the growth of *Y. pestis* in vitro

To ascertain whether the mutation affected bacterial growth rate, growth kinetics were monitored for the parental strain and its mutant derivatives at 28 and 37 °C. The *ompX* disruption did not impair growth at either temperature. In fact, the *Y. pestis* KIM6⁺ Nal^r $\Delta ompX::npt$ deletion mutant had a small but reproducible increase in growth rate at 28 °C compared to the parental and merodiploid control strains (doubling times 97, 101 and 107 min respectively; data not shown). This increased growth rate was more prominent at 37 °C; the deletion mutant had a doubling time of 118 min compared to 152 and 154 min, respectively, for the parental and merodiploid control strains. When compared to the parental and control strains, the increase in growth rate was statistically significant ($P<0.009$) for cultures grown at 37 °C, but not for those grown at 28 °C.

Disruption of *ompX* results in loss of autoaggregation and pellicle formation

Certain strains of *Y. pestis* produce characteristic flocculent growth in broth cultures (Bobrov *et al.*, 2002). This phenotype was readily observable for the parental *Y. pestis* KIM6⁺ Nal^r strain, the merodiploid *Y. pestis* KIM6⁺ Nal^r $ompX^+/ompX::npt$ control strain, and the *pgm*-deficient strain *Y. pestis* KIM6⁻ (Fig. 3a), which autoaggregated when grown at either 28 or 37 °C. Despite being subjected to shaking, the bacterial growth settled to the bottom when grown at 28 °C. Smaller aggregates formed but remained dispersed in cultures grown at 37 °C. In addition, these strains formed pellicles attaching to the sides of the tubes at the air-liquid interface. Loss of these characteristics coincided with disruption of the *ompX* gene. *Y. pestis* KIM6⁺ Nal^r $\Delta ompX::npt$ grew as a homogeneous suspension at both 28 and 37 °C, and neither clumping nor pellicle formation was observed at either temperature. Microscopic analysis revealed that the parental and merodiploid strains formed large aggregates, while the *Y. pestis* KIM6⁺ Nal^r $\Delta ompX::npt$ mutant grew as individual cells or in smaller clusters (Fig. 3b).

Fig. 2. SDS-PAGE of *Y. pestis* whole-cell lysate proteins. Cultures were grown at either 28 or 37 °C and collected during (a) mid-exponential phase (OD_{600} 0.7–0.8) or (b) near the start of the stationary (OD_{600} 2.1–2.3) phase of growth. (lanes 1, *Y. pestis* KIM6⁺ Nal^r; lanes 2, *Y. pestis* KIM6⁺ Nal^r $\Delta ompX::npt$; lanes 3, *Y. pestis* KIM6⁺ Nal^r $ompX^+/ompX::npt$; lanes 4: *Y. pestis* KIM6⁻). The lane labelled BM contains BenchMark pre-stained protein standard. The gel shows results from a representative experiment that was performed twice.

Fig. 3. Changes in autoaggregation and pellicle formation induced by disruption of the *ompX* gene. Bacteria were incubated in LB broth at 28 or 37 °C with vigorous shaking (200 r.p.m.) and collected during mid-exponential phase. (a) Macroscopic observation of cultures demonstrating autoaggregation and pellicle formation (indicated by arrows). A, *Y. pestis* KIM6⁺ Nal^r; B, *Y. pestis* KIM6⁺ Nal^r Δ *ompX*::*npt*; C, *Y. pestis* KIM6⁺ Nal^r, *ompX*⁺/*ompX*::*npt*; D, *Y. pestis* KIM6⁻. Note: Pellicle formation at 28 °C is very light and not easily visible or photographed in cultures collected during exponential growth. (b) Microscopic observation (taken at $\times 600$ magnification) of crystal-violet-stained cells confirming loss of autoaggregation upon deletion of the *ompX* gene.

OmpX protein and *Y. pestis* interactions with human epithelial cells

It was previously reported that two OmpX homologues, Ail and Rck in *Y. enterocolitica* and *Salmonella typhimurium*, respectively, are involved in adherence to and internalization by epithelial cells (Cirillo *et al.*, 1996; Heffernan *et al.*, 1994; Miller & Falkow, 1988). To determine whether *Y. pestis* OmpX performs similar functions, we examined the effect of inactivating the *ompX* gene on interactions with cultured HEp-2 human epithelial cells. The overall association of *Y. pestis* KIM6⁺ Nal^r Δ *ompX*::*npt* with HEp-2 cells was significantly impaired in comparison to *Y. pestis* KIM6⁺ Nal^r and *Y. pestis* KIM6⁺ Nal^r *ompX*⁺/*ompX*::*npt* (Fig. 4). Microscopic observation of *Y. pestis* KIM6⁺ Nal^r Δ *ompX*::*npt*-HEp-2 co-cultures revealed that very few bacterial cells were associated with the HEp-2 cells. In contrast, the parental and merodiploid control strains of *Y. pestis*, both of which express OmpX, were observed predominantly as large aggregates associated with HEp-2 cell monolayers. Bacterial cell counts from the cell-association assays were consistent with microscopic observation. *Y. pestis* KIM6⁺ Nal^r Δ *ompX*::*npt* cell numbers associated with HEp-2 cells (representing both adherent and internalized cells) were reduced by $\sim 90\%$ (11-fold reduction) compared to the parental strain and merodiploid control. This led to an $\sim 98\%$ (65.5-fold) reduction in internalization of *Y. pestis*. The data indicated

that internalization was more severely affected than adherence. Specifically, only 6% of the cell-associated *Y. pestis* KIM6⁺ Nal^r Δ *ompX*::*npt* were internalized by eukaryotic cells vs 35.75 and 32% for *Y. pestis* KIM6⁺ Nal^r and *Y. pestis* KIM6⁺ Nal^r *ompX*⁺/*ompX*::*npt*, respectively.

OmpX confers resistance to human serum

To examine whether *Y. pestis* OmpX confers resistance to killing by complement in human serum, a property of *Y. enterocolitica* and *Y. pseudotuberculosis* Ail (Bliska & Falkow, 1992; Yang *et al.*, 1996), standard assay protocols were used. The *ompX* deletion mutant was significantly more susceptible to complement-mediated killing compared to the parental and control strains (Fig. 5) grown at both 28 and 37 °C. Incubation in 50% NHS at 37 °C was essentially 100% lethal to *Y. pestis* KIM6⁺ Nal^r *ompX*::*npt* after 1 h. In contrast, *Y. pestis* KIM6⁺ Nal^r and *Y. pestis* KIM6⁺ Nal^r *ompX*⁺/*ompX*::*npt* expressing OmpX both retained complete viability during this time period. To exclude the possibility that the autoaggregation phenotype of the parental strain may significantly distort the complement-resistance results by protecting a fraction of the bacterial population from complement by making the bacterial cells inaccessible, we observed that during longer incubation with serum, bacterial aggregates were being

Fig. 4. The OmpX protein in *Y. pestis* KIM6⁺ Nal^r is required for optimal interaction with HEp-2 cells. (a) Cell-associated GFP-expressing *Y. pestis* bacteria after 1 h incubation and exhaustive washing. *Y. pestis* KIM6⁺ Nal^r ΔompX::npt pFVP25.1 (B) is visible as mostly individual cells associated with HEp-2 cells. In contrast, *Y. pestis* KIM6⁺ Nal^r pFVP25.1 (A) and *Y. pestis* KIM6⁺ Nal^r ompX⁺/ompX::npt pFVP25.1 (C) associated with the HEp-2 cells exist in bacterial aggregates and more bacteria are cell-associated compared to the ompX deletion strain. D, HEp-2 cell monolayers incubated without bacteria (taken at ×634 magnification; bar, 10 μm). (b) Quantification of cell association and internalization by: (1) *Y. pestis* KIM6⁺ Nal^r, (2) *Y. pestis* KIM6⁺ Nal^r ΔompX::npt and (3) *Y. pestis* KIM6⁺ Nal^r ompX⁺/ompX::npt. Results shown are mean ± SEM from data derived from two assays performed in duplicate (n=8) on two separate days. The asterisks (**) indicate statistically significant differences between the deletion mutant, and the parental and control strains (P<0.001).

disrupted and after 4 h the cells were more separated (data obtained from confocal microscopy of GFP-labelled bacteria; not shown). Incubation of the bacteria with serum for 4 h and subsequent addition of fresh serum did not result in any decrease in viability, which indicates that it is OmpX, not autoaggregation, which is the major factor in serum resistance.

DISCUSSION

This study characterized *Y. pestis* KIM6⁺ OmpX by inactivating the *ompX* gene and assessing the effects of the mutation on phenotypes associated with homologues in other Gram-negative organisms. We determined that *ompX* endows *Y. pestis* with the abilities to (i) efficiently associate with and become internalized by epithelial cells, (ii) resist the bactericidal activity of human serum, and (iii) promote autoaggregation.

In conducting this study we searched the *Y. pestis* KIM and CO92 databases and encountered conflicting nomenclature; the protein in the KIM6 database was designated OmpX, whereas the homologue in the CO92 strain was designated Ail. Although an Ail designation could be appropriate, because our work was conducted with *Y. pestis* KIM6⁺, we used the OmpX designation in this report to be consistent with the current database designation. Furthermore, *Y. pestis* has three additional *ail*-like genes with varying levels of predicted sequence similarity, and to designate *ompX* as *ail* would require a further qualifier as to which *ail* homologue was being addressed. Until a unified nomenclature is assigned, we have used *ompX* to limit potential confusion. However, we recommend that a standard nomenclature be assigned through future research and discussions.

Inactivation of *ompX* reduced epithelial cell association and internalization ~90 and ~98 %, respectively. At present we cannot distinguish whether the increased cell association of *Y. pestis* expressing OmpX is due directly to the effects of OmpX or indirectly due to OmpX-mediated autoaggregation. The residual cell association observed for the *ompX* deletion mutant is consistent with evidence suggesting that *Y. pestis* interaction with non-professional phagocytes is multifactorial. Kukleva *et al.* (2000) noted that a 22 kDa protein in the *Y. pestis* outer membrane promoted adherence to phagocytes; the protein in their study was not identified, nor were its effects on non-professional phagocytes analysed. Other proteins are involved in adhesion and invasion by *Y. pestis* (Cowan *et al.*, 2000; Lahteenmaki *et al.*, 1998, 2001a, b, 2003; Leigh *et al.*, 2005; Liu *et al.*, 2006). The role of the pCP1 plasmid, encoding the plasminogen activator (Pla protease), in interaction with HeLa cells has been reported (Cowan *et al.*, 2000). Although the reduced cell association and internalization resulting from pCP1 curing was less than that caused by inactivating *ompX* in *Y. pestis* KIM6⁺, one cannot make direct comparisons since different cell culture systems and

Fig. 5. OmpX promotes resistance to human serum bactericidal activity when the bacteria are grown at either 28 °C (a) or 37 °C (b). Bacteria were incubated in 50 % NHS or HIS. ▲, *Y. pestis* KIM6⁺ Nal^I; ●, *Y. pestis* KIM6⁺ Nal^I ΔompX::npt; ■, *Y. pestis* KIM6⁺ Nal^I ompX⁺/ompX::npt. The number of surviving bacteria after incubation with NHS is presented as a percentage of the number of bacteria incubated in HIS (100 %). Data in parentheses represent bacterial cell numbers for each strain (×10⁵) following incubation in HIS for 0, 0.5, and 1 h respectively. These values represent 100 % survival in HIS at each time point. Results are the mean ± SEM from data derived from two assays performed on separate days (n=6). * and ** indicate statistically significant differences ($P<0.002$ and $P<0.001$, respectively) at the 0.5 and 1 h time points between the deletion mutant and either of the other two strains tested.

strains were used. Psa fimbriae are also adhesins, but have not been reported to promote internalization (Liu *et al.*, 2006).

Y. pestis OmpX, in contrast to *Y. enterocolitica* Ail, mediates adherence and internalization when bacteria are grown at <30 °C. We propose two potential explanations for this difference. The first is that *Y. enterocolitica* Ail expression is temperature-regulated; low levels of Ail are produced at 28 °C and much higher levels are produced at 37 °C (Bliska & Falkow, 1992; Pierson & Falkow, 1993). However, *Y. pestis* OmpX expression is constitutive. The second is that artificial overexpression of *Y. enterocolitica* Ail at 28 °C does not promote internalization of *Y. enterocolitica*, indicating that additional factors contribute to Ail-mediated internalization (Bliska & Falkow, 1992). As shown previously, adherence conferred by *Y. enterocolitica* Ail is affected by the length or structure of O side-chains in lipopolysaccharide (LPS) which are temperature-regulated (Pierson, 1994). Presumably because of steric interference, only bacteria with shortened O side-chains (*Y. enterocolitica* grown at 37 °C) can promote interaction between Ail and the host cell (Bliska & Falkow, 1992; Pierson, 1994). In contrast to *Y. enterocolitica*, *Y. pestis* LPS, due to mutation, lacks O-antigen regardless of growth temperature (Prior *et al.*, 2001). This lack of O-antigen interference could explain OmpX-mediated internalization at low temperature. The length of O-antigen has been previously shown to regulate functions of other proteins such as Pla and PgtE (Kukkonen *et al.*, 2004). LPS structural diversity (Skurnik *et al.*, 2000; Zhou *et al.*, 2004) might also explain differences between OmpX in *Y. pestis* and Ail in *Y. pseudotuberculosis*. Despite sharing nearly 100 % sequence

homology with *Y. pestis* OmpX, Ail fails to confer adherence of *Y. pseudotuberculosis* to HEp-2 cells (Yang *et al.*, 1996). Another possible explanation of this discrepancy is the one substitution (Phe vs Val) in the predicted third (out of four) surface-exposed-loop regions of *Y. pseudotuberculosis* Ail. Resolution of this issue by complementing the ompX::npt mutant with ail is a goal of future studies.

Like *Y. enterocolitica* Ail and *S. typhimurium* Rck, OmpX mediates serum resistance (Heffernan *et al.*, 1994; Miller *et al.*, 1989, 1990). Disruption of ompX in this study increased sensitivity of *Y. pestis* cultured at either 28 or 37 °C. Resistance was attributed to a direct protective effect by OmpX from complement-mediated killing, rather than shielding of the bacteria from serum components within cell aggregates. Incubating the parental strain in the presence of serum caused complete dissociation of the cell aggregates by 4 h; yet the cells remained resistant to fresh serum added to the suspension at this point (results not shown). The protection by OmpX at either 28 or 37 °C contrasts with *Y. enterocolitica* Ail, which confers this trait only at 37 °C (Bliska & Falkow, 1992; Pierson & Falkow, 1993; Pierson, 1994), reflecting temperature regulation of Ail expression in that organism (Bliska & Falkow, 1992). OmpX expression at both temperatures probably helps ensure *Y. pestis* complement resistance by organisms multiplying in humans or following growth at ambient temperatures. It is clear that OmpX plays a significant role in serum resistance. Since the strains used in this study are wild-type with respect to Pla protease, our results are consistent with a previous report which demonstrated that deletion of Pla protease, known to degrade C3, does not

apparently increase sensitivity to serum (Sodeinde *et al.*, 1992). Comparative studies of another member of the Ail family, Rck of *S. typhimurium*, may suggest that the mode of OmpX action is through inhibition of C9 polymerization and formation of the mature membrane attack complex (MAC) on the bacterial surface (Heffernan *et al.*, 1992b). It has also been reported that *Y. pestis* binds the complement regulatory protein C4bp as observed by Ngampasutadol *et al.* (2005). Whether *Y. pestis* OmpX is directly involved in these latter two processes is currently being examined.

The autoaggregation phenotype associated with pellicle formation and flocculent growth is characteristic of certain *Y. pestis* strains and connected with virulence (Laird & Cavanaugh, 1980). The parental KIM6⁺ strain has this property, which was lost by *Y. pestis* KIM6⁺ Δ ompX::npt grown at either 28 or 37 °C. These phenotypes have been associated with biofilms in some bacteria such as *Campylobacter jejuni* (Joshua *et al.*, 2006), *Mycobacterium smegmatis* (Chen *et al.*, 2006), and *Pseudomonas aeruginosa* (Friedman & Kolter, 2004) and are attributed to resistance to host defence mechanisms (Schembri *et al.*, 2003). Other *Y. pestis* genetic loci, including *hmsHFRS* (in the *pgm* locus) and *hmsT* (Hare & McDonough, 1999), have been correlated with autoaggregation. The presence of *hmsHFRS* and *hmsT* in the *Y. pestis* strains used in this study was confirmed by growth on Congo red agar and by PCR, respectively (data not shown), excluding the possibility that the altered autoaggregation phenotype was attributed to loss of either *hmsHFRS* or *hmsT*. We also demonstrated that *Y. pestis* KIM6⁻, lacking *pgm*, autoaggregated. This finding supports studies by Podladchikova & Rykova (2006) on derivatives of the *Y. pestis* EV76 *hms*⁻ pYV strain in which an unidentified bacterial cell surface protein of ~17 kDa was involved in autoaggregation.

In summary, we have demonstrated a prominent role for *Y. pestis* OmpX in several potential virulence processes. Of three genes (*inv*, *yadA* and *ail*) associated with these processes in enteropathogenic *Yersinia*, *yadA* (Rosqvist *et al.*, 1988) and *inv* (Simonet *et al.*, 1996) are inactive in *Y. pestis*. Experiments are ongoing to investigate other roles of OmpX in pathogenesis and its potential application as a vaccine candidate.

ACKNOWLEDGEMENTS

This work was supported by the National Institutes of Health, grants P20 RR15587, P20 RR016454, and U54AI57141, and the Idaho Agricultural Experimental Station. We appreciate advice and technical assistance provided by Ann Norton and Katarzyna Dziewanowska.

REFERENCES

Bendtsen, J. D., Nielsen, H., von Heijne, G. & Brunak, S. (2004). Improved prediction of signal peptides: SignalP 3.0. *J Mol Biol* **340**, 783–795.

- Bliska, J. B. & Falkow, S. (1992). Bacterial resistance to complement killing mediated by the Ail protein of *Yersinia enterocolitica*. *Proc Natl Acad Sci U S A* **89**, 3561–3565.
- Bobrov, A. G., Geoffroy, V. A. & Perry, R. D. (2002). Yersiniabactin production requires the thioesterase domain of HMWP2 and YbtD, a putative phosphopantetheinylate transferase. *Infect Immun* **70**, 4204–4214.
- Bockmann, R. A. & Cafilisch, A. (2005). Spontaneous formation of detergent micelles around the outer membrane protein OmpX. *Biophys J* **88**, 3191–3204.
- Caspi, R., Foerster, H., Fulcher, C. A., Hopkinson, R., Ingraham, J., Kaipa, P., Krummenacker, M., Paley, S., Pick, J. & other authors (2006). MetaCyc: a multiorganism database of metabolic pathways and enzymes. *Nucleic Acids Res* **34**, D511–D516.
- Chen, J. M., German, G. J., Alexander, D. C., Ren, H., Tan, T. & Liu, J. (2006). Roles of Lsr2 in colony morphology and biofilm formation of *Mycobacterium smegmatis*. *J Bacteriol* **188**, 633–641.
- Cirillo, D. M., Heffernan, E. J., Wu, L., Harwood, J., Fierer, J. & Guiney, D. G. (1996). Identification of a domain in Rck, a product of the *Salmonella typhimurium* virulence plasmid, required for both serum resistance and cell invasion. *Infect Immun* **64**, 2019–2023.
- Cowan, C., Jones, H. A., Kaya, Y. H., Perry, R. D. & Straley, S. C. (2000). Invasion of epithelial cells by *Yersinia pestis*: evidence for a *Y. pestis*-specific invasin. *Infect Immun* **68**, 4523–4530.
- Datsenko, K. A. & Wanner, B. L. (2000). One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc Natl Acad Sci U S A* **97**, 6640–6645.
- de Lorenzo, V., Herrero, M., Jakubzik, U. & Timmis, K. N. (1990). Mini-Tn5 transposon derivatives for insertion mutagenesis, promoter probing, and chromosomal insertion of cloned DNA in gram-negative eubacteria. *J Bacteriol* **172**, 6568–6572.
- Dupont, M., De, E., Chollet, R., Chevalier, J. & Pages, J. M. (2004). Enterobacter aerogenes OmpX, a cation-selective channel *mar*- and osmo-regulated. *FEBS Lett* **569**, 27–30.
- Dziewanowska, K., Carson, A. R., Patti, J. M., Deobald, C. F., Bayles, K. W. & Bohach, G. A. (2000). Staphylococcal fibronectin binding protein interacts with heat shock protein 60 and integrins: role in internalization by epithelial cells. *Infect Immun* **68**, 6321–6328.
- Friedman, L. & Kolter, R. (2004). Genes involved in matrix formation in *Pseudomonas aeruginosa* PA14 biofilms. *Mol Microbiol* **51**, 675–690.
- Gasteiger, E. H. C., Gattiker, A., Duvaud, S., Wilkins, M. R., Appel, R. D. & Bairoch, A. (2005). Protein Identification and Analysis Tools on the ExPASy Server. In *The Proteomics Protocols Handbook*, pp. 571–607. Edited by J. M. Walker. Totowa, NJ: Humana Press.
- Gay, P., Le Coq, D., Steinmetz, M., Ferrari, E. & Hoch, J. A. (1983). Cloning structural gene *sacB*, which codes for exoenzyme levansucrase of *Bacillus subtilis*: expression of the gene in *Escherichia coli*. *J Bacteriol* **153**, 1424–1431.
- Hare, J. M. & McDonough, K. A. (1999). High-frequency RecA-dependent and -independent mechanisms of Congo red binding mutations in *Yersinia pestis*. *J Bacteriol* **181**, 4896–4904.
- Heffernan, E. J., Harwood, J., Fierer, J. & Guiney, D. (1992a). The *Salmonella typhimurium* virulence plasmid complement resistance gene *rck* is homologous to a family of virulence-related outer membrane protein genes, including *pagC* and *ail*. *J Bacteriol* **174**, 84–91.
- Heffernan, E. J., Reed, S., Hackett, J., Fierer, J., Roudier, C. & Guiney, D. (1992b). Mechanism of resistance to complement-mediated killing of bacteria encoded by the *Salmonella typhimurium* virulence plasmid gene *rck*. *J Clin Invest* **90**, 953–964.

- Heffernan, E. J., Wu, L., Louie, J., Okamoto, S., Fierer, J. & Guiney, D. G. (1994). Specificity of the complement resistance and cell association phenotypes encoded by the outer membrane protein genes *rck* from *Salmonella typhimurium* and *ail* from *Yersinia enterocolitica*. *Infect Immun* **62**, 5183–5186.
- Joshua, G. W., Guthrie-Irons, C., Karlyshev, A. V. & Wren, B. W. (2006). Biofilm formation in *Campylobacter jejuni*. *Microbiology* **152**, 387–396.
- Karp, P. D., Ouzounis, C. A., Moore-Kochlacs, C., Goldovsky, L., Kaipa, P., Ahrén, D., Tsoka, S., Darzentas, N., Kunin, V. & López-Bigas, N. (2005). Expansion of the BioCyc collection of pathway/genome databases to 160 genomes. *Nucleic Acids Res* **33**, 6083–6089.
- Katayama, H., Nagasu, T. & Oda, Y. (2001). Improvement of in-gel digestion protocol for peptide mass fingerprinting by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry. *Rapid Commun Mass Spectrom* **15**, 1416–1421.
- Kukkonen, M., Suomalainen, M., Kyllönen, P., Lahteenmaki, K., Lang, H., Virkola, R., Helander, I. M., Holst, O. & Korhonen, T. K. (2004). Lack of O-antigen is essential for plasminogen activation by *Yersinia pestis* and *Salmonella enterica*. *Mol Microbiol* **51**, 215–225.
- Kukleva, L. M., Zadnova, S. P., Shcherbakov, A. A. & Protchenko, O. A. (2000). The role of the protein with molecular weight of 22 kD in the phagocytosis of vaccine strain of *Yersinia pestis* EV. *Zh Mikrobiol Epidemiol Immunobiol* 55–58.
- Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680–685.
- Lahteenmaki, K., Virkola, R., Saren, A., Emody, L. & Korhonen, T. K. (1998). Expression of plasminogen activator Pla of *Yersinia pestis* enhances bacterial attachment to the mammalian extracellular matrix. *Infect Immun* **66**, 5755–5762.
- Lahteenmaki, K., Kukkonen, M. & Korhonen, T. K. (2001a). The Pla surface protease/adhesin of *Yersinia pestis* mediates bacterial invasion into human endothelial cells. *FEBS Lett* **504**, 69–72.
- Lahteenmaki, K., Kuusela, P. & Korhonen, T. K. (2001b). Bacterial plasminogen activators and receptors. *FEMS Microbiol Rev* **25**, 531–552.
- Lahteenmaki, K., Kukkonen, M., Jaatinen, S., Suomalainen, M., Soranummi, H., Virkola, R., Lang, H. & Korhonen, T. K. (2003). *Yersinia pestis* Pla has multiple virulence-associated functions. *Adv Exp Med Biol* **529**, 141–145.
- Laird, W. J. & Cavanaugh, D. C. (1980). Correlation of autoagglutination and virulence of *Yersinia*. *J Clin Microbiol* **11**, 430–432.
- Lee, K., Wang, T., Paszczynski, A. J. & Daoud, S. S. (2006). Expression proteomics to p53 mutation reactivation with PRIMA-1 in breast cancer cells. *Biochem Biophys Res Commun* **349**, 1117–1124.
- Leigh, S. A., Forman, S., Perry, R. D. & Straley, S. C. (2005). Unexpected results from the application of signature-tagged mutagenesis to identify *Yersinia pestis* genes required for adherence and invasion. *Microb Pathog* **38**, 259–266.
- Lindler, L. E., Klempner, M. S. & Straley, S. C. (1990). *Yersinia pestis* pH 6 antigen: genetic, biochemical, and virulence characterization of a protein involved in the pathogenesis of bubonic plague. *Infect Immun* **58**, 2569–2577.
- Liu, F., Chen, H., Galvan, E. M., Lasaro, M. A. & Schifferli, D. M. (2006). Effects of Psa and F1 on the adhesive and invasive interactions of *Yersinia pestis* with human respiratory tract epithelial cells. *Infect Immun* **74**, 5636–5644.
- McEvedy, C. (1988). The bubonic plague. *Sci Am* **258**, 118–123.
- Mecsas, J., Welch, R., Erickson, J. W. & Gross, C. A. (1995). Identification and characterization of an outer membrane protein, OmpX, in *Escherichia coli* that is homologous to a family of outer membrane proteins including Ail of *Yersinia enterocolitica*. *J Bacteriol* **177**, 799–804.
- Miller, V. L. & Falkow, S. (1988). Evidence for two genetic loci in *Yersinia enterocolitica* that can promote invasion of epithelial cells. *Infect Immun* **56**, 1242–1248.
- Miller, S. I., Kukral, A. M. & Mekalanos, J. J. (1989). A two-component regulatory system (*phoP phoQ*) controls *Salmonella typhimurium* virulence. *Proc Natl Acad Sci U S A* **86**, 5054–5058.
- Miller, V. L., Bliska, J. B. & Falkow, S. (1990). Nucleotide sequence of the *Yersinia enterocolitica* *ail* gene and characterization of the Ail protein product. *J Bacteriol* **172**, 1062–1069.
- Miller, V. L., Beer, K. B., Heussipp, G., Young, B. M. & Wachtel, M. R. (2001). Identification of regions of Ail required for the invasion and serum resistance phenotypes. *Mol Microbiol* **41**, 1053–1062.
- Ngampasutadol, J., Ram, S., Blom, A. M., Jarva, H., Jerse, A. E., Lien, E., Goguen, J., Gulati, S. & Rice, P. A. (2005). Human C4b-binding protein selectively interacts with *Neisseria gonorrhoeae* and results in species-specific infection. *Proc Natl Acad Sci U S A* **102**, 17142–17147.
- Nielsen, H. & Krogh, A. (1998). Prediction of signal peptides and signal anchors by a hidden Markov model. *Proc Int Conf Intell Syst Mol Biol* **6**, 122–130.
- Pierson, D. E. (1994). Mutations affecting lipopolysaccharide enhance Ail-mediated entry of *Yersinia enterocolitica* into mammalian cells. *J Bacteriol* **176**, 4043–4051.
- Pierson, D. E. & Falkow, S. (1993). The *ail* gene of *Yersinia enterocolitica* has a role in the ability of the organism to survive serum killing. *Infect Immun* **61**, 1846–1852.
- Podladchikova, O. N. & Rykova, V. A. (2006). Identification of the autoagglutination factor of Hms⁺ cells of the plague agent. *Mol Gen Microbiol Virusol* **25**–29.
- Prior, J. L., Hitchen, P. G., Williamson, D. E., Reason, A. J., Morris, H. R., Dell, A., Wren, B. W. & Titball, R. W. (2001). Characterization of the lipopolysaccharide of *Yersinia pestis*. *Microb Pathog* **30**, 49–57.
- Rebeil, R., Ernst, R. K., Jarrett, C. O., Adams, K. N., Miller, S. I. & Hinnebusch, B. J. (2006). Characterization of late acyltransferase genes of *Yersinia pestis* and their role in temperature-dependent lipid A variation. *J Bacteriol* **188**, 1381–1388.
- Rosqvist, R., Skurnik, M. & Wolf-Watz, H. (1988). Increased virulence of *Yersinia pseudotuberculosis* by two independent mutations. *Nature* **334**, 522–524.
- Schembri, M. A., Kjaergaard, K. & Klemm, P. (2003). Global gene expression in *Escherichia coli* biofilms. *Mol Microbiol* **48**, 253–267.
- Seo, K. S., Lee, S. U., Park, Y. H., Davis, W. C., Fox, L. K. & Bohach, G. A. (2007). Long-term staphylococcal enterotoxin C1 exposure induces soluble factor-mediated immunosuppression by bovine CD4⁺ and CD8⁺ T cells. *Infect Immun* **75**, 260–269.
- Shaffer, J. & Goldin, M. (1974). *Clinical Diagnosis by Laboratory Methods*, 15th edn. Philadelphia, PA: W.B. Saunders Company.
- Shevchenko, A., Wilm, M., Vorm, O. & Mann, M. (1996). Mass spectrometric sequencing of proteins from silver-stained polyacrylamide gels. *Anal Chem* **68**, 850–858.
- Simonet, M., Riot, B., Fortineau, N. & Berche, P. (1996). Invasin production by *Yersinia pestis* is abolished by insertion of an IS200-like element within the *inv* gene. *Infect Immun* **64**, 375–379.
- Skurnik, M., Peippo, A. & Ervela, E. (2000). Characterization of the O-antigen gene clusters of *Yersinia pseudotuberculosis* and the cryptic O-antigen gene cluster of *Yersinia pestis* shows that the plague bacillus is most closely related to and has evolved from *Y. pseudotuberculosis* serotype O:1b. *Mol Microbiol* **37**, 316–330.
- Smith, M. J. (2000). *Genetic regulation of type III secretion systems in Yersinia enterocolitica*, p. 148. PhD thesis, University of Idaho.

- Sodeinde, O. A., Subrahmanyam, Y. V., Stark, K., Quan, T., Bao, Y. & Goguen, J. D. (1992).** A surface protease and the invasive character of plague. *Science* **258**, 1004–1007.
- Stoorvogel, J., van Bussel, M. J., Tommassen, J. & van de Klundert, J. A. (1991).** Molecular characterization of an *Enterobacter cloacae* outer membrane protein (OmpX). *J Bacteriol* **173**, 156–160.
- Surgalla, M. J. & Beesley, E. D. (1969).** Congo red-agar plating medium for detecting pigmentation in *Pasteurella pestis*. *Appl Microbiol* **18**, 834–837.
- Titball, R. W. & Williamson, E. D. (2004).** *Yersinia pestis* (plague) vaccines. *Expert Opin Biol Ther* **4**, 965–973.
- Vandenbosch, J. L., Rabert, D. K. & Jones, G. W. (1987).** Plasmid-associated resistance of *Salmonella typhimurium* to complement activated by the classical pathway. *Infect Immun* **55**, 2645–2652.
- Wren, B. W. (2003).** The yersiniae – a model genus to study the rapid evolution of bacterial pathogens. *Nat Rev Microbiol* **1**, 55–64.
- Yang, Y., Merriam, J. J., Mueller, J. P. & Isberg, R. R. (1996).** The *psa* locus is responsible for thermoinducible binding of *Yersinia pseudotuberculosis* to cultured cells. *Infect Immun* **64**, 2483–2489.
- Zhou, D., Han, Y., Song, Y., Huang, P. & Yang, R. (2004).** Comparative and evolutionary genomics of *Yersinia pestis*. *Microbes Infect* **6**, 1226–1234.
- Zietz, B. P. & Dunkelberg, H. (2004).** The history of the plague and the research on the causative agent *Yersinia pestis*. *Int J Hyg Environ Health* **207**, 165–178.

Edited by: P. van der Ley