Mini-Review

Correspondence

José A. Lemos

Jose_Lemos@urmc.rochester.edu

A model of efficiency: stress tolerance by *Streptococcus mutans*José A. Lemos¹ and Robert A. Burne²¹Center for Oral Biology and Department of Microbiology and Immunology, University of Rochester Medical Center, Rochester, NY 14642, USA²Department of Oral Biology, University of Florida College of Dentistry, Gainesville, FL 32610, USA

The complete genome sequence of *Streptococcus mutans*, a bacterial pathogen commonly associated with human dental caries, was published in 2002. The streamlined genome (2.03 Mb) revealed an organism that is well adapted to its obligately host-associated existence in multispecies biofilms on tooth surfaces: a dynamic environment that undergoes rapid and substantial fluctuations. However, *S. mutans* lacks many of the sensing systems and alternative sigma factors that bacteria often use to coordinate gene expression in response to stress and changes in their environment. Over the past 7 years, functional genomics and proteomics have enhanced our understanding of how *S. mutans* has integrated the stress regulon and global transcriptional regulators to coordinate responses to environmental fluctuations with modulation of virulence in a way that ensures persistence in the oral cavity and capitalizes on conditions that are favourable for the development of dental caries. Here, we highlight advances in dissection of the stress regulon of *S. mutans* and its intimate interrelationship with pathogenesis.

Introduction

Caries is a classic biofilm disease that develops when changes in the oral environment enhance the growth of cariogenic bacteria, which are highly efficient at converting carbohydrates to the organic acids that demineralize tooth enamel. Microbiological assessment of caries-active sites and studies with experimental animals implicated *Streptococcus mutans* as the primary causative agent of human dental caries (Loesche, 1986). The virulence of *S. mutans* resides in three core attributes: its abilities to form biofilms on the tooth surface, to produce large quantities of organic acids (acidogenicity) from a wide range of carbohydrates, and to tolerate environmental stresses, particularly low pH (aciduricity) (Lemos *et al.*, 2005). In addition to dental caries, *S. mutans* is often an agent in subacute bacterial endocarditis, a life-threatening inflammation of heart valves.

Unlike most infectious diseases, in which classic virulence factors, such as a toxin, play a clear role in the damage elicited by the organism, the pathology of dental caries is associated almost exclusively with bacterial metabolism. Catabolism of the nutrients in saliva and the host's diet creates stressors in the form of acids, reactive oxygen species, and other agents that damage biomolecules. Thus, stress tolerance by the bacteria is intimately intertwined with virulence. The purpose of this review is to highlight

post-genomic research on genetic, biochemical and physiological mechanisms that have evolved in *S. mutans* to modulate its pathogenic potential in response to nutritional, chemical and physical stresses encountered in complex biofilms.

Seven years PG (post genome)

In 2001, the complete genome sequence of a serotype *c* strain of *S. mutans* became available (Ajdic *et al.*, 2002). Through the application of functional genomics, transcriptomics and proteomics, researchers were able to make rapid progress in dissecting the mechanisms of stress tolerance utilized by this pathogen. One theme that emerged from these studies is that *S. mutans* has streamlined its genome by using pathways that cope with environmental insults to regulate a variety of virulence attributes.

A few years after the completion of the UA159 genome sequence, *S. mutans* microarray slides became available, with generous support from the NIDCR, through the J. Craig Venter Institute (formerly The Institute for Genomic Research, TIGR). To date, microarrays have been used to probe the responses of *S. mutans* to amino acid starvation (Nascimento *et al.*, 2008), oxygen (Ahn *et al.*, 2007), sugar transport (Ajdic & Pham, 2007) and manganese depletion (Arirachakaran *et al.*, 2007); to identify genes that are differentially expressed in biofilms of *S. mutans* compared with free-living cells (Shemesh *et al.*, 2007); and to evaluate

Abbreviations: AgDS, agmatine deiminase system; TCS, two-component system.

the consequences of gene-specific mutations (Abranches *et al.*, 2006, 2008; Lemos *et al.*, 2008; Merritt *et al.*, 2005; Nascimento *et al.*, 2008; Sztajer *et al.*, 2008; Wen *et al.*, 2006).

Proteomic studies have been instrumental in identifying proteins and pathways that participate in acid tolerance and acid adaptation (Len *et al.*, 2004a, b; Rathsam *et al.*, 2005a, b; Welin *et al.*, 2003, 2004; Wilkins *et al.*, 2002, 2003). Of particular interest is a report by Nick Jacques and co-workers that used continuous culture to catalogue changes in the expression of proteins involved in energy metabolism when the growth pH was lowered from 7 to 5 (Len *et al.*, 2004b). By coupling proteomic data with measurements of end-products of carbon utilization, the authors were able to propose that *S. mutans* tolerates growth at low pH by expending energy to extrude H^+ , by modulating the production of acid end-products, and by using branched-chain amino acid biosynthesis as a potential mechanism to reduce acid production and moderate intracellular pH (Len *et al.*, 2004b).

Comparison of the proteome of mature biofilm and planktonic cells of *S. mutans* cells grown at neutral pH revealed that multiple proteins associated with carbon uptake and cell division were downregulated in biofilms, whereas proteins required for the development of genetic competence were upregulated (Rathsam *et al.*, 2005a), the latter finding being consistent with the observation that the transformation efficiency of *S. mutans* is optimal during biofilm growth (Li *et al.*, 2001b). This observation is thought to have significance in terms of plaque ecology. Specifically, co-ordinated production of bacteriocins from *S. mutans* and the development of competence have been documented in high-cell-density environments, suggesting that the organism could use competence-induced cell lysis to acquire DNA from neighbouring species (Kreth *et al.*, 2005, 2006, 2007). Notably, a study with *S. mutans* implicated the presence of DNA released from competence-induced cell lysis in the extracellular matrix with proper biofilm maturation (Petersen *et al.*, 2005). It remains to be determined whether *S. mutans* biofilms acquire DNA from the external environment to provide a nutrient source, to increase genetic diversity, or both

(Spoering & Gilmore, 2006). In addition to the potential impact on commensal organisms in oral biofilms, a direct correlation has been noted between production of the competence-stimulating peptide (CSP) and activation of autolytic pathways with biofilm formation and persistence of *S. mutans*. In particular, when administered at doses beyond the levels necessary to induce competence, CSP of *S. mutans* was found to induce cell lysis (Qi *et al.*, 2005), suggesting the presence of an altruistic programmed cell death pathway. In this case, the 'sacrifice' of a subset of cells may enable the establishment and survival of the remainder of the population.

Stress survival pathways

Bacteria in dental plaque experience a wide range of stresses. The intermittent ingestion of food by the host results in dramatic changes in nutrient availability and pH, but significant variability in oxygen tension and osmolality are also observed (Lemos *et al.*, 2005). Although it has been noted that oral biofilms experience a 'feast or famine' lifestyle, organisms residing in the oral cavity are not really exposed to severely oligotrophic environments. For this reason, studies of the stress responses of *S. mutans* have served as an excellent model to reveal critical differences in the ways that obligately host-associated bacteria cope with environmental stresses when compared with bacteria that have both free-living and host-associated lifestyles. These studies have revealed important contrasts in the way this organism copes with stresses and uses stress regulons/enzymes to coordinate virulence and survival compared to more widely studied bacterial paradigms (Fig. 1).

Δ pH homeostasis and metabolic pathways

The membrane-bound F-ATPase (H^+ -translocating ATPase) is considered the primary determinant of acid tolerance of *S. mutans* because it allows the organism to maintain a cytoplasmic pH that is more alkaline than the extracellular environment (Lemos *et al.*, 2005). Work from the laboratory of Marquis correlated acid tolerance in oral streptococci with the pH optima and absolute level of

Fig. 1. Mechanisms of acid stress tolerance in *S. mutans*.

activity of the F_1F_0 -ATPase (Bender *et al.*, 1986). Moreover, it was recently demonstrated that the F-ATPase of *S. mutans*, and other oral bacteria, can function as an ATP synthase in starved cells grown at low pH (Sheng & Marquis, 2006). The authors demonstrated that in starved cells, a sudden drop in pH results in a rapid increase in ATP, followed by a rapid decline, that enhances protection against acid killing. By using specific inhibitors of F-ATPase, the authors were able to demonstrate that this increase in ATP comes from the enzyme acting as an ATP synthase. Thus, the F-ATPase may play a dual role in acid tolerance – extruding protons from the cells and, under certain conditions, generating ATP for growth and persistence (Fig. 2).

A mechanism for acid resistance used by many oral streptococci is the production of ammonia by urease enzymes or the arginine deiminase system (ADS) (Burne & Marquis, 2000). Organisms carrying these enzymes can convert urea or arginine, respectively, to produce CO_2 and ammonia, which can neutralize acids and give the organisms a competitive advantage in acidified biofilms. Although *S. mutans* is not capable of generating significant quantities of alkali as it lacks urease and ADS pathways (Ajdic *et al.*, 2002), an agmatine deiminase system (AgDS), analogous to the ADS, was characterized in *S. mutans* UA159 (Griswold *et al.*, 2006). The AgDS converts agmatine, a decarboxylated derivative of arginine that is found in human dental plaque, to putrescine, ammonia and CO_2 . Whereas the ADS and urease pathways catalyse substantial environmental alkalization and appear to be associated with caries resistance, the AgDS of *S. mutans* is

expressed at relatively low levels and is unlikely to elicit a significant alkalization of the environment. However, the production of ammonia from agmatine is believed to contribute to the competitive fitness of *S. mutans* at low pH by increasing the cytoplasmic pH and generating ATP that can be used for growth or to extrude protons (Fig. 2) (Griswold *et al.*, 2006).

Another contributor to acid tolerance of *S. mutans* is malolactic fermentation, which catalyses the conversion of the dicarboxylic L-malate, a major acid in fruits such as apple, to the monocarboxylic lactic acid and CO_2 . It was demonstrated that although malate did not serve as a catabolite for growth of *S. mutans*, it did serve to protect the organism against acid killing by increasing the pH of the cytoplasm via production of CO_2 (Fig. 2) (Sheng & Marquis, 2007).

Protection, repair and quality control of macromolecules

One consequence of exposure to environmental stresses is the accumulation of abnormal proteins due to increased errors in transcription and translation. Moreover, ageing cells present in mature biofilms are prone to mistranslation and aggregation. In this context, molecular chaperones and proteases, which modulate the stability of proteins and prevent accumulation of misfolded proteins, are central to physiological homeostasis. In support of this concept, proteome analysis of *S. mutans* grown at steady state in continuous culture at pH 7 or 5 identified several molecular chaperones, proteases and DNA repair enzymes as being upregulated during growth at low pH (Len *et al.*, 2004a).

Fig. 2. Metabolic pathways that contribute to acid tolerance by increasing the cytoplasmic pH and/or by generating ATP. AgDS, agmatine deiminase system; BCAA, branched-chain amino acid; MLF, malolactic fermentation.

The GroEL and DnaK chaperones take part in several cellular processes, including protein folding, renaturation, and presentation of proteins for degradation. In *S. mutans*, DnaK and GroEL appear to be indispensable, and the essential nature of these chaperones was confirmed by forced downregulation of *groEL* and *dnaK* expression (Lemos *et al.*, 2007b). Lowering of DnaK levels resulted in impaired capacity to form biofilms in the presence of glucose and rendered the strain more sensitive to low pH, elevated temperature and H₂O₂ (Lemos *et al.*, 2007b). The acid sensitivity of the DnaK knock-down strain was attributed, at least in part, to the DnaK chaperone participating in the biogenesis or stabilization of the F-ATPase complex (Lemos *et al.*, 2007b). Downregulation of GroEL also resulted in high temperature sensitivity and impaired capacity to form biofilms, but did not affect growth at low pH or in the presence of H₂O₂ (Lemos *et al.*, 2007b). Wen *et al.* (2005) showed that the ribosome-associated peptidyl-prolyl isomerase RopA (trigger factor) is important for adherence and formation of biofilms and for tolerance to low pH and H₂O₂. Inactivation of the surface-associated HtrA protease or the cytoplasmic ClpP peptidase generated multiple stress-sensitive phenotypes in *S. mutans*, and was also linked to altered biofilm formation and reduced genetic competence (Ahn *et al.*, 2005; Biswas & Biswas, 2005; Deng *et al.*, 2007b; Lemos & Burne, 2002). Notably, in all these cases, a unifying theme is the intimate connection between stress responses and biofilm formation, suggesting that the stress regulon of *S. mutans* may be responsible for controlling a broader set of biological functions when compared to organisms with more complex genomes.

Many of the stresses encountered by oral bacteria induce DNA damage; in particular, acid and oxidative stresses increase the formation of abasic sites in DNA. Earlier reports demonstrated an overlap between DNA repair systems and stress-response pathways, including RecA, the endonuclease Smn and the UV repair excinuclease UvrA (Hahn *et al.*, 1999; Hanna *et al.*, 2001; Quivey *et al.*, 1995). More recently, Faustoferri *et al.* (2005) characterized the Smx exonuclease in *S. mutans* and showed that an *smx* mutant strain was highly sensitive to DNA damage caused by the production of hydroxyl radicals via Fenton reaction.

Cell envelope alterations

The importance of cell membrane integrity and composition in relation to changes that affect proton permeability and F-ATPase activity in *S. mutans* has been documented (Lemos *et al.*, 2005). Fozo & Quivey (2004b) showed that, in response to the acidification of its environment, *S. mutans* increases the proportion of monounsaturated membrane fatty acids, which is predicted to decrease proton permeability. Inactivation of a gene responsible for biosynthesis of monounsaturated fatty acids, *fabM*, resulted in a strain that was extremely sensitive to low

pH and unable to maintain ΔpH (Fozo & Quivey, 2004a). Rats infected with the *fabM* mutant exhibited substantially reduced caries, as compared to the parent strain (Fozo *et al.*, 2007).

The significance of membrane protein biogenesis to stress tolerance was demonstrated in a study with mutated strains lacking the signal recognition particle translocation (SRP) pathway or the membrane-localized chaperone YidC, both involved in the translocation and assembly of membrane proteins. Once considered essential for the viability of all organisms, the SRP pathway was found to be dispensable in *S. mutans* (Hasona *et al.*, 2005), although mutants lacking proteins of the SRP pathway or YidC were impaired in growth under a variety of stress conditions (Hasona *et al.*, 2005). The authors observed that YidC and a functional SRP pathway are necessary for optimal insertion of membrane proteins, including the F-ATPase, providing a partial explanation for the diminished acid tolerance of strains lacking YidC or components of the SRP pathway (Hasona *et al.*, 2005, 2007). Notably, mutations in SRP-related genes were also associated with decreases in biofilm formation, providing another example of the overlap between pathways that govern stress tolerance and biofilm formation (Hasona *et al.*, 2007).

Finally, the surface-associated protein BrpA was found to play a role in biofilm development, autolysis, cell division and stress tolerance (Wen *et al.*, 2006). A comparison of the transcriptomes of a *brpA* mutant and its parent revealed significant alterations in the expression of genes involved in cell wall biogenesis, stress tolerance and adherence (Wen *et al.*, 2006). Although the function of BrpA has not been defined, increased autolysis in the Δ*brpA* strain indicates that this protein may play a role in modulating cell wall integrity through modulation of autolytic activities, which could mechanistically link BrpA to the increases in susceptibility to acid and oxidative stresses observed in the BrpA-deficient strain (Wen *et al.*, 2006).

Nutritional regulators and alteration of catabolic pathways

In order to thrive in dental plaque, where there is considerable fluctuation in the nutrient pools, *S. mutans* must be able to adjust its metabolism and gene expression patterns to maximize the use of available substrates (Lemos *et al.*, 2005). Despite the need to endure periods of nutrient limitation, abrupt exposure to an excess amount of carbohydrate in the diet can result in the rapid accumulation of toxic glycolytic intermediates, acidification of the environment, and osmotic stress. To survive nutrient starvation, to cope with the detrimental effects of glycolytic intermediates, and to maintain proper NAD/NADH⁺ balances, *S. mutans* has developed a sophisticated regulatory network that combines transcriptional regulation with allosteric modulation of enzyme activities to coordinate optimal flow of carbohydrates.

Carbohydrate source and availability are key factors affecting the pathogenic potential of oral biofilms. The sugar phosphotransferase system (PTS) is the major carbohydrate transport system in oral streptococci, especially under carbohydrate-limiting conditions. In addition to participating in sugar uptake, PTS components influence many other cellular processes. Mutations in the ManL PTS permease influenced biofilm development, regulation of acid tolerance and global control of gene expression, in particular carbon catabolite repression (Abranches *et al.*, 2006, 2008). Two global regulators of central metabolism genes, CcpA and CodY, have been shown to influence acid tolerance and the expression of other virulence traits of *S. mutans* (Abranches *et al.*, 2008; Lemos *et al.*, 2008). CcpA, a regulator of carbon metabolism in Gram-positive bacteria, has been shown to globally regulate transcription in response to carbohydrate availability, and a CcpA-deficient strain was substantially more acid resistant than its parent (Abranches *et al.*, 2008). The enhanced acid tolerance of the CcpA mutant has been associated with increases in the expression of the PTS that result in higher rates of ATP generation through glycolysis. Microarrays revealed that CodY, a regulator that helps cells to adapt to poor nutritional conditions, is indeed a global regulator of gene expression in *S. mutans* (Lemos *et al.*, 2008). Phenotypic studies revealed that the *codY* mutant had reduced capacity to form biofilms and was more sensitive to growth at low pH (Lemos *et al.*, 2008).

The nutritional alarmone (p)ppGpp also appears to play an important role in orchestrating an appropriate response to multiple environmental and physiological inputs that *S. mutans* encounters in the oral cavity (Fig. 3). When limited for essential amino acids, bacteria accumulate (p)ppGpp by enzymic phosphorylation of GDP and GTP, resulting in downregulation of genes for macromolecular biosynthesis and upregulation of genes for amino acid biosynthesis and stress tolerance. In Gram-positive bacteria, RelA is a bifunctional enzyme with potent (p)ppGpp-synthetic and -degradative activities. In *S. mutans*, RelA was shown to play major roles in the regulation of phenotypic traits that

are required for establishment, persistence and survival (Lemos *et al.*, 2004; Nascimento *et al.*, 2008), further supporting an overlap between circuits that govern nutrient starvation, general stress tolerance and biofilm formation. Until recently, RelA was considered the sole enzyme responsible for synthesis and degradation of (p)ppGpp in Gram-positive bacteria. However, our group recently identified two novel enzymes, designated RelP and RelQ, with (p)ppGpp-synthase activities in *S. mutans* that could be found in a number of related Gram-positive bacteria (Lemos *et al.*, 2007a). A *relAPQ* triple mutant was auxotrophic for the branched-chain amino acids leucine and valine, but not isoleucine, a phenotype that was directly related to CodY-dependent repression of genes involved in the synthesis of branched-chain amino acids (Lemos *et al.*, 2008). Interestingly, RelP is co-transcribed with, and apparently regulated by, the RelRS two-component system (Lemos *et al.*, 2007a) suggesting that *S. mutans* may use environmental signals to optimize cell growth and survival in a manner that allows the organism to balance growth during dietary intake by the host with the capacity to rapidly mount an adaptive response during fasting periods. Consistent with the role of (p)ppGpp in bacteria, homologues of RelRS in *Streptococcus pyogenes*, designated SptRS, were shown to be critical for this bacterium to survive in saliva (Shelburne *et al.*, 2005).

Two-component signal transduction systems

S. mutans lives almost exclusively in densely populated biofilms that form on the tooth surface. The structure and composition of these biofilms are influenced by the capacity of its constituents to adapt to environmental changes. As is typical of bacteria with specialized niches, there are very few alternative sigma factors in the UA159 genome (Ajdic *et al.*, 2002). Thus, regulatory systems, such as two-component systems (TCSs), which integrate various chemical and physical signals to coordinate appropriate gene expression patterns, play a central role in stress tolerance and are viewed as desirable targets for the development of new antimicrobial therapies.

TCSs are composed of a transmembrane sensor kinase that detects environmental changes and a cytosolic response regulator, which is a DNA-binding protein that modulates expression of target genes when phosphorylated by the kinase. In streptococcal species, the number of TCSs is small compared to organisms with a free-living life-style, ranging from as few as 10 in *Streptococcus thermophilus*, to more than 20 in *S. agalactiae*. Sequence analysis initially revealed the presence of 13 TCSs in *S. mutans* UA159 (Ajdic *et al.*, 2002), but the Biswas laboratory identified an additional pair in this same strain (Biswas *et al.*, 2008).

Over the past few years, studies that evaluated the role of TCSs in *S. mutans* have shown that they regulate virulence gene expression, induction of competence, biofilm development, bacteriocin production and stress tolerance (Biswas *et al.*, 2008; Chen *et al.*, 2008; Deng *et al.*, 2007a;

Fig. 3. Illustration of physiological processes regulated by (p)ppGpp in *S. mutans*.

Levesque *et al.*, 2007; Li *et al.*, 2001a, 2002a, b; Qi *et al.*, 2004; Senadheera *et al.*, 2005; Zeng *et al.*, 2006). In particular, two studies from independent laboratories systematically inactivated the genes encoding sensor kinases of all TCSs and evaluated their role in stress tolerance by *S. mutans* (Biswas *et al.*, 2008; Levesque *et al.*, 2007). In the study by Levesque *et al.* (2007), *smu1814c* (*scnK*) and *smu1965c* (*levS*) mutants displayed significantly slower growth at pH 5.5, whereas the *smu1128c* (*ciaH*) mutant grew better than the parental strain in the presence of NaCl or H₂O₂. Biswas *et al.* (2008) found that inactivation of three sensor kinases, *smu486* (*liaS*), *smu1128c* (*ciaH*) and *smu1516c* (*vicK*), affected stress tolerance of strain UA159. However, the *vicK* mutant showed an increased tolerance to puromycin, which causes premature chain termination during protein synthesis (Biswas *et al.*, 2008). The *liaS* and *ciaH* mutants showed reduced growth when incubated in aerobic conditions or on agar medium supplemented with H₂O₂ (Biswas *et al.*, 2008). The *liaS* and *ciaH* mutants also showed increased sensitivity to puromycin, while the *ciaH* mutant showed significant reduction of growth at pH 5 and increased sensitivity to DNA damage caused by mitomycin C (Biswas *et al.*, 2008). Notably, previous reports have also shown that inactivation of *ciaH* resulted in an acid-sensitive phenotype in strains UA159 and UA140 (Ahn *et al.*, 2006; Qi *et al.*, 2004). The *S. mutans* VicRK system was shown to respond to, and protect against, oxidative stress in one particular study (Deng *et al.*, 2007a). A role in oxidative stress response was also assigned to *ScnRK*, as *scnRK* mutants were more sensitive to H₂O₂ and more susceptible to phagocytic killing in non-activated macrophages (Chen *et al.*, 2008). Studies from the Cvitkovitch laboratory have shown that inactivation of *LiaS* or *ComDE* conferred an acid-sensitive phenotype upon strains NG8 and BM71, respectively (Li *et al.*, 2001a, 2002a), although *comD* (*smu1916c*) or *comE* (*smu1917c*) do not appear to affect acid tolerance in strain UA159 (Ahn *et al.*, 2006). Finally, the *smu927-smu928* TCS, designated *relRS*, is co-transcribed with the *relP* (p)ppGpp-synthetase, and has been implicated in survival and persistence as it may help regulate (p)ppGpp metabolism (Lemos *et al.*, 2007a).

In *S. pyogenes*, the TCS *CovRS* regulates expression of approximately 15 % of the genome, including key virulence genes (Graham *et al.*, 2002). In *S. mutans* UA159, *CovR* is an orphan response regulator that controls expression of genes related to biofilm formation and virulence (Biswas *et al.*, 2007; Biswas & Biswas, 2006). Similar to what has been observed for the *S. pyogenes covRS*, expression of the *S. mutans covR* is autoregulated, optimal during exponential growth and induced by addition of Mg²⁺ in a dose-dependent manner (Chong *et al.*, 2008). The extent of the genes controlled by *CovR* in *S. mutans* is not known, but based on the findings obtained in other streptococci, it is expected that *CovR* participates in the stress responses.

Collectively, these data support the idea that there may be substantial heterogeneity among strains in the role of

specific TCSs, not only in the genes they regulate, but also in the external stimuli to which they respond. Nevertheless, *CiaRH* have been consistently found to play a role in the stress responses of *S. mutans*. Moreover, *CiaRH* have also been implicated in competence development, bacteriocin production and biofilm formation (Ahn *et al.*, 2006; Levesque *et al.*, 2007; Qi *et al.*, 2004). More recently, it was demonstrated that the *ciaRH* operon of *S. mutans* consists of three genes, with the first gene, *ciaX*, encoding a small, double-glycine signalling peptide that allows *CiaRH* to modulate its own expression in response to calcium (He *et al.*, 2008). Inactivation of *ciaX*, or point mutations in its calcium-binding domain, resulted in diminished biofilm formation that was rescued by addition of calcium. Human saliva is saturated in calcium (Agha-Hosseini *et al.*, 2006) and calcium is the principal cation in tooth enamel, so calcium signalling may be an important regulator, through *CiaRH*, of stress responses and virulence in *S. mutans*.

Other regulators

Metal ions, including iron and manganese, have been implicated in the regulation of virulence expression by *S. mutans*. In particular, the *SloR* metalloregulator was shown to modulate *S. mutans* biofilm formation, genetic competence and oxidative stress tolerance in response to manganese availability (Rolerson *et al.*, 2006). Work from the Spatafora laboratory has linked *SloR* repression of the transcriptional regulator *gcrR* with acid stress tolerance (Dunning *et al.*, 2008). More specifically, a *gcrR* mutant was more sensitive to low pH and this phenotype was linked to the inability of the mutant to maintain Δ pH homeostasis.

As mentioned above, the *AgDS* has been proposed to enhance acid resistance through alkalinization of the cytoplasm (Griswold *et al.*, 2006). The *AgDS* of *S. mutans* is subject to complex regulation by substrate, catabolite control, and relevant environmental stresses. A *LuxR*-like transcriptional regulator, named *aguR*, was identified upstream of the *aguBDAC* operon. Inactivation of *aguR* decreased *AgD* activity and eliminated agmatine induction, indicating that *AguR* is a major regulator of *AgDS* (Griswold *et al.*, 2006).

Concluding remarks

Genomic and proteomic studies have enabled researchers to make rapid progress in the identification of genes, proteins and pathways that are associated with stress tolerance in *S. mutans*. Because there is a strong overlap between stress tolerance and biofilm development pathways, some of these gene products are attractive targets for the development of new anti-caries therapies (Matsushita & Janda, 2002). In particular, strategies that short-circuit regulatory pathways used by *S. mutans* to sense and respond to environmental signals may have a potent capacity to disrupt cariogenic biofilms.

Acknowledgements

We would like to thank Jacqueline Abranches and Jessica Kajfasz for critical reading of this manuscript. The authors' research in this area was supported by grants RO1 DE13239, DE10362 and DE12236 from the NIDCR.

References

- Abranches, J., Candella, M. M., Wen, Z. T., Baker, H. V. & Burne, R. A. (2006). Different roles of EIIAB^{Man} and EII^{Glc} in regulation of energy metabolism, biofilm development, and competence in *Streptococcus mutans*. *J Bacteriol* **188**, 3748–3756.
- Abranches, J., Nascimento, M. M., Zeng, L., Browngardt, C. M., Wen, Z. T., Rivera, M. F. & Burne, R. A. (2008). CcpA regulates central metabolism and virulence gene expression in *Streptococcus mutans*. *J Bacteriol* **190**, 2340–2349.
- Agha-Hosseini, F., Dizgah, I. M. & Amirkhani, S. (2006). The composition of unstimulated whole saliva of healthy dental students. *J Contemp Dent Pract* **7**, 104–111.
- Ahn, S. J., Lemos, J. A. & Burne, R. A. (2005). Role of HtrA in growth and competence of *Streptococcus mutans* UA159. *J Bacteriol* **187**, 3028–3038.
- Ahn, S. J., Wen, Z. T. & Burne, R. A. (2006). Multilevel control of competence development and stress tolerance in *Streptococcus mutans* UA159. *Infect Immun* **74**, 1631–1642.
- Ahn, S. J., Wen, Z. T. & Burne, R. A. (2007). Effects of oxygen on virulence traits of *Streptococcus mutans*. *J Bacteriol* **189**, 8519–8527.
- Ajdic, D. & Pham, V. T. (2007). Global transcriptional analysis of *Streptococcus mutans* sugar transporters using microarrays. *J Bacteriol* **189**, 5049–5059.
- Ajdic, D., McShan, W. M., McLaughlin, R. E., Savić, G., Chang, J., Carson, M. B., Primeaux, C., Tian, R., Kenton, S. & other authors (2002). Genome sequence of *Streptococcus mutans* UA159, a cariogenic dental pathogen. *Proc Natl Acad Sci U S A* **99**, 14434–14439.
- Airachakaran, P., Benjavongkulchai, E., Luengpailin, S., Ajdic, D. & Banas, J. A. (2007). Manganese affects *Streptococcus mutans* virulence gene expression. *Caries Res* **41**, 503–511.
- Bender, G. R., Sutton, S. V. & Marquis, R. E. (1986). Acid tolerance, proton permeabilities, and membrane ATPases of oral streptococci. *Infect Immun* **53**, 331–338.
- Biswas, S. & Biswas, I. (2005). Role of HtrA in surface protein expression and biofilm formation by *Streptococcus mutans*. *Infect Immun* **73**, 6923–6934.
- Biswas, S. & Biswas, I. (2006). Regulation of the glucosyltransferase (*gtfBC*) operon by CovR in *Streptococcus mutans*. *J Bacteriol* **188**, 988–998.
- Biswas, I., Drake, L. & Biswas, S. (2007). Regulation of *gbpC* expression in *Streptococcus mutans*. *J Bacteriol* **189**, 6521–6531.
- Biswas, I., Drake, L., Erkin, D. & Biswas, S. (2008). Involvement of sensor kinases in the stress tolerance response of *Streptococcus mutans*. *J Bacteriol* **190**, 68–77.
- Burne, R. A. & Marquis, R. E. (2000). Alkali production by oral bacteria and protection against dental caries. *FEMS Microbiol Lett* **193**, 1–6.
- Chen, P. M., Chen, H. C., Ho, C. T., Jung, C. J., Lien, H. T., Chen, J. Y. & Chia, J. S. (2008). The two-component system ScnRK of *Streptococcus mutans* affects hydrogen peroxide resistance and murine macrophage killing. *Microbes Infect* **10**, 293–301.
- Chong, P., Drake, L. & Biswas, I. (2008). Modulation of *covR* expression in *Streptococcus mutans* UA159. *J Bacteriol* **190**, 4478–4488.
- Deng, D. M., Liu, M. J., ten Cate, J. M. & Crielaard, W. (2007a). The VicRK system of *Streptococcus mutans* responds to oxidative stress. *J Dent Res* **86**, 606–610.
- Deng, D. M., ten Cate, J. M. & Crielaard, W. (2007b). The adaptive response of *Streptococcus mutans* towards oral care products: involvement of the ClpP serine protease. *Eur J Oral Sci* **115**, 363–370.
- Dunning, D. W., McCall, L. W., Powell, W. F., Jr, Arscott, W. T., McConocha, E. M., McClurg, C. J., Goodman, S. D. & Spatafora, G. A. (2008). SloR modulation of the *Streptococcus mutans* acid tolerance response involves the GcrR response regulator as an essential intermediary. *Microbiology* **154**, 1132–1143.
- Faustoferri, R. C., Hahn, K., Weiss, K. & Quivey, R. G., Jr (2005). Smx nuclease is the major, low-pH-inducible apurinic/apyrimidinic endonuclease in *Streptococcus mutans*. *J Bacteriol* **187**, 2705–2714.
- Fozo, E. M. & Quivey, R. G., Jr (2004a). The *fabM* gene product of *Streptococcus mutans* is responsible for the synthesis of monounsaturated fatty acids and is necessary for survival at low pH. *J Bacteriol* **186**, 4152–4158.
- Fozo, E. M. & Quivey, R. G., Jr (2004b). Shifts in the membrane fatty acid profile of *Streptococcus mutans* enhance survival in acidic environments. *Appl Environ Microbiol* **70**, 929–936.
- Fozo, E. M., Scott-Anne, K., Koo, H. & Quivey, R. G., Jr (2007). Role of unsaturated fatty acid biosynthesis in virulence of *Streptococcus mutans*. *Infect Immun* **75**, 1537–1539.
- Graham, M. R., Smoot, L. M., Migliaccio, C. A., Virtaneva, K., Sturdevant, D. E., Porcella, S. F., Federle, M. J., Adams, G. J., Scott, J. R. & Musser, J. M. (2002). Virulence control in group A *Streptococcus* by a two-component gene regulatory system: global expression profiling and in vivo infection modeling. *Proc Natl Acad Sci U S A* **99**, 13855–13860.
- Griswold, A. R., Jameson-Lee, M. & Burne, R. A. (2006). Regulation and physiologic significance of the agmatine deiminase system of *Streptococcus mutans* UA159. *J Bacteriol* **188**, 834–841.
- Hahn, K., Faustoferri, R. C. & Quivey, R. G., Jr (1999). Induction of an AP endonuclease activity in *Streptococcus mutans* during growth at low pH. *Mol Microbiol* **31**, 1489–1498.
- Hanna, M. N., Ferguson, R. J., Li, Y. H. & Cvitkovitch, D. G. (2001). *uvrA* is an acid-inducible gene involved in the adaptive response to low pH in *Streptococcus mutans*. *J Bacteriol* **183**, 5964–5973.
- Hasona, A., Crowley, P. J., Levesque, C. M., Mair, R. W., Cvitkovitch, D. G., Bleiweis, A. S. & Brady, L. J. (2005). Streptococcal viability and diminished stress tolerance in mutants lacking the signal recognition particle pathway or YidC2. *Proc Natl Acad Sci U S A* **102**, 17466–17471.
- Hasona, A., Zuobi-Hasona, K., Crowley, P. J., Abranches, J., Ruelf, M. A., Bleiweis, A. S. & Brady, L. J. (2007). Membrane composition changes and physiological adaptation by *Streptococcus mutans* signal recognition particle pathway mutants. *J Bacteriol* **189**, 1219–1230.
- He, X., Wu, C., Yarbrough, D., Sim, L., Niu, G., Merritt, J., Shi, W. & Qi, F. (2008). The *cia* operon of *Streptococcus mutans* encodes a unique component required for calcium-mediated autoregulation. *Mol Microbiol* **70**, 112–126.
- Kreth, J., Merritt, J., Shi, W. & Qi, F. (2005). Co-ordinated bacteriocin production and competence development: a possible mechanism for taking up DNA from neighbouring species. *Mol Microbiol* **57**, 392–404.
- Kreth, J., Merritt, J., Zhu, L., Shi, W. & Qi, F. (2006). Cell density- and ComE-dependent expression of a group of mutacin and mutacin-like genes in *Streptococcus mutans*. *FEMS Microbiol Lett* **265**, 11–17.

- Kreth, J., Hung, D. C., Merritt, J., Perry, J., Zhu, L., Goodman, S. D., Cvitkovitch, D. G., Shi, W. & Qi, F. (2007). The response regulator ComE in *Streptococcus mutans* functions both as a transcription activator of mutacin production and repressor of CSP biosynthesis. *Microbiology* **153**, 1799–1807.
- Lemos, J. A. & Burne, R. A. (2002). Regulation and physiological significance of ClpC and ClpP in *Streptococcus mutans*. *J Bacteriol* **184**, 6357–6366.
- Lemos, J. A., Brown, T. A., Jr & Burne, R. A. (2004). Effects of RelA on key virulence properties of planktonic and biofilm populations of *Streptococcus mutans*. *Infect Immun* **72**, 1431–1440.
- Lemos, J. A., Abranches, J. & Burne, R. A. (2005). Responses of cariogenic streptococci to environmental stresses. *Curr Issues Mol Biol* **7**, 95–107.
- Lemos, J. A., Lin, V. K., Nascimento, M. M., Abranches, J. & Burne, R. A. (2007a). Three gene products govern (p)ppGpp production by *Streptococcus mutans*. *Mol Microbiol* **65**, 1568–1581.
- Lemos, J. A., Luzardo, Y. & Burne, R. A. (2007b). Physiologic effects of forced down-regulation of *dnaK* and *groEL* expression in *Streptococcus mutans*. *J Bacteriol* **189**, 1582–1588.
- Lemos, J. A., Nascimento, M. M., Lin, V. K., Abranches, J. & Burne, R. A. (2008). Global regulation by (p)ppGpp and CodY in *Streptococcus mutans*. *J Bacteriol* **190**, 5291–5299.
- Len, A. C., Harty, D. W. & Jacques, N. A. (2004a). Stress-responsive proteins are upregulated in *Streptococcus mutans* during acid tolerance. *Microbiology* **150**, 1339–1351.
- Len, A. C., Harty, D. W. & Jacques, N. A. (2004b). Proteome analysis of *Streptococcus mutans* metabolic phenotype during acid tolerance. *Microbiology* **150**, 1353–1366.
- Levesque, C. M., Mair, R. W., Perry, J. A., Lau, P. C., Li, Y. H. & Cvitkovitch, D. G. (2007). Systemic inactivation and phenotypic characterization of two-component systems in expression of *Streptococcus mutans* virulence properties. *Lett Appl Microbiol* **45**, 398–404.
- Li, Y. H., Hanna, M. N., Svensater, G., Ellen, R. P. & Cvitkovitch, D. G. (2001a). Cell density modulates acid adaptation in *Streptococcus mutans*: implications for survival in biofilms. *J Bacteriol* **183**, 6875–6884.
- Li, Y. H., Lau, P. C., Lee, J. H., Ellen, R. P. & Cvitkovitch, D. G. (2001b). Natural genetic transformation of *Streptococcus mutans* growing in biofilms. *J Bacteriol* **183**, 897–908.
- Li, Y. H., Lau, P. C., Tang, N., Svensater, G., Ellen, R. P. & Cvitkovitch, D. G. (2002a). Novel two-component regulatory system involved in biofilm formation and acid resistance in *Streptococcus mutans*. *J Bacteriol* **184**, 6333–6342.
- Li, Y. H., Tang, N., Aspiras, M. B., Lau, P. C., Lee, J. H., Ellen, R. P. & Cvitkovitch, D. G. (2002b). A quorum-sensing signaling system essential for genetic competence in *Streptococcus mutans* is involved in biofilm formation. *J Bacteriol* **184**, 2699–2708.
- Loesche, W. J. (1986). Role of *Streptococcus mutans* in human dental decay. *Microbiol Rev* **50**, 353–380.
- Matsushita, M. & Janda, K. D. (2002). Histidine kinases as targets for new antimicrobial agents. *Bioorg Med Chem* **10**, 855–867.
- Merritt, J., Kreth, J., Shi, W. & Qi, F. (2005). LuxS controls bacteriocin production in *Streptococcus mutans* through a novel regulatory component. *Mol Microbiol* **57**, 960–969.
- Nascimento, M. M., Lemos, J. A., Abranches, J., Lin, V. K. & Burne, R. A. (2008). Role of RelA of *Streptococcus mutans* in global control of gene expression. *J Bacteriol* **190**, 28–36.
- Petersen, F. C., Tao, L. & Scheie, A. A. (2005). DNA binding-uptake system: a link between cell-to-cell communication and biofilm formation. *J Bacteriol* **187**, 4392–4400.
- Qi, F., Merritt, J., Lux, R. & Shi, W. (2004). Inactivation of the *ciaH* gene in *Streptococcus mutans* diminishes mutacin production and competence development, alters sucrose-dependent biofilm formation, and reduces stress tolerance. *Infect Immun* **72**, 4895–4899.
- Qi, F., Kreth, J., Levesque, C. M., Kay, O., Mair, R. W., Shi, W., Cvitkovitch, D. G. & Goodman, S. D. (2005). Peptide pheromone induced cell death of *Streptococcus mutans*. *FEMS Microbiol Lett* **251**, 321–326.
- Quivey, R. G., Jr, Faustoferri, R. C., Clancy, K. A. & Marquis, R. E. (1995). Acid adaptation in *Streptococcus mutans* UA159 alleviates sensitization to environmental stress due to RecA deficiency. *FEMS Microbiol Lett* **126**, 257–261.
- Rathsam, C., Eaton, R. E., Simpson, C. L., Browne, G. V., Berg, T., Harty, D. W. & Jacques, N. A. (2005a). Up-regulation of competence-but not stress-responsive proteins accompanies an altered metabolic phenotype in *Streptococcus mutans* biofilms. *Microbiology* **151**, 1823–1837.
- Rathsam, C., Eaton, R. E., Simpson, C. L., Browne, G. V., Valova, V. A., Harty, D. W. & Jacques, N. A. (2005b). Two-dimensional fluorescence difference gel electrophoretic analysis of *Streptococcus mutans* biofilms. *J Proteome Res* **4**, 2161–2173.
- Rolerson, E., Swick, A., Newlon, L., Palmer, C., Pan, Y., Keeshan, B. & Spatafora, G. (2006). The SloR/Dlg metalloregulator modulates *Streptococcus mutans* virulence gene expression. *J Bacteriol* **188**, 5033–5044.
- Senadheera, M. D., Guggenheim, B., Spatafora, G. A., Huang, Y. C., Choi, J., Hung, D. C., Treglown, J. S., Goodman, S. D., Ellen, R. P. & Cvitkovitch, D. G. (2005). A VicRK signal transduction system in *Streptococcus mutans* affects *gtfBCD*, *gbpB*, and *fff* expression, biofilm formation, and genetic competence development. *J Bacteriol* **187**, 4064–4076.
- Shelburne, S. A., III, Sumby, P., Sitkiewicz, I., Granville, C., DeLeo, F. R. & Musser, J. M. (2005). Central role of a bacterial two-component gene regulatory system of previously unknown function in pathogen persistence in human saliva. *Proc Natl Acad Sci U S A* **102**, 16037–16042.
- Shemesh, M., Tam, A. & Steinberg, D. (2007). Differential gene expression profiling of *Streptococcus mutans* cultured under biofilm and planktonic conditions. *Microbiology* **153**, 1307–1317.
- Sheng, J. & Marquis, R. E. (2006). Enhanced acid resistance of oral streptococci at lethal pH values associated with acid-tolerant catabolism and with ATP synthase activity. *FEMS Microbiol Lett* **262**, 93–98.
- Sheng, J. & Marquis, R. E. (2007). Malolactic fermentation by *Streptococcus mutans*. *FEMS Microbiol Lett* **272**, 196–201.
- Spoering, A. L. & Gilmore, M. S. (2006). Quorum sensing and DNA release in bacterial biofilms. *Curr Opin Microbiol* **9**, 133–137.
- Sztajer, H., Lemme, A., Vilchez, R., Schulz, S., Geffers, R., Yip, C. Y., Levesque, C. M., Cvitkovitch, D. G. & Wagner-Dobler, I. (2008). Autoinducer-2-regulated genes in *Streptococcus mutans* UA159 and global metabolic effect of the *luxS* mutation. *J Bacteriol* **190**, 401–415.
- Welin, J., Wilkins, J. C., Beighton, D., Wrzesinski, K., Fey, S. J., Mose-Larsen, P., Hamilton, I. R. & Svensater, G. (2003). Effect of acid shock on protein expression by biofilm cells of *Streptococcus mutans*. *FEMS Microbiol Lett* **227**, 287–293.
- Welin, J., Wilkins, J. C., Beighton, D. & Svensater, G. (2004). Protein expression by *Streptococcus mutans* during initial stage of biofilm formation. *Appl Environ Microbiol* **70**, 3736–3741.
- Wen, Z. T., Suntharaligham, P., Cvitkovitch, D. G. & Burne, R. A. (2005). Trigger factor in *Streptococcus mutans* is involved in stress tolerance, competence development, and biofilm formation. *Infect Immun* **73**, 219–225.

Wen, Z. T., Baker, H. V. & Burne, R. A. (2006). Influence of BrpA on critical virulence attributes of *Streptococcus mutans*. *J Bacteriol* **188**, 2983–2992.

Wilkins, J. C., Homer, K. A. & Beighton, D. (2002). Analysis of *Streptococcus mutans* proteins modulated by culture under acidic conditions. *Appl Environ Microbiol* **68**, 2382–2390.

Wilkins, J. C., Beighton, D. & Homer, K. A. (2003). Effect of acidic pH on expression of surface-associated proteins of *Streptococcus oralis*. *Appl Environ Microbiol* **69**, 5290–5296.

Zeng, L., Wen, Z. T. & Burne, R. A. (2006). A novel signal transduction system and feedback loop regulate fructan hydrolase gene expression in *Streptococcus mutans*. *Mol Microbiol* **62**, 187–200.