Isolation and analysis of differentially expressed genes in *Penicillium glabrum* subjected to thermal stress

L. Nevarez,¹ V. Vasseur,¹ G. Le Dréan,¹ A. Tanguy,² I. Guisle-Marsollier,³ R. Houlgatte³ and G. Barbier¹

Correspondence

V. Vasseur

valerie.vasseur@univ-brest.fr

¹Laboratoire Universitaire de Biodiversité et Ecologie Microbienne, Université Européenne de Bretagne, Ecole Supérieure de Microbiologie et Sécurité Alimentaire de Brest, Technopôle Brest-Iroise, 28280 Plouzané, France

²Evolution et Génétique des Populations Marines, UMR CNRS 7144, Université Pierre et Marie Curie, Station Biologique de Roscoff, Place Georges Teissier, 29682 Roscoff Cedex, France

³Plate-forme Transcriptomique Ouest-Génopôle, Institut du Thorax INSERM U533, 1 Rue Gaston Veil, BP 53508, 44035 Nantes, Cedex 1, France

Penicillium glabrum is a filamentous fungus frequently involved in food contamination. Numerous environmental factors (temperature, humidity, atmospheric composition, etc.) or food characteristics (water activity, pH, preservatives, etc.) could represent potential sources of stress for micro-organisms. These factors can directly affect the physiology of these spoilage micro-organisms: growth, conidiation, synthesis of secondary metabolites, etc. This study investigated the transcriptional response to temperature in *P. glabrum*, since this factor is one of the most important for fungal growth. Gene expression was first analysed by using suppression subtractive hybridization to generate two libraries containing 445 different up- and downregulated expressed sequence tags (ESTs). Expression of these ESTs was then assessed for different thermal stress conditions, with cDNA microarrays, resulting in the identification of 35 and 49 significantly up- and downregulated ESTs, respectively. These ESTs encode heat-shock proteins, ribosomal proteins, superoxide dismutase, trehalose-6-phosphate synthase and a large variety of identified or unknown proteins. Some of these may be molecular markers for thermal stress response in *P. glabrum*. To our knowledge, this work represents the first study of the transcriptional response of a food spoilage filamentous fungus under thermal stress conditions.

Received 16 June 2008

Revised 29 August 2008

Accepted 3 September 2008

INTRODUCTION

Fungi are ubiquitous micro-organisms often associated with spoilage and biodeterioration of a large variety of foods and feedstuffs. *Penicillium* is one of the most widespread fungal genera isolated from food products. In addition to the economic losses caused by these contaminants, many *Penicillium* species can produce mycotoxins that represent a potential health risk for humans and animals (Pitt & Hocking, 1997; Samson *et al.*, 2004). Due to its ability to disperse a large number of spores in the environment, *Penicillium glabrum* is very frequently encountered in the food manufacturing industry (Pitt & Hocking, 1997). This fungus is responsible for spoilage of

many different products, such as cheese (Northolt *et al.*, 1980), stored cereals (Borjesson *et al.*, 1992), maize (Mislivec & Tuite, 1970), rice (Kurata *et al.*, 1968), jam (Udagawa *et al.*, 1977), commercially marketed chestnuts (Overy *et al.*, 2003), soda (Ancasi *et al.*, 2006) and mineral water (Cabral & Fernandez Pinto, 2002). Numerous intrinsic factors (water activity, pH, preservatives, etc.) in foods provide ideal conditions for micro-organisms to develop. Combined with extrinsic factors (temperature, humidity, atmospheric composition, etc.), modifications of these conditions are potential sources of stress that can directly affect the physiology (growth, sporulation, synthesis of secondary metabolites, etc.) of spoilage micro-organisms. *In vitro* data obtained with different *Penicillium* strains have shown significant variation in the production of secondary metabolites among substrates (Chang *et al.*, 1991; Kokkonen *et al.*, 2005). Environmental conditions may also modify mycotoxin production, as has been reported for *Fusarium* species (Magan *et al.*, 2002). Some

Abbreviations: aRNA, amplified RNA; EST, expressed sequence tag; HSP, heat-shock protein; qRT-PCR, real-time quantitative reverse-transcription PCR; SSH, suppression subtractive hybridization.

Two supplementary tables with details of the up- and downregulated ESTs are available with the online version of this paper.

modifications, such as osmotic stress, have been shown to influence the synthesis of secondary metabolites in *Penicillium notatum* (Fiedurek, 1997).

Temperature and water activity are the most important factors that determine the ability of moulds to grow (Dantigny *et al.*, 2005). Fungi, and more generally every organism in each kingdom, have developed strategies to respond rapidly to environmental fluctuations. Among them, elevated temperatures are one of the foremost and best-known stress inducers at the cellular level. The general response to thermal stress is characterized by acquired thermotolerance, which increases survival capacity during subsequent exposure to high temperatures (Piper, 1993). Response to thermal stress has been described as a tolerance mechanism against adverse growth conditions and associated with synthesis of heat-shock proteins (HSPs) (Lindquist, 1986). This mechanism aims to stabilize cell metabolism and prevents the accumulation of denatured proteins, which can be toxic to the cell. Heat-shock response appears to be widespread, spanning many taxonomic ranks; for example, it is found in *Drosophila* (Ashburner & Bonner, 1979), rabbit (Cosgrove & Brown, 1983), *Escherichia coli* (Yamamori & Yura, 1982), *Saccharomyces cerevisiae* (McAlister & Finkelstein, 1980) and *Aspergillus nidulans* (Newbury & Peberdy, 1996). To our knowledge, no data are available on the heat-shock transcriptional response of food-contaminating moulds, such as *P. glabrum*. From a food industry perspective, understanding the effects of various factors on the growth and physiology of a spoilage mould is crucial to better manage the risk of alteration and toxicity of food products. It would therefore be useful to gain more insight into the mechanisms involved in thermal stress and to identify the genes that are differentially regulated in response to thermal stress.

In the present study, the effects of temperature on the transcriptional response of *P. glabrum* were investigated by using two complementary techniques: suppression subtractive hybridization (SSH) and cDNA microarrays. Transcriptional response to heat shock was compared between thermal stress conditions (40 °C, 120 min) and the optimal growth temperature (25 °C). SSH is a powerful method based on suppression PCR, where subtracted cDNA libraries can be constructed to identify genes differentially expressed in response to an experimental variation in physiological, environmental or other factors (Diatchenko *et al.*, 1996). This method has been previously used in fungi to isolate differentially expressed genes associated with different morphologies or growth conditions. It has been applied in *Aspergillus nidulans* to characterize genes specifically expressed in conidiating cultures and mature conidia (Osharov *et al.*, 2002). SSH has been also used to investigate the differential gene expression between mycelium and yeast forms of the human pathogens *Penicillium marneffei* (Liu *et al.*, 2007) and *Paracoccidioides brasiliensis* (Marques *et al.*, 2004) with respect to temperature (25° or 37 °C).

Expressed sequence tags (ESTs) isolated by SSH were then investigated, in different thermal stress conditions, by assessing their expression with the cDNA microarray technique and confirming these results with real-time quantitative reverse transcription PCR (qRT-PCR) on a few selected ESTs. The advent of cDNA microarrays has provided the possibility of analysing the global changes in gene expression that occur in a cell, under different conditions (Shalon *et al.*, 1996). Transcriptional studies using whole-genome microarrays have been conducted on a wide range of environmental conditions (e.g. thermal, osmotic and oxidative stresses) in the yeasts *Sac. cerevisiae* (Gasch *et al.*, 2000; Causton *et al.*, 2001; Sakaki *et al.*, 2003) and *Schizosaccharomyces pombe* (Chen *et al.*, 2003). As the whole genome of *P. glabrum* is not available, we constructed a custom-made cDNA microarray and we report here the results obtained. This work represents, to our knowledge, the first study investigating the transcriptional response of a food spoilage fungus under thermal stress.

METHODS

Biological material and culture conditions. The fungal strain used in this study was isolated from contaminated aromatized mineral water and precisely characterized as *Penicillium glabrum* (Wehmer) Westling according to the reference method used for classifying *Penicillium* species (Pitt, 2000). It was registered as LMSA 1.01.421 in the fungi collection 'Souchothèque de Bretagne' (Brest, France). For conservation and collection of spores, the fungus was cultured in tubes of potato dextrose agar medium (PDA) (Difco) at 25 °C. Spores were collected from 7-day-old mycelium by adding 2 ml sterile water containing 0.01 % Tween 80 (Sigma-Aldrich) to each tube and flooding, with agitation at 250 r.p.m. Mycelia were cultured from a suspension of 5×10^6 spores of *P. glabrum* inoculated in 250 ml Erlenmeyer flasks containing 50 ml potato dextrose broth medium (PDB) (Difco). For SSH, *P. glabrum* was exposed briefly to two different conditions: stress (40 °C, 120 min) and control (25 °C), hereafter referred to as S and C, respectively. In both cases, mycelia were first grown in 50 ml PDB medium at 25 °C, shaken at 120 r.p.m. for 48 h, then sterile-filtered, transferred into 50 ml sterile PDB and grown for 120 min at 25 °C or 40 °C with shaking at 120 r.p.m. Mycelia were then aseptically harvested by filtration, washed thoroughly with sterile water, quickly frozen in liquid nitrogen and stored at -80 °C. For microarray experiments, five different thermal stress conditions were studied, as defined in a reduced central composite design (CCD) (Box *et al.*, 1978): S1 (33 °C, 42 min), S2 (33 °C, 198 min), S3 (40 °C, 120 min), S4 (47 °C, 42 min) and S5 (47 °C, 198 min) (Fig. 1). These five culture conditions were defined from a previous CCD used for physiological investigations of thermal stress in *P. glabrum* (unpublished data). For each condition, mycelia were grown for 48 h in triplicate (a, b and c), heat-shocked according to the different conditions and harvested following the method described above. Mycelia referred as 'Reference' (R) and 'Control' (C) were also grown in triplicate, at 25 °C, in order to constitute respectively the unique reference pool (pool R) or the Ca, Cb and Cc samples used for statistical comparison with results obtained under stress conditions.

Suppression subtractive hybridization. For each sample, the frozen mycelium was reduced to a fine powder by grinding in a mortar containing liquid nitrogen and total RNA was extracted with

Fig. 1. The central composite design used to investigate transcriptional effects of thermal stress in *Penicillium glabrum*.

Trizol (Invitrogen) according to the manufacturer's instructions. Total RNA was quantified using a Nanodrop spectrophotometer (Labtech) and its integrity was verified with a Bioanalyser (Agilent Technologies) and RNA Nano LabChip (Agilent Technologies). Poly(A)⁺ RNAs were then isolated with the PolyATtract kit (Promega) according to the manufacturer's instructions. The Poly(A)⁺ RNAs were then precipitated overnight at -20°C with 0.1 vol. 3 M sodium acetate (pH 5.2) and 1 vol. 2-propanol. Poly(A)⁺ RNAs were then washed twice in 75% (v/v) ethanol and finally resuspended with 12 μl RNase-free water. The Poly(A)⁺ RNAs extracted were then quantified using the Nanodrop spectrophotometer.

Two reciprocal subtractive cDNA libraries were constructed by SSH on poly(A)⁺ RNAs corresponding to C (25°C) and S (40°C , 120 min) conditions using the PCR-Select cDNA subtraction kit (BD Biosciences–Clontech) according to the manufacturer's instructions. After subtraction, isolated cDNA fragments were amplified by PCR with an Advantage cDNA PCR kit (BD Biosciences–Clontech), purified with a QIAquick PCR purification kit (Qiagen) and cloned into pGEM-T easy vector (Promega) in *E. coli* JM109 (Promega) according to the manufacturer's instructions. Bacterial clones of both forward and reverse libraries were cultured for 24 h at 37°C , in Luria–Bertani (LB) medium (supplemented with 100 mg ampicillin l^{-1}). Plasmids were then extracted from these bacterial clones using an alkaline lysis plasmid miniprep protocol (Birnbom & Doly, 1979). Positive clones containing a single insert were then selected after *Eco*RI digestion and 1% agarose gel electrophoresis.

DNA manipulations and sequence analysis. The cDNAs isolated from positive clones were sequenced with BigDye Terminator chemistry on an AB 3130 xl sequencer (Applied Biosystems). All sequencing was performed with the Sp6 primer (5'-GATTAGGTGACACTATAG-3', T_m 56°C) at a concentration of 5 μM for each reaction. The raw sequences obtained were trimmed using Phred software (Ewing *et al.*, 1998); plasmid sequences were eliminated with SeqClean software (TIGR software tools, <http://www.tigr.org/tdb/tgi/software>). The cleaned sequences were then compared using the BLASTX program (Altschul *et al.*, 1990) with non-redundant GenBank and Swiss-Prot protein sequence databases. Sequences of cDNA contained in both subtracted libraries were annotated and the corresponding ESTs were classified based on the function of their putative encoded proteins. Functional classification used in this work comes from a similar study on *Paracoccidioides brasiliensis* (Felipe *et al.*, 2005) and is originally based on Munich Information Center for Protein Sequences (MIPS) clustering (Ruepp *et al.*, 2004).

cDNA microarray construction. In order to normalize the amount of cDNA used for microarrays, the bacterial clones used for the

subtracted libraries were cultured in LB medium (with 100 mg ampicillin l^{-1}) for 6 h at 37°C , at 140 r.p.m. A 20 μl volume of each culture was diluted in 80 μl sterile water, then incubated at 95°C for 8 min and 4°C for 10 min. A 2 μl volume of each lysed bacterial clone solution was used in duplicate as a cDNA template for PCR amplification in 50 μl final reaction volume with an initial denaturation at 94°C for 3 min, 40 cycles of 94°C for 20 s, 56°C for 30 s, 68°C for 2 min and a final elongation step at 68°C for 5 min. The reaction was performed with a Taq DNA polymerase kit (Amersham-Pharmacia Biotech): 3 U Taq polymerase per reaction, 0.25 mM dNTP, 0.01 μM T7 primer (5'-TAATACGACTCACTATAGGG-3', T_m 56°C) and 0.01 μM Sp6 primer (5'-GATTAGGTGACACTATAG-3', T_m 56°C). PCR product duplicates were then pooled. Amplified cDNAs were purified using Multiscreen plate MSNU 03010 (Millipore) and resuspended in $3\times$ saline sodium citrate (SSC; Invitrogen) containing 1.5 M betaine (Sigma-Aldrich). A semiquantitative estimation was then performed by agarose gel electrophoresis with Smart Ladder SF (Eurogentec) in order to validate, for each clone, the presence of a single amplified cDNA band and its normalized intensity. Purified PCR products were then spotted in triplicate onto GAPS II-coated slides (Corning) with a Eurogrid Spotter (Eurogentec).

Target preparation and hybridization. For target preparation, total RNAs were extracted from biological triplicates (a, b, c) of each mycelium sample grown in control (C and R) and stress conditions (S1, S2, S3, S4 and S5), as previously described. Total RNA from each sample was extracted with Trizol (Invitrogen) and genomic DNA leftovers were eliminated by DNase I treatment with the RNase-free DNase set (Qiagen) and the RNeasy MinElute Cleanup kit (Qiagen) according to the manufacturer's instructions. Purified total RNAs were then analysed with a Nanodrop spectrophotometer (Labtech) and Bioanalyser (Agilent Technologies). Total RNAs extracted from the three distinct samples (Ra, Rb and Rc) were then pooled in order to form a single reference pool (pool R). This pooled sample was used as a unique reference for hybridizations against samples corresponding to different conditions (C, S1, S2, S3, S4 and S5). Five hundred nanograms of each RNA sample was used to prepare cDNAs by reverse transcription (RT). Those cDNAs were then 'in vitro transcribed' and labelled in amplified RNA (aRNA) targets with an Amino-Allyl MessageAmp II kit (Ambion) and Cy3/Cy5 dyes (Amersham-Pharmacia Biotech), according to the manufacturer's instructions.

Before target hybridization, the cDNA microarray slides were first hydrated (10 s), dried (10 s) and fixed for 20 s at 60 000 μJ with a Spectrolinker UV cross-linker (Spectroliner). Microarrays were then agitated for 15 min at 100 r.p.m. in a solution containing 130 ml 1-

methyl-2-pyrrolidone (Sigma-Aldrich), 0.22 M succinic anhydride (Sigma-Aldrich) and 10 ml 1 M borate buffer, pH 8. The arrayed cDNA fragments were then denatured by placing the slides in MilliQ water at 95 °C for 2 min, washed in 95 % (v/v) ethanol (Prolabo) and immediately dried by centrifugation at 700 r.p.m. for 3 min at room temperature. Slides were then pre-hybridized by incubating at 42 °C for 1 h in a solution containing 3 × SSC, 0.3 % SDS, 1 % BSA sterilized by 0.22 µm filtration. Microarrays were then washed in five different MilliQ water baths and then immediately dried by centrifugation at 700 r.p.m. for 3 min at room temperature.

Competitive hybridizations were then performed by first mixing 500 ng Cy3-labelled aRNA targets (S) and 500 ng Cy5-labelled aRNA targets (C), denaturing for 2 min at 100 °C and incubating for 30 min at 37 °C. Mixed targets were then transferred onto the microarray surface, a coverslip was applied on top and microarrays were incubated for 12 h at 42 °C. After hybridization, microarrays were rinsed for 2 min at 100 r.p.m. in four successive baths (respectively 2 × SSC, 0.1 % SDS; 1 × SSC and 0.2 × SSC twice) and immediately dried by centrifugation at 700 r.p.m. for 3 min at 25 °C.

Microarray data analysis. Hybridized slides were scanned at 10 µm resolution with a ScanArray 4000 (Perkin Elmer) and ScanArray Express software (Perkin-Elmer). Pixel fluorescence intensity and differential expression ratios were determined from the picture files (.tif) for both channels (Cy3 and Cy5) by using GenePix Pro 6.0 software (Molecular Devices). Lowess fitness regression (Yang *et al.*, 2002) was applied for global normalization of raw expression ratios. Normalized data for the 913 cDNAs were log₂ transformed and where an EST was represented by more than one cDNA, the median value was recorded. A statistical analysis of the results obtained for each experimental condition was carried out with Microsoft Excel. A global ANOVA on results obtained for C, S1 and S2 conditions, followed by a LSD test, was applied in order to isolate ESTs that were significantly ($P \leq 0.05$) differentially expressed between S1 and C and S2 and C. Among these ESTs, only those with S:C expression ratios corresponding to differences of at least 0.7 log₂ (i.e. a 1.62-fold change) were selected for inclusion in the up- and downregulated EST lists.

qRT-PCR experiments. Reverse transcription (RT) was performed on 50 µg poly(A)⁺ RNA purified from total RNA samples used for preparing microarray labelled targets. The ImProm-II Reverse Transcription System (Promega) was used according to the manufacturer's instructions. Real-time quantitative PCR (qPCR) experiments were performed with a Mini-Opticon thermocycler (Bio-Rad). Experiments were performed on 1 µl of two twice-diluted cDNA samples with 0.5 M of each primer and 7.5 µl 2 × iQ SYBR Green Supermix (Bio-Rad) in a final reaction volume of 15 µl. Cycling conditions were 2 min at 50 °C, 2 min at 95 °C, 40 cycles of (15 s at 95 °C, 1 min at 60 °C) and 15 s at 95 °C.

In the first step of the qRT-PCR experiments, the internal normalization control was determined. A set of eight genes was chosen and consisting of β -actin (frequently used as a housekeeping gene) and seven other genes presenting relatively high thermal stability according to microarray results. This group contained seven heterogeneous genes encoding adenosine kinase, a 14-3-3 protein homologue, FAD-dependent thiol oxidase, copper-fist DNA-binding domain protein, ATP synthase gamma chain, pre-mRNA splicing factor ini1 and S-phase kinase-associated protein 1A. The sequences of these genes in *P. glabrum* were unavailable in public databases. However, for the β -actin gene partial sequences were obtained from a previous study. Moreover, the seven other genes had been previously cloned and partially sequenced for SSH experiments.

Another set of ten ESTs was also selected from the microarray analysis in order to assess their expression by qRT-PCR and validate our

results. These ESTs were selected because their expression appeared to vary significantly with thermal stress. These ESTs encode HSP 70, HSP 98/104, trehalose-6-phosphate synthase, import inner-membrane translocase tim50, NADP-dependent alcohol dehydrogenase, DMP phosphatase, hypothetical protein AN5480.2, hypothetical protein EAL 90164.1, AAA family ATPase and transcriptional activator HAA1.

For each gene, a pair of specific primers was designed using Primer 3 software (<http://frodo.wi.mit.edu/cgi-bin/primer3/primer3.cgi>) (Rozen & Skaletsky, 2000) in order to produce 120–180 bp amplicons (Table 1a, b).

Each primer pair was first tested with qRT-PCR in order to check that its PCR amplification efficiency was >0.90. For each gene, the melting curve profile was first inspected in order to verify amplification specificity. Standard curves were then determined by plotting the threshold cycle (C_t) values of serial 10-fold dilutions (5, 50, 500, 5000) of a calibrator cDNA template as a linear function of the log of the dilution factor. The calibrator cDNA was obtained by reverse transcription on a mix of different stressed or control RNAs. Expression values were then calculated from the threshold cycle (C_t) of the fluorescence measures obtained. Normalized data were calculated with the $2^{-\Delta\Delta C_t}$ method (Livak & Schmittgen, 2001) using the expression of an internal control gene. Microarray and qRT-PCR results obtained for the ten selected genes were then compared using a Bland–Altman plot (Bland & Altman, 1996) as implemented by others (Steenman *et al.*, 2005).

RESULTS

Subtracted cDNA libraries

More than 1300 clones were randomly selected from the forward and reverse cDNA subtracted libraries constructed for *P. glabrum* after comparison between C (25 °C) and S (40 °C, 120 min). Among them, 1056 positive clones were sequenced and finally 913 clean and useful cDNA sequences were obtained. Using BLASTX on the GenBank and Swiss-Prot protein databases, 80 % of these sequences matched with ORFs of other fungi: *Aspergillus nidulans*, *Aspergillus fumigatus*, *Emericella nidulans*, *Neurospora crassa*, *Sac. cerevisiae* and *Sch. pombe* (Fig. 2a). Analysis of these data allowed isolation of 445 non-redundant ESTs with an *E* value lower than 10^{-5} . Only eight of the 445 ESTs were represented by more than eight different cDNA clones. The two reciprocal subtracted libraries contained 197 and 275 potentially up- and downregulated ESTs, respectively; 27 of them were shared by both libraries. The results obtained are available from our website at http://pagesperso.univ-brest.fr/~vasseur/Nevarez_et_al_2008/ and are also provided as supplementary data with the online version of this paper. The 197 potentially upregulated ESTs were composed of 108 identified ESTs, 48 ESTs with hypothetical functions and 41 ESTs that appeared to be unknown in the databases. Of the 275 ESTs contained in the downregulated library, 195 matched with annotated sequences, 47 had hypothetical functions and 33 appeared to be unknown. Of all the isolated ESTs, 64 % showed similarity with annotated ORFs, 20 % encoded hypothetical proteins and 16 % did not match any sequence in the public databases (Fig. 2b).

Table 1. Sequences of primers used in this work

EST	Primer	Primer sequence 5'–3'
(a) Primer sequences for the set of eight potential housekeeping genes		
β -actin	β -actin-F	ACGTTGTCCCATCTACGAA
	β -actin-R	GCTCAGCCAGGATCTTCATC
Adenosine kinase	AK-F	CTGGAGGACTCCGTTGACAT
	AK-R	TCCCAGAGTGGAAAATCCTG
14-3-3 protein homologue	14-3-3-F	CCGAGCGCTATGAAGAGATG
	14-3-3-R	ACTCCTCCTTCTGCTCGATG
FAD-dependent thiol oxidase	FAD-F	AACCTCATTATGCGCCTCAC
	FAD-R	GGGCTTCCTGAAGCTTTTCT
Copper-fist DNA-binding domain protein	Copper fist-F	AGGTAGTCCGACCAATGACG
	Copper fist-R	GGCGATATTTCCGAAAGTTGA
ATP synthase gamma chain	ATP synth-F	AACCTCATTATGCGCCTCAC
	ATP synth-R	GGGCTTCCTGAAGCTTTTCT
Pre-mRNA splicing factor ini1	Pre-mRNA-F	TACCCCTCCACTCCACAAAG
	Pre-mRNA-R	GGTACACGCCTTGAGAAGGA
S-phase kinase-associated protein 1A	S-Phase-F	ATGATGACAATCGTCGCAAG
	S-Phase-R	GGGGATTGCCCCTAATCAT
(b) Primer sequences for the set of 10 genes selected for validating the microarray results by qRT-PCR		
HSP 70	Hsp 70-F	CCCCTCATTTGAGGTTGAGTT
	Hsp 70-R	TGGGAGTCGTTGAAGTAGGC
HSP 98/104	Hsp 98-F	GAATGATTGGTGACCTCCT
	Hsp 98-R	CCATCAGCTGCAAAAGAACA
Trehalose-6-phosphate synthase	TpsA-F	TGGGTGTCTTGCAATTGTGAT
	TpsA-R	CTTTTCAGGGTCGATTCCAA
Import inner membrane translocase tim50	Tim50-F	AGGGTAGCCAGATCCCTGTT
	Tim50-R	AGACCTCGCCATCCATAGTG
NADP-dependent alcohol dehydrogenase	NADP Alcohol DH-F	TGACAGGCTTGATTTGGTGA
	NADP Alcohol DH-F	GGTGACCTGAAGGTTCCGTA
2,3-Diketo-5-methylthio-1-phosphopentane phosphatase	2,3 phosphatase-F	TCAAGCCAGCACTGATAACG
	2,3 phosphatase-R	CGGTCTTCCGTGAAGTCTC
Conserved hypothetical protein AN5480.2	AN5480.2-F	AGAATCTTCCAGCCGTTCAA
	AN5480.2-R	ATCATGACTCGCGGAGAAAG
Hypothetical protein EAL90164.1	EAL90164.1-F	CCTCCCAACCTCAAACCTCAA
	EAL90164.1-R	GTCATCATCGGACTGGGACT
AAA family ATPase	AAA ATPase-F	GCGACACTCTCAGCAGTCAG
	AAA ATPase-R	CAAGGGTCAGGAGGACTTCA
Transcriptional activator HAA1	HAA1-F	TCGCATAAAGCGAAAAGTTC
	HAA1-R	TGTGAATTGCGGTTGTGTT

The analysis of both subtracted libraries showed a wide variety of ESTs encoding proteins of different functional classes (Fig. 2c, d). Several ESTs coding for proteins often associated with cellular stress (HSPs, trehalose-6-phosphate synthase and superoxide dismutase) appeared to be upregulated after thermal stress (Fig. 2c). ESTs encoding other categories of proteins (metabolism, protein fate, transport proteins, etc.) showed relatively similar representation in both libraries. Some ESTs appeared to be downregulated in *P. glabrum* after heat shock (Fig. 2d), such as those encoding proteins involved in transcription (RNA polymerase, initiation factor, splicing factors, etc.) or implicated in protein synthesis (ribosomal proteins, translation initiation factors, elongation factors, etc.). Other ESTs isolated in each library coded for hypothetical or unknown proteins

cDNA microarray construction and RNA target preparation

The differential expression of ESTs isolated from SSH was analysed using custom-made cDNA microarrays. Amplified cDNAs isolated from up- and downregulated subtracted libraries were spotted on cDNA microarrays. Each slide contained, in triplicate, the 913 subtracted cDNAs corresponding to 445 different ESTs isolated from SSH. Many ESTs were represented by one to four different cDNA sequences and some were even represented by up to eight cDNAs. This redundancy was useful for evaluating the reproducibility of the expression results obtained for each EST.

Before target preparation, the quality of total RNAs extracted was first verified. Analysis of total RNA triplicates

Fig. 2. Transcriptional analysis of *Penicillium glabrum* under thermal stress (40 °C, 120 min) with SSH. (a) Global distribution of BLASTX best hits of up- and downregulated subtracted libraries among organisms; (b) Global view of proteins encoded by ESTs isolated in both up- and downregulated subtracted libraries. (c) Functional classification of proteins encoded by ESTs isolated in the upregulated library and (d) in the downregulated subtracted library.

from C conditions revealed a convenient profile with three different and clearly defined peaks, corresponding to migration marker, 18S and 28S rRNAs. Similar results were obtained for total RNA triplicates extracted from S1, S2, S3 and S4 (data not shown). Analysis of total RNA samples from the more drastic S5 condition (47 °C, 42 min) showed a migration marker peak and a large amount of very small RNA fragments indicating substantial total RNA degradation. We thus excluded the S5 stress condition from the central composite design.

Determination of housekeeping genes for validation of gene expression using qRT-PCR

The final validation of the microarray results by qRT-PCR requires previous selection of internal reference genes for qRT-PCR normalization. Expression of the eight potential housekeeping genes selected (β -actin, adenosine kinase, 14-3-3 protein homologue, FAD-dependent thiol oxidase, copper-fist DNA-binding domain protein, ATP synthase gamma chain, pre-mRNA splicing factor ini1 and S-phase kinase-associated protein 1A) was assessed under the different experimental conditions (C, S1, S2, S3 and S4) with qRT-PCR. For each primer pair tested, PCR specificity was confirmed by melting curve analysis (data not shown). Expression results obtained for the eight genes tested were comparable, where C_t values were stable under C, S1 and S2 conditions but substantially increased under the S3 and S4 conditions (Fig. 3). Given these results, normalization of qRT-PCR results could not be applied to the S3 and S4 conditions since stable expression of internal control genes

Fig. 3. Results obtained with qRT-PCR (C_t) for potential housekeeping genes in control (C) and thermal stress conditions (S1, S2, S3 and S4). $\square$, ATP synthase gamma chain; $\blacktriangle$, β -actin; $\blacklozenge$, adenosine kinase; $\circ$, S-phase kinase-associated protein 1A; $\blacksquare$, 14-3-3 protein homologue; $\triangle$, copper-fist DNA-binding domain protein; $\bullet$, FAD-dependent thiol oxidase; $\diamond$, pre-mRNA splicing factor ini1.

is required for each tested condition. As microarray data could not be validated for these thermal stress conditions, analysis was only done on the S1 and S2 conditions. β -Actin was selected as a housekeeping gene for qRT-PCR normalization, owing to its stable expression under C, S1 and S2 conditions and its frequent use in the literature. ATP synthase gamma chain and adenosine kinase were also useful genes and were used to further confirm qRT-PCR normalization (data not shown).

Microarray results for S1 and S2 thermal stress conditions

Based on the 913 cDNA sequences spotted in triplicate representing 445 different ESTs, the statistical analysis of results with a global ANOVA and LSD test indicated 35 and 49 significantly up- and downregulated ESTs. Their S:C expression ratios corresponded to at least a $\pm 0.7 \log_2$ (1.62-fold change). Expression of most of the isolated ESTs represented an approximately ± 2 -fold change, but a few ESTs showed up to $+4.3$ - and even a -4.0 -fold change for up- and downregulated ESTs, respectively.

Analysis of the microarray results revealed 49 highly significant downregulated ESTs under thermal stress conditions S1 and S2 (Table 2). In this set, 26 ESTs encoded ribosomal proteins and accounted for 53 % of the results. This large group also contained ESTs encoding proteins implicated in transcription (bZIP transcription factor CpcA, transcriptional activator HAA1, histone H4, etc.), RNA splicing (U6 small nuclear RNA-associated protein, mago nashi protein homologue) and protein synthesis (translation elongation factor G2). A few proteases [glutamate carboxypeptidase-like protein, candidapepsin 2 precursor (aspartate protease)] and proteins involved in general metabolism (xanthine phosphoribosyltransferase 1 and ornithine decarboxylase) were also observed. A few downregulated ESTs also encoded miscellaneous proteins and hypothetical or unknown proteins.

From the microarray results, 35 highly significantly upregulated ESTs were identified for thermal stress conditions S1 and S2 (Table 3). In this set, six and four ESTs encoded heat-shock proteins (HSP 30, HSP 60, HSP 70, mitochondrial HSP 70, HSP 90 and HSP 98/104) and proteins with antioxidant functions (superoxide dismutase, glutathione S-transferase 3, cytochrome *c* oxidase polypeptide VIa, flavohaemoprotein), respectively. The upregulated EST set also included seven other ESTs coding for metabolism proteins (L-xylulose reductase, mannitol-1-phosphate-5-dehydrogenase, 2,3-diketo-5-methylthio-1-phosphopentane phosphatase, carboxylesterase, NADP-dependent alcohol dehydrogenase and trehalose-6-phosphate synthase). Upregulated ESTs were also completed by few ESTs encoding proteases (aspergillopepsin A and vacuolar protease A) and other miscellaneous proteins. About 37 % of the identified upregulated ESTs coded for hypothetical or unknown proteins.

Table 2. List of ESTs identified using microarrays as being significantly downregulated under thermal stress conditions S1 and S2

Gene homologue	Accession number		Fold change		Significance
	Swiss-Prot	GenBank	(S1-C)	(S2-C)	
Ribosomal proteins					
40S ribosomal protein S2	P25443		−1.99	−1.87	***
40S ribosomal protein S5 a/b	O14277/Q9P3T6		−2.20	−1.77	***
40S ribosomal protein S8 a/b	O14049/Q9P7B2		−2.21	−1.69	***
40S ribosomal protein S9	P52810		−2.44	−1.87	***
40S ribosomal protein S11	P79013		−2.05	−1.65	***
40S ribosomal protein S12	O59936		−2.85	−2.20	***
40S ribosomal protein S16	Q42340		−2.39	−1.85	***
40S ribosomal protein S17	P27770		−2.12	−1.53	***
40S ribosomal protein S23	Q873W8		−2.56	−2.19	***
40S ribosomal protein S25	Q7SC06		−2.03	−1.68	***
60S ribosomal protein L2	Q75AP7		−1.68	−1.40	***
60S ribosomal protein L10a	Q7RZS0		−3.02	−2.42	***
60S ribosomal protein L11	Q758S7		−1.75	−1.76	***
60S ribosomal protein L14 a	P36105		−2.49	−1.83	***
60S ribosomal protein L15	O13418		−2.92	−2.21	***
60S ribosomal protein L18b	Q8TFH1		−2.17	−1.90	***
60S ribosomal protein L23	P04451		−2.50	−1.85	***
60S ribosomal protein L26b	Q9FJX2		−2.38	−1.92	***
60S ribosomal protein L30	Q7S7F1		−4.02	−2.27	***
60S ribosomal protein L34 b	P40525		−3.76	−2.76	***
60S ribosomal protein L35	Q8L805		−2.31	−1.65	***
60S ribosomal protein L38	Q9C2B9		−2.90	−2.15	***
60S ribosomal protein L44	P52809		−2.83	−2.25	***
60S ribosomal protein L5	O59953		−1.83	−1.46	***
60S ribosomal protein L8	O13672		−1.90	−1.55	***
Ribosome biogenesis protein RPF2	P36160		−1.72	−1.46	**
Transcription and translation proteins					
Histone H4	P04914		−1.82	−1.34	***
Translation elongation factor G2		EAL91882.1	−1.85	−1.50	***
Translationally controlled tumour protein homologue	P35691		−2.06	−1.58	***
Transcriptional activator HAA1	Q12753		−2.77	−2.90	**
bZIP transcription factor CpcA		EAL89546.1	−1.93	−1.96	**
U6 snRNA-associated protein LSm3 (splicing)	Q9Y7M4		−1.92	−1.57	***
Mago nashi protein homologue (splicing)	P49030		−1.95	−1.64	*
Metabolism proteins					
Xanthine phosphoribosyltransferase 1	P47165		−1.85	−1.54	**
Ornithine decarboxylase	P27121		−2.58	−2.14	*
Proteases					
Candidapepsin 2 precursor (aspartate protease)	P28871		−2.01	−1.97	***
Glutamate carboxypeptidase-like protein	Q9P6I2		−1.63	−1.37	*
Miscellaneous proteins					
AAA family ATPase		EAL86805.1	−1.73	−2.19	***
Guanine nucleotide-binding protein (protein G)	Q01369		−2.01	−1.53	***
Prefoldin subunit 6 (actin chaperone protein)		EAL92646.1	−2.18	−1.58	***
Hypothetical proteins/unknown proteins					
Hypothetical protein Afu1g07570		EAL88465.1	−1.64	−1.47	***
Hypothetical protein AN0409.2		XP_658013.1	−2.13	−1.81	***
Hypothetical protein AN5226.2		XP_662830.1	−2.21	−1.64	**
Hypothetical protein MG04362.4		XP_361917.1	−2.21	−2.07	*
Unknown protein F17			−1.94	−1.41	***
Unknown protein F26			−1.74	−1.49	***
Unknown protein F6			−2.84	−2.60	***
Unknown protein F9			−2.08	−1.77	***
Unknown protein R16			−3.70	−2.71	***

*, Significant ($P < 0.05$); **, highly significant ($P < 0.01$); ***, very highly significant ($P < 0.001$).

Table 3. List of ESTs isolated using microarrays as being significantly upregulated under thermal stress conditions S1 and S2

Gene homologue	Accession number		Fold change		Significance
	Swiss-Prot	GenBank	(S1-C)	(S2-C)	
Chaperones					
HSP 30	P40920		+ 2.67	+ 2.51	***
HSP 60	O60008		+ 1.74	+ 1.43	***
HSP 70	Q92260		+ 2.98	+ 1.93	***
HSP 90	P40292		+ 3.55	+ 2.06	***
HSP 98/104	P31540		+ 2.28	+ 2.06	***
Mitochondrial Hsp70 chaperone (Ssc70)		EAL93290.1	+ 1.66	+ 1.35	**
Antioxidant proteins					
Cytochrome <i>c</i> oxidase polypeptide VIa	P32799		+ 1.65	+ 1.52	**
Superoxide dismutase	Q9UQX0		+ 1.65	+ 1.62	***
Flavo-haemoprotein (Haemoglobin-like protein)		EAL84490.1	+ 1.69	+ 1.26	***
Glutathione <i>S</i> -transferase 3		AAX07321	+ 1.87	+ 1.60	***
Metabolism proteins					
Trehalose-6-phosphate synthase tpsA	O59921		+ 2.36	+ 1.67	***
2,3-diketo-5-methylthio-1-phosphopentane phosphatase		EAL86676.1	+ 4.34	+ 2.33	***
Carboxylesterase		EAL92659.1	+ 2.14	+ 1.60	***
L-Xylulose reductase	Q8NK50		+ 3.00	+ 1.92	***
Mannitol-1-phosphate 5-dehydrogenase	Q6AHH7		+ 3.22	+ 2.18	***
Plasma membrane ATPase (proton pump)	P24545		+ 1.69	+ 1.42	***
NADP-dependent alcohol dehydrogenase		EAL84214.1	+ 2.64	+ 1.77	***
Proteases					
Aspergillopepsin A	Q06902		+ 1.68	+ 1.63	**
Vacuolar protease A	Q01294		+ 1.78	+ 1.63	***
Miscellaneous proteins					
Metalloreductase, putative		EAL85757.1	+ 1.71	+ 1.60	**
Import inner membrane translocase subunit tim50	Q4WI16		+ 1.66	+ 1.66	**
Protein bli-3 (blue light-inducible protein)		EAL90736.1	+ 2.30	+ 2.05	**
Hypothetical proteins/unknown proteins					
Integral membrane protein		EAL87419.1	+ 2.51	+ 2.02	***
Predicted protein		XP_322336.1	+ 1.71	+ 1.86	***
Conserved hypothetical protein		EAL90164.1	+ 3.58	+ 2.26	***
Hypothetical protein AN0677.2		XP_658281.1	+ 1.76	+ 1.46	**
Hypothetical protein AN5056.2		XP_662660.1	+ 2.42	+ 2.00	**
Hypothetical protein AN5480.2		XP_663084.1	+ 3.23	+ 1.99	***
Hypothetical protein AN5537.2		XP_663141.1	+ 2.12	+ 1.68	***
Hypothetical protein AN9515.2		XP_868897.1	+ 3.01	+ 1.87	***
Unknown protein F18			+ 1.77	+ 1.66	***
Unknown protein F23			+ 1.70	+ 1.78	**
Unknown protein R10			+ 2.57	+ 1.71	***
Unknown protein R29			+ 1.96	+ 1.60	**
Unknown protein R36			+ 2.93	+ 1.91	**

*, Significant ($P < 0.05$); **, highly significant ($P < 0.01$); ***, very highly significant ($P < 0.001$).

Validation of microarray results with qRT-PCR

The microarray results obtained for S1 and S2 conditions were validated by assessing the expression level of a set of 10 differentially expressed ESTs selected from the microarray analysis using qRT-PCR. Relative expression ratios of the 10 selected ESTs were normalized against β -actin gene expression and results obtained were comparable to results from the microarray (Table 4). The Bland–Altman plot

(Fig. 4) carried out with values obtained for S1 and S2 conditions showed that all of the $\log_2$ -transformed microarray data and qRT-PCR data lay within the 95% confidence intervals, delimited by a difference of 1. Moreover, more than 65% of the comparisons differed by less than 0.5. This similarity between both sets of data validates the microarray results for S1 and S2 conditions. Reliability of the data was further verified, as similar results

Table 4. Microarray expression ratios and relative expression results obtained with qRT-PCR under S1 and S2 conditions for the 10 selected ESTs

β -Actin gene expression was used for normalization of qRT-PCR results.

EST	Accession number	S1-C		S2-C	
		Relative expression		Relative expression	
		Microarrays	qRT-PCR	Microarrays	qRT-PCR
HSP 70	Q92260	+ 2.98	+ 4.41	+ 1.93	+ 2.71
HSP 98/104	P31540	+ 2.28	+ 3.34	+ 2.06	+ 3.68
Trehalose-6-phosphate synthase	O59921	+ 2.36	+ 4.22	+ 1.67	+ 2.07
Import inner membrane translocase subunit tim50	Q4WI16	+ 1.66	+ 1.87	+ 1.66	+ 1.98
NADP-dependent alcohol dehydrogenase	EAL84214.1	+ 2.64	+ 4.17	+ 1.77	+ 2.49
2,3-Diketo-5-methylthio-1-phosphopentane phosphatase	EAL86676.1	+ 4.34	+ 3.80	+ 2.33	+ 2.17
Hypothetical protein AN5480.2	XP_663084.1	+ 3.23	+ 2.19	+ 1.99	+ 1.24
Conserved hypothetical protein EAL90164.1	EAL90164.1	+ 3.58	+ 3.84	+ 2.26	+ 1.80
AAA family ATPase	EAL86805.1	- 1.73	- 2.25	- 2.19	- 2.81
Transcriptional activator HAA1	Q12753	- 2.77	- 2.67	- 2.90	- 2.91

were also obtained by normalizing qRT-PCR results with expression of the two other housekeeping genes selected: adenosine kinase and ATP synthase (data not shown).

DISCUSSION

In the first part of the study, SSH was applied to *P. glabrum* to isolate 913 cDNA sequences corresponding to 445 differentially expressed ESTs in response to thermal stress (40 °C, 120 min). SSH is frequently employed to isolate genes differentially expressed in response to varying nutritional or environmental conditions. SSH overcomes the problem of transcript abundance differences in the mRNA population by incorporating a normalization step which equalizes the relative quantity of the cDNA. It enhances the probability of identifying the increased expression of low-abundance transcripts (Diatchenko *et al.*, 1996; Gurskaya *et al.*, 1996).

Many of the isolated ESTs encoded proteins with a wide variety of functions while some of them corresponded to hypothetical or unknown proteins. The large number of ESTs is remarkable compared with results obtained from similar studies on other fungi. For example, SSH was used to isolate specific transcripts in either yeast or mycelial forms of *Ophiostoma piceae* or *Penicillium marneffeii* and only revealed 50 and 43 genes, respectively (Dogra & Breuil, 2004; Liu *et al.*, 2007). Other studies conducted on the effects of different nutritional growth conditions isolated 37 genes implicated in growth with penicillin-repressing or non-repressing carbon sources in *Penicillium chrysogenum* (Castillo *et al.*, 2006). The large number of ESTs isolated here in *P. glabrum* under heat-shock conditions, and their wide functional classification, could be explained by the fact that temperature is involved in general cellular mechanisms and influences many functional categories. It has been shown that two-thirds of the

Fig. 4. Bland–Altman plot showing the agreement between expression ratios obtained under S1 and S2 conditions, with qRT-PCR and microarray analysis for the 10 selected genes.

Sac. cerevisiae genome is involved in response to environmental changes such as temperature, oxidation, nutrients, pH and osmolarity (Causton *et al.*, 2001).

Expression of ESTs contained in both subtracted libraries was further assessed, using microarrays to confirm their differential expression under thermal stress conditions. Custom cDNA microarrays are a very convenient tool for studying subtracted libraries in detail and are more appropriate than Northern blot or qRT-PCR methods for assessing the expression of a large number of ESTs. The combination of SSH and custom cDNA microarrays has been previously employed in *Gibberella zeae* to identify genes involved in sexual reproduction (mediated by ascospores) and implicated in disease occurring in cereal crops (Lee *et al.*, 2006). In our study, the custom microarray used in triplicates of the 913 subtracted cDNA sequences isolated corresponded to 445 different ESTs. Beyond just confirming the subtracted library approach, microarrays were also employed to analyse expression of these ESTs under five different thermal stress conditions (S1, S2, S3, S4 and S5).

Degradation of total RNA observed under S5 conditions (47 °C, 198 min) led us to eliminate this experiment from the study. This result may be correlated with a decrease in mycelium growth observed under these conditions: fungal dry weight decreased about 35 % compared with dry weight under optimal growth conditions (unpublished data). Microarray analysis for the four other stress conditions required final validation by qRT-PCR and prior selection of an internal reference gene for normalization. As no housekeeping genes have yet been defined for *P. glabrum* or closely related fungal species, comparative studies and microarrays were used. Other than the seven potential housekeeping genes selected after microarray analysis, the β -actin gene was also chosen among other well-known internal reference genes such as β -tubulin, glyceraldehyde-3-phosphate dehydrogenase and 18S rRNA (Bustin, 2000; Livak & Schmittgen, 2001). Selection of β -actin was based on its frequent use in qRT-PCR normalization in *Candida albicans* (Nailis *et al.*, 2006), *Phytophthora infestans* (Armstrong *et al.*, 2005) and *Neurospora crassa* (Mohsenzadeh *et al.*, 1998). A systematic increase in C_t values was observed using qRT-PCR for every gene tested under S3 and S4 conditions compared with C, S1 and S2 conditions in spite of the fact that mRNA quantity and quality was validated before RT. Variation in cDNA abundance can be partially explained by mRNA instability after exposure to high temperatures (Bond, 2006). For example, substantial decay of Hsp70 mRNA was reported in *Sac. cerevisiae* after a 3 h exposure to 48 °C (Kapoor *et al.*, 1995). mRNA decay primarily involves deadenylation or decapping (Tucker & Parker, 2000; Wilusz *et al.*, 2001). These processes probably affect mRNA stability and reverse transcription (Fleige & Pfaffl, 2006) without any change in the global quantity of mRNA. Given the increase in C_t values, housekeeping genes were not stable and microarray analyses were not possible for

the S3 and S4 conditions. This result was surprising since a preliminary study in *P. glabrum* showed only a slight decrease in growth under similar conditions (40 °C, 120 min; unpublished data). Analysis of the quality of the total RNAs extracted from the S3 and S4 conditions indicated that RNA integrity was conserved under both conditions. In addition, other thermal stress studies on fungi have been successfully conducted at temperatures close to 40 °C: 37 °C for *Sac. cerevisiae* (Gasch *et al.*, 2000; Causton *et al.*, 2001; Sakaki *et al.*, 2003), 39 °C for *Sch. pombe* (Chen *et al.*, 2003), 42 °C for *N. crassa* (Mohsenzadeh *et al.*, 1998), 45 °C for *C. albicans* (Zeuthen & Howard, 1989) and 50 °C for *Aspergillus niger* (Bai *et al.*, 2003). Nevertheless, results obtained for the S3 and S4 conditions clearly showed that defining a standard 'housekeeping' gene can be difficult for experiments conducted under diverse conditions, as reported by some authors (Schmittgen & Zakrajsek, 2000; Yan & Liou, 2006).

Given these results, both S3 and S4 conditions were excluded from the microarray analysis and results obtained for S1 and S2 were then confirmed using qRT-PCR on 10 genes as observed in the Bland–Altman plot (see Fig. 4). Statistical treatment of the microarray data identified 35 significantly upregulated ESTs and 49 significantly downregulated. Some of the downregulated ESTs encoded proteases or general metabolism proteins. To our knowledge none of these genes have ever been described as being downregulated in previous thermal stress studies performed on fungi. Another group of downregulated ESTs encoded proteins implicated in transcription, RNA splicing and protein synthesis. Several studies carried out on *Sac. cerevisiae* exposed to thermal stress (25–37 °C) have also reported downregulation of genes coding for histone H4 (Gasch *et al.*, 2000; Causton *et al.*, 2001; Sakaki *et al.*, 2003), translation elongation factor G2 (Sakaki *et al.*, 2003) and U6 small nuclear RNA-associated protein (Gasch *et al.*, 2000). The group of ESTs involved in translation included a large set of 27 ESTs encoding ribosomal proteins. Downregulation of these ESTs and repression of the synthesis of the corresponding proteins have been frequently reported in the literature. In *Sac. cerevisiae*, it has been shown that ribosomal protein synthesis is repressed after exposure to 33 °C for 90 min (Gorenstein & Warner, 1976), seemingly linked to a decrease of their corresponding mRNAs (Hereford & Rosbash, 1977) caused by inhibition of their transcription (Kim & Warner, 1983). Microarray studies at 25–37 °C in *Sac. cerevisiae* and at 30–39 °C in *Sch. pombe* (Gasch *et al.*, 2000; Causton *et al.*, 2001; Chen *et al.*, 2003; Sakaki *et al.*, 2003) also confirmed downregulation of these genes in response to thermal stress. For example, in *Sac. cerevisiae* mRNAs have been reported to represent an important part of total mRNAs; repression of these genes may therefore allow energy to be dedicated to increased expression of genes involved in the protective response (Jelinsky & Samson, 1999). This last group of ESTs encoding ribosomal proteins, may contain some candidates for molecular markers of thermal stress in *P. glabrum*.

In addition, we identified 35 upregulated ESTs that encoded proteases and some proteins implicated in metabolism. As for downregulated ESTs, to our knowledge, none of these genes have been previously reported in the literature on fungi, except that encoding trehalose-6-phosphate synthase (*tpsA*), frequently described as being induced under heat-shock conditions in yeasts (Gasch *et al.*, 2000; Causton *et al.*, 2001; Chen *et al.*, 2003; Sakaki *et al.*, 2003). Trehalose has been described as acting as a protectant against various environmental stresses and has been shown to stabilize cellular structures under stress conditions (Tereshina, 2005). This disaccharide has been previously described in *Sac. cerevisiae* to play a role in the acquisition of stress tolerance (De Virgilio *et al.*, 1994; Hottiger *et al.*, 1994; Singer & Lindquist, 1998). For *P. glabrum*, although it has never been reported in the literature, an accumulation of trehalose concurs with our preliminary results (unpublished data).

A group of upregulated ESTs encoding HSPs was also identified in our study. These highly conserved proteins are certainly the most investigated element of the thermal stress response and an increase in expression of the corresponding genes has been frequently observed in thermal stress studies (Lindquist, 1986). These proteins are classified into several families according to their molecular mass and, despite their different roles and properties, most HSPs have been shown to generally act as molecular chaperones. They are mainly involved in folding, assembly, regulation and degradation of other proteins and play an important role in thermotolerance (Craig, 1985; Feder & Hofmann, 1999).

Exposure to elevated temperatures has been previously reported to generate reactive oxygen species (ROS) responsible for oxidative damage in yeasts and filamentous fungi (Davidson *et al.*, 1996; Noventa-Jordao *et al.*, 1999; Bai *et al.*, 2003). As observed in our results, antioxidant systems appear to be extremely important for detoxifying cells. The synthesis of enzymes such as superoxide dismutase, glutathione S-transferase and cytochrome *c* oxidase VIa may serve to reduce the level of ROS in cells (Choi *et al.*, 1998; Marques *et al.*, 2004). In *Sac. cerevisiae*, it was shown that the synthesis of superoxide dismutase is enhanced in thermal stress conditions (Piper, 1995) and plays a role in resistance to heat shock from 30 °C to 50 °C (Davidson *et al.*, 1996). Microarray studies on the yeasts *Sac. cerevisiae* and *Sch. pombe* have also confirmed induction of genes encoding superoxide dismutase, cytosolic glutathione S-transferase and cytochrome *c* oxidase popypeptide VIb (which has similarity to the cytochrome *c* oxidase popypeptide VIa gene isolated in this study) (Gasch *et al.*, 2000; Causton *et al.*, 2001; Chen *et al.*, 2003). These ESTs coding for antioxidant proteins also appeared to be induced in heat-shocked *P. glabrum* and could be molecular markers of thermal stress.

Considering these results, this study provides a preliminary basis for analysis of the genetic regulation of the thermal

stress response in *P. glabrum*. Temperature is one of the most important factors for fungal growth and may represent a potential source of stress that can directly affect the physiology of fungi (growth, conidiation, synthesis of secondary metabolites, etc.). The transcriptional analysis performed here combined efficient and complementary molecular techniques, making it possible to isolate 84 heat-regulated ESTs with a wide variety of functions and to assess their expression under thermal stress conditions. Some of these ESTs have already been associated with different stress-related responses as they encode HSPs, trehalose-6-phosphate synthase, superoxide dismutase, ribosomal proteins or proteins involved in transcription/translation. In *P. glabrum*, they may be 'common environmental stress response' genes and respond in a similar manner to many different environmental changes to those identified in studies on yeasts (Gasch *et al.*, 2000; Chen *et al.*, 2003). On the other hand, a large number of isolated ESTs encoded general metabolism proteins, proteases, miscellaneous proteins and hypothetical or unknown proteins. To our knowledge, a large majority of them have never been associated with fungal stress before and may represent new stress markers to characterize the physiological state of *P. glabrum* (stressed or unstressed). Nevertheless, as reported by some authors, caution should be exercised when analysing microarray results as mRNA stability and decay induced by stress treatments may vary and have some influence on gene expression profiling (Monje-Casas *et al.*, 2004; Michan *et al.*, 2005). Further investigations may study more precisely the stability of each mRNA and investigate the response of the different potential markers isolated.

ACKNOWLEDGEMENTS

This work benefited from financial support from the Brittany Regional Council and the invaluable help of the sequencing and microarray facilities of Ouest-Génopôle located in Roscoff and Nantes, respectively. The authors also thank Dr Jean-Marie Heslan (Inserm, Nantes) and Dr David Mazurais (Ifremer, Brest) for their helpful advice and technical support for qRT-PCR experiments. The authors would also like to thank Emeric Dubois (Inserm, Nantes) for his assistance with the microarray analysis.

REFERENCES

- Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. (1990). Basic local alignment search tool. *J Mol Biol* **215**, 403–410.
- Ancasi, E. G., Carrillo, L. & Benitez Ahrendts, M. R. (2006). Moulds and yeasts in bottled water and soft drinks. *Rev Argent Microbiol* **38**, 93–96.
- Armstrong, M. R., Whisson, S. C., Pritchard, L., Bos, J. I., Venter, E., Avrova, A. O., Rehmany, A. P., Bohme, U., Brooks, K. & other authors (2005). An ancestral oomycete locus contains late blight avirulence gene *Avr3a*, encoding a protein that is recognized in the host cytoplasm. *Proc Natl Acad Sci U S A* **102**, 7766–7771.
- Ashburner, M. & Bonner, J. J. (1979). The induction of gene activity in *Drosophila* by heat shock. *Cell* **17**, 241–254.

- Bai, Z., Harvey, L. M. & McNeil, B. (2003). Oxidative stress in submerged cultures of fungi. *Crit Rev Biotechnol* **23**, 267–302.
- Birnboim, H. C. & Doly, J. (1979). A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res* **7**, 1513–1523.
- Bland, J. M. & Altman, D. G. (1996). Measurement error. *BMJ* **313**, 744.
- Bond, U. (2006). Stressed out! Effects of environmental stress on mRNA metabolism. *FEMS Yeast Res* **6**, 160–170.
- Borjesson, T., Stollman, U. & Schnurer, J. (1992). Volatile metabolites produced by six fungal species compared with other indicators of fungal growth on cereal grains. *Appl Environ Microbiol* **58**, 2599–2605.
- Box, G. E. P., Hunter, W. G. & Hunter, J. S. (1978). *Statistics for Experimenters. An Introduction to Design Data Analysis and Model Building*. New York: Wiley.
- Bustin, S. A. (2000). Absolute quantification of mRNA using real-time reverse transcription polymerase chain reaction assays. *J Mol Endocrinol* **25**, 169–193.
- Cabral, D. & Fernandez Pinto, V. E. (2002). Fungal spoilage of bottled mineral water. *Int J Food Microbiol* **72**, 73–76.
- Castillo, N. I., Fierro, F., Gutierrez, S. & Martin, J. F. (2006). Genome-wide analysis of differentially expressed genes from *Penicillium chrysogenum* grown with a repressing or a non-repressing carbon source. *Curr Genet* **49**, 85–96.
- Causton, H. C., Ren, B., Koh, S. S., Harbison, C. T., Kanin, E., Jennings, E. G., Lee, T. I., True, H. L., Lander, E. S. & other authors (2001). Remodeling of yeast genome expression in response to environmental changes. *Mol Biol Cell* **12**, 323–337.
- Chang, S. C., Wei, Y. H., Wei, D. L., Chen, Y. Y. & Jong, S. C. (1991). Factors affecting the production of eremofortin C and PR toxin in *Penicillium roqueforti*. *Appl Environ Microbiol* **57**, 2581–2585.
- Chen, D., Toone, W. M., Mata, J., Lyne, R., Burns, G., Kivinen, K., Brazma, A., Jones, N. & Bahler, J. (2003). Global transcriptional responses of fission yeast to environmental stress. *Mol Biol Cell* **14**, 214–229.
- Choi, J. H., Lou, W. & Vancura, A. (1998). A novel membrane-bound glutathione S-transferase functions in the stationary phase of the yeast *Saccharomyces cerevisiae*. *J Biol Chem* **273**, 29915–29922.
- Cosgrove, J. W. & Brown, I. R. (1983). Heat shock protein in mammalian brain and other organs after a physiologically relevant increase in body temperature induced by D-lysergic acid diethylamide. *Proc Natl Acad Sci U S A* **80**, 569–573.
- Craig, E. A. (1985). The heat shock response. *CRC Crit Rev Biochem* **18**, 239–280.
- Dantigny, P., Guilmar, A. & Bensoussan, M. (2005). Basis of predictive mycology. *Int J Food Microbiol* **100**, 187–196.
- Davidson, J. F., Whyte, B., Bissinger, P. H. & Schiestl, R. H. (1996). Oxidative stress is involved in heat-induced cell death in *Saccharomyces cerevisiae*. *Proc Natl Acad Sci U S A* **93**, 5116–5121.
- De Virgilio, C., Hottiger, T., Dominguez, J., Boller, T. & Wiemken, A. (1994). The role of trehalose synthesis for the acquisition of thermotolerance in yeast. I. Genetic evidence that trehalose is a thermoprotectant. *Eur J Biochem* **219**, 179–186.
- Diatchenko, L., Lau, Y. F., Campbell, A. P., Chenchik, A., Moqadam, F., Huang, B., Lukyanov, S., Lukyanov, K., Gurskaya, N. & other authors (1996). Suppression subtractive hybridization: a method for generating differentially regulated or tissue-specific cDNA probes and libraries. *Proc Natl Acad Sci U S A* **93**, 6025–6030.
- Dogra, N. & Breuil, C. (2004). Suppressive subtractive hybridization and differential screening identified genes differentially expressed in yeast and mycelial forms of *Ophiostoma piceae*. *FEMS Microbiol Lett* **238**, 175–181.
- Ewing, B., Hillier, L., Wendl, M. C. & Green, P. (1998). Base-calling of automated sequencer traces using Phred. I. Accuracy assessment. *Genome Res* **8**, 175–185.
- Feder, M. E. & Hofmann, G. E. (1999). Heat-shock proteins, molecular chaperones, and the stress response: evolutionary and ecological physiology. *Annu Rev Physiol* **61**, 243–282.
- Felipe, M. S., Andrade, R. V., Arraes, F. B., Nicola, A. M., Maranhao, A. Q., Torres, F. A., Silva-Pereira, I., Pocas-Fonseca, M. J., Campos, E. G. & other authors (2005). Transcriptional profiles of the human pathogenic fungus *Paracoccidioides brasiliensis* in mycelium and yeast cells. *J Biol Chem* **280**, 24706–24714.
- Fiedurek, J. (1997). Enhancement of β -galactosidase production and secretion by high osmotic stress in *Penicillium notatum*. *Microbiol Res* **153**, 65–69.
- Fleige, S. & Pfaffl, M. W. (2006). RNA integrity and the effect on the real-time qRT-PCR performance. *Mol Aspects Med* **27**, 126–139.
- Gasch, A. P., Spellman, P. T., Kao, C. M., Carmel-Harel, O., Eisen, M. B., Storz, G., Botstein, D. & Brown, P. O. (2000). Genomic expression programs in the response of yeast cells to environmental changes. *Mol Biol Cell* **11**, 4241–4257.
- Gorenstein, C. & Warner, J. R. (1976). Coordinate regulation of the synthesis of eukaryotic ribosomal proteins. *Proc Natl Acad Sci U S A* **73**, 1547–1551.
- Gurskaya, N. G., Diatchenko, L., Chenchik, A., Siebert, P. D., Khaspekov, G. L., Lukyanov, K. A., Vagner, L. L., Ermolaeva, O. D., Lukyanov, S. A. & other authors (1996). Equalizing cDNA subtraction based on selective suppression of polymerase chain reaction: cloning of Jurkat cell transcripts induced by phytohemagglutinin and phorbol 12-myristate 13-acetate. *Anal Biochem* **240**, 90–97.
- Hereford, L. M. & Rosbash, M. (1977). Regulation of a set of abundant mRNA sequences. *Cell* **10**, 463–467.
- Hottiger, T., De Virgilio, C., Hall, M. N., Boller, T. & Wiemken, A. (1994). The role of trehalose synthesis for the acquisition of thermotolerance in yeast. II. Physiological concentrations of trehalose increase the thermal stability of proteins *in vitro*. *Eur J Biochem* **219**, 187–193.
- Jelinsky, S. A. & Samson, L. D. (1999). Global response of *Saccharomyces cerevisiae* to an alkylating agent. *Proc Natl Acad Sci U S A* **96**, 1486–1491.
- Kapoor, M., Curle, C. A. & Runham, C. (1995). The *hsp70* gene family of *Neurospora crassa*: cloning, sequence analysis, expression, and genetic mapping of the major stress-inducible member. *J Bacteriol* **177**, 212–221.
- Kim, C. H. & Warner, J. R. (1983). Mild temperature shock alters the transcription of a discrete class of *Saccharomyces cerevisiae* genes. *Mol Cell Biol* **3**, 457–465.
- Kokkonen, M., Jestoi, M. & Rizzo, A. (2005). The effect of substrate on mycotoxin production of selected *Penicillium* strains. *Int J Food Microbiol* **99**, 207–214.
- Kurata, H., Sakabe, F., Udagawa, S., Ichino, M. & Suzuki, M. (1968). A mycological examination for the presence of mycotoxin-producers on the 1954–1967's stored rice grains. *Eisei Shikenjo Hokoku* **86**, 183–188.
- Lee, S. H., Lee, S., Choi, D., Lee, Y. W. & Yun, S. H. (2006). Identification of the down-regulated genes in a *mat1-2*-deleted strain of *Gibberella zeae*, using cDNA subtraction and microarray analysis. *Fungal Genet Biol* **43**, 295–310.
- Lindquist, S. (1986). The heat-shock response. *Annu Rev Biochem* **55**, 1151–1191.

- Liu, H., Xi, L., Zhang, J., Li, X., Liu, X., Lu, C. & Sun, J. (2007). Identifying differentially expressed genes in the dimorphic fungus *Penicillium marneffe* by suppression subtractive hybridization. *FEMS Microbiol Lett* **270**, 97–103.
- Livak, K. J. & Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta C_t}$ method. *Methods* **25**, 402–408.
- Magan, N., Hope, R., Colleate, A. & Baxter, E. S. (2002). Relationship between growth and mycotoxin production by *Fusarium* species, biocides and environment. *Eur J Plant Pathol* **108**, 685–690.
- Marques, E. R., Ferreira, M. E., Drummond, R. D., Felix, J. M., Menossi, M., Savoldi, M., Travassos, L. R., Puccia, R., Batista, W. L. & other authors (2004). Identification of genes preferentially expressed in the pathogenic yeast phase of *Paracoccidioides brasiliensis*, using suppression subtraction hybridization and differential macroarray analysis. *Mol Genet Genomics* **271**, 667–677.
- McAlister, L. & Finkelstein, D. B. (1980). Heat shock proteins and thermal resistance in yeast. *Biochem Biophys Res Commun* **93**, 819–824.
- Michan, C., Monje-Casas, F. & Pueyo, C. (2005). Transcript copy number of genes for DNA repair and translesion synthesis in yeast: contribution of transcription rate and mRNA stability to the steady-state level of each mRNA along with growth in glucose-fermentative medium. *DNA Repair* **4**, 469–478.
- Mislivec, P. B. & Tuite, J. (1970). Species of *Penicillium* occurring in freshly-harvested and in stored dent corn kernels. *Mycologia* **62**, 67–74.
- Mohsenzadeh, S., Saupe-Thies, W., Steier, G., Schroeder, T., Fracella, F., Ruoff, P. & Rensing, L. (1998). Temperature adaptation of house keeping and heat shock gene expression in *Neurospora crassa*. *Fungal Genet Biol* **25**, 31–43.
- Monje-Casas, F., Michan, C. & Pueyo, C. (2004). Absolute transcript levels of thioredoxin- and glutathione-dependent redox systems in *Saccharomyces cerevisiae*: response to stress and modulation with growth. *Biochem J* **383**, 139–147.
- Nailis, H., Coenye, T., Van Nieuwerburgh, F., Deforce, D. & Nelis, H. J. (2006). Development and evaluation of different normalization strategies for gene expression studies in *Candida albicans* biofilms by real-time PCR. *BMC Mol Biol* **7**, 25.
- Newbury, J. & Peberdy, J. F. (1996). Characterization of the heat shock response in protoplasts of *Aspergillus nidulans*. *Mycol Res* **100**, 1325–1332.
- Northolt, M. D., van Egmond, H. P., Soentoro, P. & Deijl, E. (1980). Fungal growth and the presence of sterigmatocystin in hard cheese. *J Assoc Off Anal Chem* **63**, 115–119.
- Noventa-Jordao, M. A., Couto, R. M., Goldman, M. H., Aguirre, J., Iyer, S., Caplan, A., Terenzi, H. F. & Goldman, G. H. (1999). Catalase activity is necessary for heat-shock recovery in *Aspergillus nidulans* germlings. *Microbiology* **145**, 3229–3234.
- Osharov, N., Mathew, J., Romans, A. & May, G. S. (2002). Identification of conidial-enriched transcripts in *Aspergillus nidulans* using suppression subtractive hybridization. *Fungal Genet Biol* **37**, 197–204.
- Overy, D. P., Seifert, K. A., Savard, M. A. & Frisvad, J. C. (2003). Spoilage fungi and their mycotoxins in commercially marketed chestnuts. *Int J Food Microbiol* **88**, 69–77.
- Piper, P. W. (1993). Molecular events associated with acquisition of heat tolerance by the yeast *Saccharomyces cerevisiae*. *FEMS Microbiol Rev* **11**, 339–355.
- Piper, P. W. (1995). The heat shock and ethanol stress response of yeast exhibit extensive similarity and functional overlap. *FEMS Microbiol Lett* **134**, 121–127.
- Pitt, J. I. (2000). *A Laboratory Guide to Common Penicillium Species*, 3rd edn. North Ryde, Australia: Food Science.
- Pitt, J. I. & Hocking, A. D. (1997). *Fungi and Food Spoilage*. London: Blackie Academic and Professional.
- Rozen, S. & Skaletsky, H. (2000). Primer3 on the WWW for general users and for biologist programmers. *Methods Mol Biol* **132**, 365–386.
- Ruepp, A., Zollner, A., Maier, D., Albermann, K., Hani, J., Mokrejs, M., Tetko, I., Guldener, U., Mannhaupt, G. & other authors (2004). The FunCat, a functional annotation scheme for systematic classification of proteins from whole genomes. *Nucleic Acids Res* **32**, 5539–5545.
- Sakaki, K., Tashiro, K., Kuhara, S. & Mihara, K. (2003). Response of genes associated with mitochondrial function to mild heat stress in yeast *Saccharomyces cerevisiae*. *J Biochem* **134**, 373–384.
- Samson, R. A., Hoekstra, E. S., Frisvad, J. C. & Filtenborg, O. (2004). *Introduction to Food and Airborne Fungi*. Utrecht, The Netherlands: Centraalbureau voor Schimmelfcultures (CBS).
- Schmittgen, T. D. & Zakrajsek, B. A. (2000). Effect of experimental treatment on housekeeping gene expression: validation by real-time, quantitative RT-PCR. *J Biochem Biophys Methods* **46**, 69–81.
- Shalon, D., Smith, S. J. & Brown, P. O. (1996). A DNA microarray system for analyzing complex DNA samples using two-color fluorescent probe hybridization. *Genome Res* **6**, 639–645.
- Singer, M. A. & Lindquist, S. (1998). Thermotolerance in *Saccharomyces cerevisiae*: the Yin and Yang of trehalose. *Trends Biotechnol* **16**, 460–468.
- Steenman, M., Lamirault, G., Le Meur, N., Le Cunff, M., Escande, D. & Leger, J. J. (2005). Distinct molecular portraits of human failing hearts identified by dedicated cDNA microarrays. *Eur J Heart Fail* **7**, 157–165.
- Tereshina, V. M. (2005). Thermotolerance in fungi: the role of heat shock proteins and trehalose. *Microbiology* **74**, 293–304.
- Tucker, M. & Parker, R. (2000). Mechanisms and control of mRNA decapping in *Saccharomyces cerevisiae*. *Annu Rev Biochem* **69**, 571–595.
- Udagawa, S. I., Kobatake, M. & Kurata, H. (1977). Re-estimation of preservation effectiveness of potassium sorbate (food additive) in jams and marmalade. *Eisei Shikenjo Hokoku* **88–92** (in Japanese).
- Wilusz, C. J., Gao, M., Jones, C. L., Wilusz, J. & Peltz, S. W. (2001). Poly(A)-binding proteins regulate both mRNA deadenylation and decapping in yeast cytoplasmic extracts. *RNA* **7**, 1416–1424.
- Yamamori, T. & Yura, T. (1982). Genetic control of heat-shock protein synthesis and its bearing on growth and thermal resistance in *Escherichia coli* K-12. *Proc Natl Acad Sci U S A* **79**, 860–864.
- Yan, H. Z. & Liou, R. F. (2006). Selection of internal control genes for real-time quantitative RT-PCR assays in the oomycete plant pathogen *Phytophthora parasitica*. *Fungal Genet Biol* **43**, 430–438.
- Yang, Y. H., Dudoit, S., Luu, P., Lin, D. M., Peng, V., Ngai, J. & Speed, T. P. (2002). Normalization for cDNA microarray data: a robust composite method addressing single and multiple slide systematic variation. *Nucleic Acids Res* **30**, e15.
- Zeuthen, M. L. & Howard, D. H. (1989). Thermotolerance and the heat-shock response in *Candida albicans*. *J Gen Microbiol* **135**, 2509–2518.

Edited by: B. A. Horwitz