Plant host and sugar alcohol induced exopolysaccharide biosynthesis in the *Burkholderia cepacia* complex

S. Josefin Bartholdson,^{1,2} Alan R. Brown,² Ben R. Mewburn,³ David J. Clarke,¹ Stephen C. Fry,³ Dominic J. Campopiano¹ and John R. W. Govan²

Correspondence

John R. W. Govan
john.r.w.govan@ed.ac.uk

¹School of Chemistry, University of Edinburgh, Edinburgh EH9 3JJ, UK

²Centre for Infectious Diseases, University of Edinburgh, Edinburgh EH16 4SB, UK

³Institute of Molecular Plant Sciences, University of Edinburgh, Edinburgh EH9 3JH, UK

The species that presently constitute the *Burkholderia cepacia* complex (Bcc) have multiple roles; they include soil and water saprophytes, bioremediators, and plant, animal and human pathogens. Since the first description of pathogenicity in the Bcc was based on sour skin rot of onion bulbs, this study returned to this plant host to investigate the onion-associated phenotype of the Bcc. Many Bcc isolates, which were previously considered to be non-mucoid, produced copious amounts of exopolysaccharide (EPS) when onion tissue was provided as the sole nutrient. EPS production was not species-specific, was observed in isolates from both clinical and environmental sources, and did not correlate with the ability to cause maceration of onion tissue. Chemical analysis suggested that the onion components responsible for EPS induction were primarily the carbohydrates sucrose, fructose and fructans. Additional sugars were investigated, and all alcohol sugars tested were able to induce EPS production, in particular mannitol and glucitol. To investigate the molecular basis for EPS biosynthesis, we focused on the highly conserved *bce* gene cluster thought to be involved in cepacian biosynthesis. We demonstrated induction of the *bce* gene cluster by mannitol, and found a clear correlation between the inability of representatives of the *Burkholderia cenocepacia* ET12 lineage to produce EPS and the presence of an 11 bp deletion within the *bceB* gene, which encodes a glycosyltransferase. Insertional inactivation of *bceB* in *Burkholderia ambifaria* AMMD results in loss of EPS production on sugar alcohol media. These novel and surprising insights into EPS biosynthesis highlight the metabolic potential of the Bcc and show that a potential virulence factor may not be detected by routine laboratory culture. Our results also highlight a potential hazard in the use of inhaled mannitol as an osmolyte to improve mucociliary clearance in individuals with cystic fibrosis.

Received 6 August 2007

Revised 2 May 2008

Accepted 5 May 2008

INTRODUCTION

The genus *Burkholderia* includes three closely related microbial species that highlight diverse evolutionary adaptation to different niches and hosts. *Burkholderia mallei* is a soliped-specific pathogen that only occasionally infects humans. *Burkholderia pseudomallei* is a free-living soil microbe and the causative agent of the subtropical human disease melioidosis. The species that presently

constitute the *Burkholderia cepacia* complex (Bcc) have multiple roles; they include soil and water saprophytes, rhizosphere parasites, bioremediators, plant growth promoters and plant, animal and human pathogens. Members of the Bcc are particularly associated with life-threatening respiratory infections in patients with chronic granulomatous disease (CGD), and are the most potentially virulent, transmissible and inherently resistant microbes to have emerged as cystic fibrosis (CF) pathogens in recent decades (Govan, 2006; Mahenthiralingam *et al.*, 2005).

Although most species within the Bcc produce a variety of putative virulence factors, the role of these factors in the pathogenesis of human infection is unclear (Mahenthiralingam *et al.*, 2005). Evidence from various model systems (mouse, rat, plant and nematode) suggests

Abbreviations: Bcc, *Burkholderia cepacia* complex; CF, cystic fibrosis; CGD, chronic granulomatous disease; EPS, exopolysaccharide; HPAE-PAD, high-performance anion-exchange chromatography with pulsed amperometric detection; TFA, trifluoroacetic acid.

Three supplementary tables and a supplementary figure are available with the online version of this paper.

that the importance of individual virulence factors, or combinations of factors, depends on the infection model used (Bernier *et al.*, 2003). In addition, studies of Bcc infections in CF patients also suggest a key role of host-pathogen interactions, since clinical outcome in individual patients cannot be predicted even during epidemic outbreaks when multiple patients are infected by the same strain (Govan *et al.*, 1993). The first description of pathogenicity in the Bcc was based on sour skin rot of onion bulbs (Burkholder, 1950). In this study, we returned to this plant host to investigate the onion-associated phenotype of the Bcc, and reveal a link between growth conditions and exopolysaccharide (EPS) production.

EPS is a putative Bcc virulence factor that is involved in persistence of the bacteria in CF lungs (Conway *et al.*, 2004), interactions with antimicrobial peptides (Herasimenka *et al.*, 2005) and the formation of biofilms (Cunha *et al.*, 2004). The EPSs of *Burkholderia* species have recently been comprehensively reviewed (Goldberg, 2007). Recent studies (Zlosnik *et al.*, 2008) have also challenged the previous belief that mucoid, EPS-producing colonial morphotypes of Bcc are rare in both environmental and clinical isolates (Govan & Deretic, 1996). Other studies have shown that mucoid Bcc isolates mostly synthesize one type of EPS, with a highly branched heptasaccharide repeating unit, which was named cepacian (Moreira *et al.*, 2003; Sist *et al.*, 2003). EPS production has been shown to increase when the Bcc are grown in mannitol-rich yeast extract medium (MYEM) (Sage *et al.*, 1990; Zlosnik *et al.*, 2008).

Here we report the novel observation that many Bcc isolates, found to be non-mucoid on typical culture media, produce copious amounts of EPS when onion tissue is provided as the sole nutrient. Chemical and molecular analyses suggest that EPS biosynthesis is strain-specific and that the plant compounds responsible are primarily sugars and sugar alcohols. We show that the EPS phenotype on onion media is associated with the previously described *bce* cluster (Moreira *et al.*, 2003), thought to be involved in cepacian biosynthesis.

METHODS

Bacterial strains and culture conditions. Bcc isolates used in this study are described in Table 1 and include 16 isolates from the two published Bcc strain panels (Coenye *et al.*, 2003; Mahenthiralingam *et al.*, 2000). Additional Bcc strains investigated included *Burkholderia pyrracinia* BTS7, *Burkholderia cenocepacia* BTS2, as well as 19 *Burkholderia multivorans*, 14 *Burkholderia cenocepacia* IIIA and 11 *Burkholderia cenocepacia* IIIB isolates from our collection. Isolates were recovered from storage at -80°C by subculture on nutrient agar (NA; Columbia base agar, Oxoid) and subsequently grown on media composed of 1.5% (w/v) bacteriological agar (Oxoid) containing 2% (w/v) sugars and fractions from various isolation methods below.

Sugars and other related chemicals were purchased from Sigma-Aldrich, BDH AnalaR, MP Biochemicals, Fluka BioChemika or Acros Organics.

Onion maceration. White onion slices were placed in Petri dishes and inoculated with stationary-phase Bcc cultures (10^6 c.f.u.) that had been cultured overnight in 2.5% (w/v) nutrient broth No 2 (Oxoid) with 0.5% (w/v) yeast extract. The onions were left at 30°C for 5 days. The results were assessed by eye, and onion maceration recorded as positive or negative.

Onion extract agar. Peeled white onions (1 kg) were chopped, homogenized in a blender at room temperature, and filtered through muslin. The filtrate was filter-sterilized through a $0.22\text{ }\mu\text{m}$ filter and lyophilized to give a yellow sticky powder (typical yield 62 g). Twenty grams of lyophilized onion extract and 15 g bacteriological agar were made up to 1 l with distilled H_2O , then autoclaved at 121°C for 15 min. Strains were subcultured onto onion agar and incubated for 72 h at 30°C . Mucoidy was recorded on a scale ranging from non-mucoid (–) to very mucoid (+++).

Sugar agar. Sugar agar contained 20 g of the sugar of interest, 2 g yeast extract, and 15 g bacteriological agar dissolved in 1 l distilled H_2O (Sage *et al.*, 1990). The sugars used were as follows: D-fructose, D-galactose, D-mannitol, D-glucose, glycerol, lactose, L-rhamnose, D-mannose, maltose, sucrose, *myo*-inositol, ribitol (adonitol) and D-glucitol (sorbitol). The fructan polysaccharide inulin was also tested.

As a control, isolates were grown on bacteriological agar containing 0.2% (w/v) yeast extract alone.

Reverse-phase chromatography. Twenty millilitres of onion extract, 2% (w/v) in distilled H_2O , was loaded onto a pre-packed C8 column (10 g/60.0 ml, Varian, Anachem) and bound material was eluted stepwise using three concentrations of methanol (20%, 50% and 80%, v/v; Fisher Scientific). Each fraction was lyophilized, dissolved in distilled H_2O and incorporated into 1.5% (w/v) bacteriological agar.

Ethyl acetate partitioning. To separate any lipids, non-polar, non-acidic and polar compounds, the resolubilized onion extract was brought to pH 7.0 using NaOH and partitioned against ethyl acetate (Fisher Scientific) at 1:1 (v/v). The phases were separated, and the aqueous phase was brought to pH 2.0 using HCl and the extraction was repeated. Following both extractions, the organic and aqueous layers were evaporated or lyophilized respectively, and then redissolved in distilled H_2O , pH adjusted to 7.0, and incorporated into 1.5% (w/v) bacteriological agar.

Acid hydrolysis. The aqueous phase residue of the ethyl acetate partitioned onion extract was redissolved in distilled H_2O (50 mg ml^{-1}) and was hydrolysed in 2 M TFA (trifluoroacetic acid; Sigma) at 60°C or 120°C for 1 h.

Paper electrophoresis. To fractionate the extract based on the presence or absence of functional groups, the freeze-dried aqueous phase of the onion extract was weighed and resuspended in distilled H_2O to a final concentration of 50 mg ml^{-1} . One millilitre was loaded at a central origin on a Whatman no.1 paper ($57 \times 42\text{ cm}$). The following standards were added in the margins of the paper: glucose 6-phosphate, glucose, glucosamine, methyl green (Sigma) and orange G (BDH). Electrophoresis was conducted at 1 kV for 20 min, in volatile buffers at pH 2.0 or 6.5 with white spirit or toluene as coolant (Fry, 2000).

Paper chromatography. One millilitre aliquots of each of the redissolved aqueous phase of the onion extract and the TFA-hydrolysed samples (50 mg ml^{-1}) were chromatographed on Whatman no. 1 paper alongside markers (ferulic acid, rhamnose, glucose, lactose, mannose, galactose, fructose, mannitol and glycerol) in butan-1-ol/acetic acid/water (12:3:5, by vol.) for up to 60 h.

Table 1. EPS biosynthesis of *B. cepacia* complex species when grown on agar supplemented with various substrates

All strains tested are from the two published Bcc panels (Coenye *et al.*, 2003; Mahenthiralingam *et al.*, 2000), except for BTS2 and BTS7, which were donated by Paola Cescutti. EPS production was scored on a scale from – (no EPS) to + + + (very mucoid). Mucoid growth described as + + + is shown in Fig. 1(b). YE, yeast extract agar.

Species	Strain	Source	YE	Onion	Glucose	Sucrose	Fructose	Inulin	Glycerol	Mannitol	Glucitol	Ribitol	Inositol	Mannose
<i>B. cepacia</i>	ATCC 25416	Onion	–	++	–	++	+++	–	+++	+++	+++	+++	+++	+
<i>B. cepacia</i>	CEP509	CF	–	–	–	+	++	+	++	++	+	+	+	–
<i>B. multivorans</i>	C1576	CF	–	++	–	+	+++	+	++	+++	+++	+++	+++	–
<i>B. multivorans</i>	ATCC 17616	Soil	–	++	–	+	+++	+	++	+++	+++	+++	+++	–
<i>B. cenocepacia</i>	J2315*	CF	–	–	–	–	–	–	–	–	–	–	–	–
<i>B. cenocepacia</i>	K56-2*	CF	–	–	–	–	–	–	–	–	–	–	–	–
<i>B. cenocepacia</i>	BC7*	CF	–	–	–	–	–	–	–	–	–	–	–	–
<i>B. cenocepacia</i>	PC184	CF	–	++	–	+	++	+	+	++	+	+	+	–
<i>B. cenocepacia</i>	BTS2	CF	–	++	–	+	++	+	+	++	++	+	++	–
<i>B. stabilis</i>	LMG14294	CF	–	–	–	–	+	–	–	+	–	–	–	–
<i>B. vietnamiensis</i>	LMG10929	Rice	–	+++	–	++	+++	+	+++	+++	+	–	+++	++
<i>B. vietnamiensis</i>	PC259	CF	–	++	–	+	+++	+	++	+++	+	–	–	–
<i>B. dolosa</i>	E12	CF	–	++	–	–	+++	–	++	++	–	+	+++	–
<i>B. ambifaria</i>	AMMD	Soil	–	+++	–	+	+++	+	+++	+++	+++	+++	+++	–
<i>B. anthina</i>	W92 ^T	Soil	–	+++	–	–	+++	+	+++	+++	+++	+++	+++	++
<i>B. anthina</i>	C1765	CF	–	+	–	–	+++	–	+++	+++	–	+++	+	+
<i>B. pyrrocinia</i>	BTS7	CF	–	+++	–	+	+++	+++	+++	+++	+++	+++	+++	+
<i>B. pyrrocinia</i>	C1469	CF	–	++	–	++	++	+	++	+++	+++	+++	++	–

*ET12 isolates.

Staining and elution methods. Electrophoretograms and chromatograms were stained with silver nitrate (Fry, 2000) to reveal monosaccharides, oligosaccharides, alditols, saccharinic acids and phenols, and with aniline hydrogen phthalate to reveal monosaccharides and reducing disaccharides. The paper strips of interest from both methods were eluted by a syringe method with distilled H₂O (Eshdat & Mirelman, 1972). The eluted material was incorporated into 1.5 % (w/v) bacteriological agar with 0.2 % (w/v) yeast extract.

High-performance anion-exchange chromatography with pulsed amperometric detection (HPAE-PAD). Twenty microlitres each of the aqueous phases of the onion extract and the TFA-hydrolysed samples (0.1 mg ml⁻¹) were analysed by HPAE-PAD (Dionex). The system consisted of an AS3500 autosampler, GP40 gradient pump, ED40 electrochemical detector, and PC10 pneumatic controller. The amperometry detector cell contained a gold electrode and a pH-Ag/AgCl combination reference electrode. CarboPac MA-1, PA-1 and PA-100 columns and guard columns were used for the separation of alditols, monosaccharides and oligosaccharides, respectively. Eluents, degassed by bubbling with helium, were as follows: MA-1, 600 mM NaOH at 0.4 ml min⁻¹ (isocratic); PA-1, 20 mM NaOH for 3 min, then H₂O for 32 min, then a 0–200 mM NaOH gradient over 10 min (all at 1.0 ml min⁻¹ with post-column addition of base); PA-100, 100 mM NaOH throughout, supplemented with a 0–200 mM NaOAc gradient over 30 min, then 200–800 mM NaOAc over 10 min (all at 1.0 ml min⁻¹). Analytes were identified by comparison of retention times to those of standards and quantified by integration of peak area with Chromeleon software (Dionex).

Investigation of conserved EPS gene clusters in Bcc species. Genome sequences representing five Bcc species were examined to determine if two previously published EPS gene clusters within *B. cenocepacia* J2315, the *bce* gene cluster (Moreira *et al.*, 2003) and the *wcb* gene cluster (Parsons *et al.*, 2003), are conserved across the Bcc. The amino acid sequences for every ORF within each of the two gene clusters within *B. cenocepacia* J2315 (genome sequence available at the Wellcome Trust Sanger Institute; <http://www.sanger.ac.uk/>) were used to search by TBLASTN the following Bcc genome sequences: *Burkholderia ambifaria* AMMD, *Burkholderia vietnamiensis* G4, *B. multivorans* ATCC 17616, *Burkholderia* sp. 383 (all sequences are available at the US Department of Energy Joint Genome Institute; <http://www.jgi.doe.gov/>) and *Burkholderia dolosa* AU0158 (available at the Broad Institute; <http://www.broad.mit.edu/>). TBLASTN searches were performed using default parameters (BLOSUM 62, Word size 3).

RT-PCR analysis of EPS biosynthetic gene clusters. The EPS-producing strain, *B. ambifaria* AMMD, was cultured in 0.2 % (w/v) yeast extract with or without supplementation with 2 % (w/v) mannitol. RNA was extracted from mid-exponential phase cultures (RNeasy Protect Bacteria Mini kit; Qiagen) and DNase I-treated (RNase-free DNase set; Qiagen), prior to reverse transcription with 1.5 µg RNA template, random primers and SuperScript III Reverse Transcriptase (Invitrogen). cDNAs and corresponding non-RT controls were used as template in PCR reactions specific for two distinct *wza* homologues (Bamb_5549 and Bamb_3621) located within two separate putative EPS biosynthetic gene clusters. Primer sequences are available upon request.

PCR analysis of *bceB* gene. Using the *bceB* gene sequence of *B. cenocepacia* J2315, *bceB* homologues were identified within the publicly available genome sequences of *B. ambifaria* AMMD, *B. cepacia* sp. 383, *B. cenocepacia* AU1054, *B. cenocepacia* HI2424, *B. cenocepacia* PC184, *B. vietnamiensis* G4 and *B. dolosa* AU0158. The eight gene sequences were aligned, and PCR primers flanking the location of the previously described 11 bp deletion in *B. cenocepacia* J2315 (Moreira *et al.*, 2003) were designed based on conserved regions

across the eight aligned sequences (For 5'-TGAAGGCGGT[G/C]GCGATCGTC-3'; Rev 5'-TCGAT[G/C]CGCACGTCGTCGAG-3'). For preparation of genomic DNA, one or two bacterial colonies were resuspended in 20 µl lysis solution [0.25 % (w/v) SDS, 0.05 M NaOH] and incubated at 95 °C for 15 min. After brief centrifugation, 180 µl sterile water was added and centrifugation performed at 13 000 g for 5 min. Two microlitres of supernatant containing genomic DNA was used as template in PCR assays. PCRs were performed in a 50 µl volume containing 300 nM forward and reverse primer, 1.5 mM MgCl₂, 260 µM of each dNTP, 4 % (v/v) DMSO, 1 U *Taq* polymerase (Invitrogen) and appropriate manufacturer's reaction buffer. Thermal cycling was performed on a GeneAmp PCR System 9700 (Applied Biosystems) with the following parameters: 94 °C 3 min; 40 cycles of 94 °C (30 s), 60 °C (30 s) and 72 °C (30 s); 72 °C 10 min. PCR products were electrophoresed on a 4 % E-Gel (Invitrogen) and visualized by UV illumination. The *B. cenocepacia* J2315 *bceB* gene sequence harbouring the 11 bp deletion yields a PCR product of 140 bp, compared to 151 bp from the wild-type sequence found in *B. cenocepacia* IST 432 (Videira *et al.*, 2005).

PCR analysis of BCESM and *cbiA*. Genomic DNA was prepared as described above. PCRs were performed as described previously (Mahenthiralingam *et al.*, 1997; Sajjan *et al.*, 1995).

Insertional inactivation of *bceB* in *B. ambifaria* AMMD. Insertional inactivation of *bceB* was performed using the pGPΩTp suicide vector, essentially as described previously (Flannagan *et al.*, 2007). In brief, a 300 bp fragment internal to the *bceB* ORF of *B. ambifaria* AMMD and flanked by *Xba*I and *Eco*RI sites was PCR-amplified and ligated into the corresponding sites in pGPΩTp following appropriate restriction. Resulting plasmids were transformed into *Escherichia coli* GT115 competent cells (InvivoGen) and subsequently introduced into *B. ambifaria* AMMD by triparental mating. Resulting exconjugants were selected using gentamicin (50 mg l⁻¹) and trimethoprim (100 mg l⁻¹), and mutants identified by PCR using a chromosomal-specific primer in conjunction with the vector-specific primer RSF1300 (Flannagan *et al.*, 2007).

RESULTS

EPS production on onion agar

When a lyophilized onion extract was incorporated into agar at 2 % (w/v) as the sole nutrient, members of the Bcc were not only able to grow, but a majority of isolates also produced copious amounts of EPS, as typified by *B. ambifaria* AMMD (Fig. 1a, b). This phenotype is not observed when the Bcc are cultured on *B. cepacia* media (Mast Diagnostics), nutrient agar and other common laboratory media. Induction of EPS biosynthesis on onion agar was observed in all Bcc species investigated, but was not observed in all strains within a species (Table 1 and Supplementary Tables S1, S2 and S3, available with the online version of this paper). The exception was the single strain of *Burkholderia stabilis* available for testing. Of particular interest was the failure of the well-characterized *B. cenocepacia* ET12 representatives J2315, K56-2 and BC7 to produce EPS on onion agar. In addition, there was no correlation between induction of Bcc EPS on onion agar and the ability of individual Bcc strains to cause maceration of onion bulbs (data not shown).

Fig. 1. Growth of *B. ambifaria* AMMD on (a) nutrient agar (non-mucoid); (b) onion agar (mucoid), and (c) comparison of AMMD *bceB* mutant (left) and AMMD wild-type (right) on mannitol agar.

The onion factor

Attempts were made to identify the 'onion factor' responsible for inducing EPS biosynthesis by use of standard biochemical methods for extracting and fractionating phytochemicals. The causative factor was retained on drying *in vacuo*, and remained in the aqueous phase after partition with ethyl acetate at pH 7.0 and pH 2.0. When further physicochemical analyses discounted the role of proteins and lipids, attention was turned to the carbohydrate content of onion extract. After preparative paper electrophoresis in buffers at pH 2.0 and 6.5, the only biological activity recovered from paper strips co-migrated with the standard glucose, indicating the absence of ionizable functional groups such as phosphate, acid or amine (data not shown). Analytical paper chromatography and HPAE-PAD identified the major carbohydrate components as sucrose, glucose, fructose and fructans (Fig. 2). The HPAE-PAD chromatograms in Fig. 2 show characteristic peaks of glucose, fructose and sucrose. Sucrose breaks down to fructose and glucose upon mild hydrolysis, as do the fructans to fructose. Fructose breaks down under complete hydrolysis as expected for a ketose sugar, whilst the aldose sugar glucose remains stable. The ability of these and related compounds to stimulate EPS biosynthesis in Bcc was then investigated (Table 1). Glycerol and mannitol were included as these sugar alcohols have previously been noted to enhance EPS biosynthesis in *Pseudomonas aeruginosa* (Whitchurch *et al.*, 1996) and the Bcc (Sage *et al.*, 1990; Zlosnik *et al.*, 2008). Glucitol was included because of its close degradative relationship with fructose and mannitol (Allenza *et al.*, 1982). These experiments showed that within a particular Bcc species, EPS biosynthesis was strain-specific and that the most potent inducers of EPS were fructose and all alditols tested, most significantly mannitol and glucitol, as well as the cyclitol *myo*-inositol. Importantly, the profile of EPS biosynthesis production with these sugars was similar to that observed with onion extract (Table 1). EPS biosynthesis was not observed on agar containing yeast extract alone, nor in the

presence of glucose (Table 1), galactose, lactose or maltose (data not shown) with any Bcc strains tested.

Investigation of the molecular basis for EPS biosynthesis

With the exception of the single *B. stabilis* strain tested, all Bcc species were shown to be capable of producing EPS when grown on onion agar (Table 1 and Supplementary Tables S1–S3), suggesting the presence of a conserved EPS biosynthetic gene cluster. Therefore genome sequences representing five Bcc species (*B. ambifaria*, *B. multivorans*, *B. vietnamiensis*, *B. dolosa* and *Burkholderia* sp. 383) were examined to determine if two putative EPS gene clusters within *B. cenocepacia* J2315, the *bce* gene cluster (Moreira *et al.*, 2003) and the *wcb* gene cluster (Parsons *et al.*, 2003), are conserved across the Bcc. The *wcb* gene cluster was found to be poorly conserved, with between one-third and one-half of J2315 ORFs having no direct homologues within the species examined (data not shown). Notably, within the EPS-producing *B. ambifaria* AMMD strain, over half of the J2315 *wcb*-associated ORFs have no homologues, and the remaining homologous ORFs are not organized within a gene cluster. In contrast, the *bce* gene cluster was conserved across all species examined, in terms of both sequence homology and organization of ORFs (Supplementary Fig. S1). In the course of these genome comparisons, a third putative polysaccharide biosynthetic gene cluster was observed on chromosome 2 of *B. cenocepacia* J2315. This cluster encodes two putative EPS transporter proteins (BCAM1330 and BCAM1331), an acyltransferase (BCAM1333), several glycosyltransferases (BCAM1335, BCAM1337, BCAM1338), a polysaccharide biosynthesis protein (BCAM1336) and a mannose-6-phosphate isomerase (BCAM1340). This cluster is conserved amongst several Bcc species, albeit to a lesser extent than the *bce* gene cluster (data not shown). In the EPS-producing *B. ambifaria* AMMD, this gene cluster maps to ORFs Bamb_3621 through to Bamb_3629.

Fig. 2. HPAE-PAD chromatograms: Carbowac PA-100 column separation of sugars in onion extract. (a) Crude onion extract; (b) onion extract hydrolysed by 2 M TFA 60 °C 1 h; (c) onion extract hydrolysed by 2 M TFA 120 °C 1 h. Glucose, fructose and sucrose peaks were clearly identified based on standards (not shown), and hydrolysis pattern. Peaks with retention times between 12 and 36 min appear to be oligosaccharides of fructose based on their degradation to fructose under mild hydrolysis.

To investigate which polysaccharide gene cluster is induced by growth on mannitol, the expression of representative genes from two distinct EPS gene clusters was assessed in *B. ambifaria* AMMD grown in the presence and absence of mannitol. The genes studied each encode homologues of the Wza EPS export protein: Bamb_5549 of the *bce* gene cluster (equivalent to *bceE*, BCAM0858 of *B. cenocepacia* J2315), and Bamb_3621 of the novel putative polysaccharide gene cluster described above (BCAM1330 of *B. cenocepacia* J2315). As shown in Fig. 3, expression of Bamb_3621 was not observed under either growth condition. In contrast, expression of the *bceE* homologue (Bamb_5549) was clearly induced by the presence of mannitol. We therefore focused on the *bce* gene cluster to investigate why some Bcc isolates, most notably those of the *B. cenocepacia* ET12 lineage, failed to produce EPS under any growth conditions.

A PCR assay was designed to screen isolates for an 11 bp deletion in the *bceB* gene that has been suggested to be responsible for loss of EPS production in the CF isolate *B.*

cenocepacia J2315 (Moreira *et al.*, 2003). Of the panel of strains shown in Table 1, only the *B. cenocepacia* ET12 isolates J2315, K56-2 and BC7 harbour the 11 bp deletion (data not shown). This result prompted us to test a panel of *B. cenocepacia* IIIA strains containing both ET12 and non-ET12 isolates. There was a clear correlation between the 11 bp deletion and the presence of both *cblA* (cable pilus) and BCESM (*B. cepacia* epidemic strain marker), indicating that this deletion is a conserved feature within the ET12 lineage (Fig. 4). In our study, with the exception of strain E3051, the presence of the deletion correlated with the lack of EPS production in all *B. cenocepacia* IIIA isolates examined (Fig. 4; Supplementary Table S1). Furthermore, the 11 bp deletion in the *bceB* gene was not observed in any *B. cenocepacia* IIIB or *B. multivorans* isolates studied (see Supplementary Tables S2 and S3).

Using proven methods for the complementation of gene function in *B. cenocepacia* (Ortega *et al.*, 2005), attempts were made to complement BceB function within *B. cenocepacia* K56-2 by introducing a wild-type *bceB* ORF

Fig. 3. Induction of the *bce* gene cluster by mannitol. RT-PCR analysis was performed to assess the expression of two distinct *wza* homologues (*bceE* and Bamb_3621) of *B. ambifaria* AMMD in the absence and presence of mannitol. The *bceE* gene (Bamb_5549) is located within the *bce* gene cluster, whilst Bamb_3621 is located within a distinct putative EPS biosynthetic gene cluster. Expression was assessed in yeast extract (YE) and yeast extract supplemented with 2 % (w/v) mannitol (YE+Man). Expression of Bamb_3621 was not detected in YE or YE+Man. In contrast, *bceE* expression was strongly induced by growth in YE+Man. For each sample, RT and non-RT reactions (+/-) are shown alongside each other. Genomic DNA positive controls are shown for each gene (*bceE* +ve, 3621 +ve).

Fig. 4. *bceB* PCR analysis of representative *B. cenocepacia* IIIA isolates. For each isolate, EPS production is indicated as mucoid (M) or non-mucoid (NM) alongside PCR analysis of *bceB*, and the epidemic strain markers *cbIA* and BCESM. All PCR products are pictured alongside the 1 kb Plus DNA Ladder (Invitrogen). NTC, no template control. The *bceB* PCR assay yields a 151 bp product from wild-type sequence and a 140 bp product from sequence harbouring an 11 bp deletion (as previously reported within J2315). With one exception (E3051), lack of EPS production within *B. cenocepacia* IIIA isolates correlated with the 11 bp deletion within *bceB*. This deletion was only observed within strains that were positive for both *cbIA* and BCESM, indicating it to be a conserved feature of the ET12 lineage. The *bceB* deletion was not observed in a strain harbouring just one of the two epidemic strain markers (E1001).

amplified from *B. cenocepacia* PC184. Whilst expression of 6His-tagged BceB was detected in *E. coli* C41(DE3) cells, previously shown to support expression of membrane-bound proteins (Miroux & Walker, 1996), we were unable to detect expression within *B. cenocepacia* K56-2 (data not shown). Consequently, we chose to disrupt the *bceB* ORF of an EPS-producer. Insertional inactivation of the *bceB* gene resulted in loss of EPS production in *B. ambifaria* AMMD when grown on mannitol agar (Fig. 1c) and on onion agar (data not shown).

DISCUSSION

In this study, we returned to the original Bcc host and report that Bcc isolates previously considered non-mucoid produce copious amounts of EPS when onion tissue is provided as the sole nutrient, highlighting the metabolic potential of this group of organisms. This novel and surprising observation is not species-specific, and is exhibited by isolates of both clinical and environmental origin. The onion components responsible for EPS induction are primarily the carbohydrates sucrose, fructose and fructans. Additionally, all alcohol sugars tested are able to induce EPS production, in particular mannitol and glucitol.

The reasons why EPS biosynthesis is readily induced by onion extracts and, in particular, by fructose and the hexitol sugars mannitol and glucitol require further investigation. An ability to respond to inulin was confined to certain strains, all of which responded well to fructose. This suggests that the fructan polysaccharide was only active in bacteria that can hydrolyse it to fructose. The ability of one strain (BTS7) to respond particularly strongly to inulin could possibly be due to its enhanced ability to hydrolyse inulin to fructose. Interestingly, the similar profile of EPS induction shown by fructose and alcohol sugars agrees well with a previous study which showed that the initial steps of utilization of these sugars in the Bcc differ from those in most other pseudomonads. Growth of Bcc on fructose involves active transport followed by fructokinase conversion to fructose 6-phosphate then degradation via the Entner–Doudoroff pathway (Allenza *et al.*, 1982). Growth on mannitol and glucitol also requires this pathway following active uptake and intracellular oxidation to fructose.

The role of Bcc EPS as a putative virulence factor is unclear. The influence of mannitol and other alcohol sugars on EPS production suggests that the prevalence of EPS in the Bcc may have been underestimated. In addition, given the variety of human, animal, plant and other models studied, the role of EPS may be influenced by the host involved and the route of administration. EPS has been associated with altered Bcc clearance in a mouse model of infection (Conway *et al.*, 2004), and EPS-deficient Bcc mutants displayed reduced mortality within a CGD mouse model (Sousa *et al.*, 2007a). Similarly, a role for EPS in persistence

in human airways is suggested by its capacity to scavenge reactive oxygen species and inhibit neutrophil chemotaxis (Bylund *et al.*, 2006). In our study, EPS biosynthesis did not correlate with the ability to cause onion rot, which is perhaps to be expected since pectinases rather than EPS are likely to play the major role in the maceration of plant tissue. Cunha *et al.* (2004) did not observe a clear correlation between EPS biosynthesis *in vitro* and the ability of Bcc strains to establish chronic infections within the CF lung. Recent evidence, however, suggests a subtle and unexpected role for Bcc EPS in CF lung infection. Consistent with our findings, Zlosnik *et al.* (2008) reported that isolates of *B. cenocepacia*, the most virulent Bcc species, are most frequently non-mucoid. They also observed a mucoid to non-mucoid conversion in sequential isolates of Bcc from chronically infected CF patients. This apparent loss of mucoidy *in vivo*, and its absence in the virulent *B. cenocepacia* ET12 lineage, provides an intriguing contrast with the characteristic non-mucoid to mucoid conversion observed with alginate-producing *P. aeruginosa*. Zlosnik and colleagues suggest that Bcc EPS could be responsible for the persistence of Bcc in CF airways whilst loss of EPS leads to increased disease severity.

In our investigation of EPS biosynthesis determinants, we focused on the highly conserved *bce* gene cluster. Previously described by Moreira *et al.* (2003), the *bce* gene cluster has had several of its encoded proteins characterized (Ferreira *et al.*, 2007; Sousa *et al.*, 2007b; Videira *et al.*, 2005) and is thought to be involved in cepacian biosynthesis. In the present study, we demonstrated induction of the *bce* gene cluster by mannitol, and found a clear correlation between the inability of representatives of the *B. cenocepacia* ET12 lineage to produce EPS and the presence of an 11 bp deletion within the *bceB* gene, originally described within the genome sequence of *B. cenocepacia* J2315 (Moreira *et al.*, 2003). Consistent with this correlation, insertional inactivation of *bceB*, which encodes a glycosyltransferase, resulted in the loss of EPS production by *B. ambifaria* AMMD when grown on onion media. Combined, these observations highlight the pivotal role of the *bce* gene cluster in onion-induced EPS biosynthesis, and suggest that the observed EPS is cepacian. However, mutations elsewhere within the *bce* gene cluster, or in other EPS-related gene clusters, must be responsible for the lack of EPS biosynthesis in *B. cenocepacia* strain E3051 (Supplementary Table S1), and in other EPS-negative Bcc strains in our study which lack the 11 bp deletion in *bceB* (Supplementary Tables S2 and S3).

The ability of hexoses and hexitols, in particular mannitol, to enhance EPS biosynthesis in the *B. cepacia* complex has disturbing implications for therapeutic intervention in CF. Recent attempts to improve airway clearance with hypertonic saline 5% (w/v) have been handicapped by the problem of salty taste and the salt-sensitive nature of many antimicrobial peptides. Thus, attention has turned to the use of non-ionic osmolytes, including inhaled mannitol

(Daviskas *et al.*, 2008; Robinson *et al.*, 1999; Wills, 2007), which is marketed as Bronchitol (Pharmaxis). Robinson and colleagues acknowledged that the majority of *P. aeruginosa* and Bcc isolates are able to utilize mannitol as a carbon and energy source. However, they felt that the nutritional influence of mannitol as a therapeutic osmolyte would be minimal given the abundance of other nutrients already present in CF respiratory secretions. On a cautionary note, they state that this potential problem would need to be confirmed by quantitative microbiology. The potential induction of virulence determinants during osmolyte therapy has to our knowledge not been considered. Our results also provide justification for the continued exclusion of CF individuals known to be infected with Bcc from ongoing trials of inhaled mannitol (<http://clinicaltrials.gov>; identifier NCT00117208 and NCT00251056).

ACKNOWLEDGEMENTS

We thank Catherine Doherty (University of Edinburgh, UK) for the strain panels, Rosanna Hennessy (University of Edinburgh, UK) for the RT-PCR data, Paola Cescutti (Università di Trieste, Italy) for *Burkholderia pyrrocinia* BTS7 and *B. cenocepacia* BTS2, and Miguel Valvano (University of Western Ontario, Canada) for the suicide vector pGP Ω Tp and complementation vector pSCRhaB3. S.J.B., A.R.B., J.R.W.G. and D.J.C. thank the CF Trust and the Big Lottery Fund, D.J.C. and D.J.C. also thank the Royal Society of Edinburgh, and S.C.F. and B.R.M. thank the Biotechnology and Biological Sciences Research Council for financial support.

REFERENCES

- Allenza, P., Lee, Y. N. & Lessie, T. G. (1982). Enzymes related to fructose utilization in *Pseudomonas cepacia*. *J Bacteriol* **150**, 1348–1356.
- Bernier, S. P., Silo-Suh, L., Woods, D. E., Ohman, D. E. & Sokol, P. A. (2003). Comparative analysis of plant and animal models for characterization of *Burkholderia cepacia* virulence. *Infect Immun* **71**, 5306–5313.
- Burkholder, W. (1950). Sour skin, a bacterial rot of onion bulbs. *Phytopathology* **40**, 115–117.
- Bylund, J., Burgess, L. A., Cescutti, P., Ernst, R. K. & Speert, D. P. (2006). Exopolysaccharides from *Burkholderia cenocepacia* inhibit neutrophil chemotaxis and scavenge reactive oxygen species. *J Biol Chem* **281**, 2526–2532.
- Coenye, T., Vandamme, P., LiPuma, J. J., Govan, J. R. & Mahenthalingam, E. (2003). Updated version of the *Burkholderia cepacia* complex experimental strain panel. *J Clin Microbiol* **41**, 2797–2798.
- Conway, B. A., Chu, K. K., Bylund, J., Altman, E. & Speert, D. P. (2004). Production of exopolysaccharide by *Burkholderia cenocepacia* results in altered cell-surface interactions and altered bacterial clearance in mice. *J Infect Dis* **190**, 957–966.
- Cunha, M. V., Sousa, S. A., Leitao, J. H., Moreira, L. M., Videira, P. A. & Sa-Correia, I. (2004). Studies on the involvement of the exopolysaccharide produced by cystic fibrosis-associated isolates of the *Burkholderia cepacia* complex in biofilm formation and in persistence of respiratory infections. *J Clin Microbiol* **42**, 3052–3058.

- Daviskas, E., Anderson, S. D., Eberl, S. & Young, I. H. (2008). Effect of increasing doses of mannitol on mucus clearance in patients with bronchiectasis. *Eur Respir J* 31, 765–772.
- Eshdat, Y. & Mirelman, D. (1972). An improved method for the recovery of compounds from paper chromatograms. *J Chromatogr* 65, 458–459.
- Ferreira, A. S., Leitao, J. H., Sousa, S. A., Cosme, A. M., Sa-Correia, I. & Moreira, L. M. (2007). Functional analysis of *Burkholderia cepacia* genes *bceD* and *bceF*, encoding a phosphotyrosine phosphatase and a tyrosine autokinase, respectively: role in exopolysaccharide biosynthesis and biofilm formation. *Appl Environ Microbiol* 73, 524–534.
- Flannagan, R. S., Aubert, D., Kooi, C., Sokol, P. A. & Valvano, M. A. (2007). *Burkholderia cenocepacia* requires a periplasmic HtrA protease for growth under thermal and osmotic stress and for survival in vivo. *Infect Immun* 75, 1679–1689.
- Fry, S. C. (2000). *The Growing Plant Cell Wall: Chemical and Metabolic Analysis*. Caldwell, NJ: The Blackburn Press.
- Goldberg, J. B. (2007). Polysaccharides of *Burkholderia* spp. In *Burkholderia: Molecular Microbiology and Genomics*, pp. 93–110. Edited by T. Coenye & P. Vandamme. Norwich, UK: Horizon Bioscience.
- Govan, J. R. W. (2006). *Burkholderia cepacia* complex and *Stenotrophomonas maltophilia*. In *Cystic Fibrosis in the 21st Century: Progress in Respiratory Research*, pp. 145–152. Edited by A. Bush, E. W. F. W. Alton, J. C. Davies, U. Griesenbach & A. Jaffe. Basel: Karger.
- Govan, J. R. & Deretic, V. (1996). Microbial pathogenesis in cystic fibrosis: mucoid *Pseudomonas aeruginosa* and *Burkholderia cepacia*. *Microbiol Rev* 60, 539–574.
- Govan, J. R., Brown, P. H., Maddison, J., Doherty, C. J., Nelson, J. W., Dodd, M., Greening, A. P. & Webb, A. K. (1993). Evidence for transmission of *Pseudomonas cepacia* by social contact in cystic fibrosis. *Lancet* 342, 15–19.
- Herasimenka, Y., Benincasa, M., Mattiuzzo, M., Cescutti, P., Gennaro, R. & Rizzo, R. (2005). Interaction of antimicrobial peptides with bacterial polysaccharides from lung pathogens. *Peptides* 26, 1127–1132.
- Mahenthalingam, E., Simpson, D. A. & Speert, D. P. (1997). Identification and characterization of a novel DNA marker associated with epidemic *Burkholderia cepacia* strains recovered from patients with cystic fibrosis. *J Clin Microbiol* 35, 808–816.
- Mahenthalingam, E., Coenye, T., Chung, J. W., Speert, D. P., Govan, J. R., Taylor, P. & Vandamme, P. (2000). Diagnostically and experimentally useful panel of strains from the *Burkholderia cepacia* complex. *J Clin Microbiol* 38, 910–913.
- Mahenthalingam, E., Urban, T. A. & Goldberg, J. B. (2005). The multifarious, multireplicon *Burkholderia cepacia* complex. *Nat Rev Microbiol* 3, 144–156.
- Miroux, B. & Walker, J. E. (1996). Over-production of proteins in *Escherichia coli*: mutant hosts that allow synthesis of some membrane proteins and globular proteins at high levels. *J Mol Biol* 260, 289–298.
- Moreira, L. M., Videira, P. A., Sousa, S. A., Leitao, J. H., Cunha, M. V. & Sa-Correia, I. (2003). Identification and physical organization of the gene cluster involved in the biosynthesis of *Burkholderia cepacia* complex exopolysaccharide. *Biochem Biophys Res Commun* 312, 323–333.
- Ortega, X., Hunt, T. A., Loutet, S., Vinion-Dubiel, A. D., Datta, A., Choudhury, B., Goldberg, J. B., Carlson, R. & Valvano, M. A. (2005). Reconstitution of O-specific lipopolysaccharide expression in *Burkholderia cenocepacia* strain J2315, which is associated with transmissible infections in patients with cystic fibrosis. *J Bacteriol* 187, 1324–1333.
- Parsons, Y. N., Banasko, R., Detsika, M. G., Duangsonk, K., Rainbow, L., Hart, C. A. & Winstanley, C. (2003). Suppression-subtractive hybridisation reveals variations in gene distribution amongst the *Burkholderia cepacia* complex, including the presence in some strains of a genomic island containing putative polysaccharide production genes. *Arch Microbiol* 179, 214–223.
- Robinson, M., Daviskas, E., Eberl, S., Baker, J., Chan, H. K., Anderson, S. D. & Bye, P. T. (1999). The effect of inhaled mannitol on bronchial mucus clearance in cystic fibrosis patients: a pilot study. *Eur Respir J* 14, 678–685.
- Sage, A., Linker, A., Evans, L. R. & Lessie, T. G. (1990). Hexose phosphate metabolism and exopolysaccharide formation in *Pseudomonas cepacia*. *Curr Microbiol* 20, 191–198.
- Sajjan, U. S., Sun, L., Goldstein, R. & Forstner, J. F. (1995). Cable (*cbl*) type II pili of cystic fibrosis-associated *Burkholderia (Pseudomonas) cepacia*: nucleotide sequence of the *cblA* major subunit pilin gene and novel morphology of the assembled appendage fibers. *J Bacteriol* 177, 1030–1038.
- Sist, P., Cescutti, P., Skerlavaj, S., Urbani, R., Leitao, J. H., Sa-Correia, I. & Rizzo, R. (2003). Macromolecular and solution properties of cepacian: the exopolysaccharide produced by a strain of *Burkholderia cepacia* isolated from a cystic fibrosis patient. *Carbohydr Res* 338, 1861–1867.
- Sousa, S. A., Ulrich, M., Bragonzi, A., Burke, M., Worlitzsch, D., Leitão, J. H., Meisner, C., Eberl, L., Sá-Correia, I. & Döring, G. (2007a). Virulence of *Burkholderia cepacia* complex strains in gp91phox^{-/-} mice. *Cell Microbiol* 9, 2817–2825.
- Sousa, S. A., Moreira, L. M., Wopperer, J., Eberl, L., Sá-Correia, I. & Leitão, J. H. (2007b). The *Burkholderia cepacia* *bceA* gene encodes a protein with phosphomannose isomerase and GDP-D-mannose pyrophosphorylase activities. *Biochem Biophys Res Commun* 353, 200–206.
- Videira, P. A., Garcia, A. P. & Sa-Correia, I. (2005). Functional and topological analysis of the *Burkholderia cenocepacia* priming glucosyl-transferase BceB, involved in the biosynthesis of the cepacian exopolysaccharide. *J Bacteriol* 187, 5013–5018.
- Whitchurch, C. B., Alm, R. A. & Mattick, J. S. (1996). The alginate regulator AlgR and an associated sensor FimS are required for twitching motility in *Pseudomonas aeruginosa*. *Proc Natl Acad Sci U S A* 93, 9839–9843.
- Wills, P. J. (2007). Inhaled mannitol in cystic fibrosis. *Expert Opin Investig Drugs* 16, 1121–1126.
- Zlosnik, J. E., Hird, T. J., Fraenkel, M. C., Moreira, L. M., Henry, D. A. & Speert, D. P. (2008). Differential mucoid exopolysaccharide production by members of the *Burkholderia cepacia* complex. *J Clin Microbiol* 46, 1470–1473.

Edited by: P. Cornelis