Characterization of the pathogenicity island protein PdpA and its role in the virulence of *Francisella novicida*

Crystal L. Schmerk,¹ Barry N. Duplantis,¹ Diana Wang,¹ Robert D. Burke,¹ Alicia Y. Chou,² Karen L. Elkins,² Jagjit S. Ludu¹ and Francis E. Nano¹

Correspondence

Francis E. Nano
fnano@uvic.ca

¹Department of Biochemistry and Microbiology, University of Victoria, Victoria, BC, Canada

²Center for Biologics Evaluation and Research, Food and Drug Administration, Rockville, MD, USA

Received 24 October 2008

Revised 29 January 2009

Accepted 30 January 2009

Francisella tularensis is a highly virulent, intracellular pathogen that causes the disease tularaemia. A research surrogate for *F. tularensis* is *Francisella novicida*, which causes a tularaemia-like disease in mice, grows similarly in macrophages, and yet is unable to cause disease in humans. Both *Francisella* species contain a cluster of genes referred to as the *Francisella* pathogenicity island (FPI). Pathogenicity determinant protein A (PdpA), encoded by the *pdpA* gene, is located within the FPI and has been associated with the virulence of *Francisella* species. In this work we examined the properties of PdpA protein expression and localization as well as the phenotype of a *F. novicida* *pdpA* deletion mutant. Monoclonal antibody detection of PdpA showed that it is a soluble protein that is upregulated in iron-limiting conditions and undetectable in an *mglA* or *mglB* mutant background. Deletion of *pdpA* resulted in a strain that was highly attenuated for virulence in chicken embryos and mice.

INTRODUCTION

Francisella tularensis is a Category A bioterrorism threat and is the causative agent of the zoonotic disease tularaemia. The closely related bacterium *Francisella novicida* is unable to cause disease in healthy individuals; however, it causes a disease in mice that is very similar to human tularaemia and, as such, is a useful tool for studying *Francisella* virulence factors. Both *F. tularensis* and *F. novicida* grow vigorously in macrophages, with a life cycle that includes transient localization in a phagosome followed by growth in the host-cell cytosol (Anthony *et al.*, 1991; Clemens *et al.*, 2004; Fortier *et al.*, 1994; Golovliov *et al.*, 2003a; Proctor *et al.*, 1975). Little information is available concerning the function of virulence factors that contribute to *F. tularensis* infection. Many of the genes that have been shown to be necessary for intracellular growth and virulence are located within the recently discovered *Francisella* pathogenicity island (FPI, see Fig. 1) (Nano *et al.*, 2004; Nano & Schmerk, 2007). The FPI is about 28 kb in length and contains 16–19 genes, depending on the biotype and strain. Recent bioinformatics analysis indicates that several genes within the FPI encode products that are homologues of components of type VI secretion systems (T6SSs) (de Bruin *et al.*, 2007;

Ludu *et al.*, 2008a). *F. novicida* contains one copy of the FPI whereas *F. tularensis* strains that have been examined contain two copies of the genomic island (Nano & Schmerk, 2007).

Disruptions of a number of genes within the FPI have been shown to severely reduce intracellular bacterial growth (de Bruin *et al.*, 2007; Golovliov *et al.*, 2003b; Gray *et al.*, 2002; Lai *et al.*, 2004; Lauriano *et al.*, 2004; Nano *et al.*, 2004; Santic *et al.*, 2005; Tempel *et al.*, 2006; Weiss *et al.*, 2007). However, in most cases the phenotype of mutants could not be unequivocally demonstrated to be the result of the inactivation of a specific gene, since polarity effects were not studied nor genetic complementation performed. It is probable that most FPI genes contribute to intracellular growth and virulence, and deletion mutagenesis and complementation studies have provided strong evidence that *iglA* and *iglC*, two of the genes in the *anmK-iglD* operon (see Fig. 1) are absolutely required for intramacrophage growth.

Regulation of FPI gene expression has been linked to several factors. MglA is the best studied of the *Francisella* global regulators (Baron & Nano, 1998; Brotcke *et al.*, 2006; Charity *et al.*, 2007). Studies of gene regulation in *Francisella* have revealed that FPI-encoded mRNA species are depressed in an *mglA* mutant. The MglA protein has been shown to bind RNA polymerase complexes, but its direct role in affecting mRNA levels of FPI genes has not yet been established. Low iron concentrations also increase

Abbreviations: FPI, *Francisella* pathogenicity island; T6SS, type VI secretion system.

The GenBank/EMBL/DDBJ accession number for the sequence of the genomic regions in the *pdpA* deletions is EU810409.

Fig. 1. Deletion of *pdpA*. (a) The *pdpA* gene is the first cistron in a low-G+C operon of the *Francisella* pathogenicity island (FPI). A newly revised nomenclature is used to show the two convergent operons in the FPI. (b) Primers that amplify regions within the *pdpA* gene were used to screen wild-type (WT) *F. novicida*, the $\Delta pdpA$ mutant and the *in cis* complement $\Delta pdpA/SKX::pdpA$. (c) The exact nucleotides removed to create the $\Delta pdpA$ mutant are shown. The in-frame deletion preserves the last 21 nucleotides of *pdpA*, which contain the ribosome-binding site (shown in italics) of the downstream gene *pdpB*.

the level of mRNA for several FPI-encoded mRNA transcripts (Deng *et al.*, 2006). Mass spectroscopy analysis has shown that IglA, IglB, IglC, IglD and PdpB are all increased in low-iron conditions (Deng *et al.*, 2006; Lenco *et al.*, 2007). Immunoblot analysis has shown that both iron concentration and MglA play a role in IglA expression. It is clear that other regulatory proteins, such as FevR, SspA and PmrA, also affect expression of FPI genes (Brotcke & Monack, 2008; Charity *et al.*, 2007; Mohapatra *et al.*, 2007).

The *pdpA* gene is 2.46 kb in size, representing one of the largest ORFs in the FPI (Nano & Schmerk, 2007). A gene replacement in *pdpA* was made as part of the initial characterization of the FPI, and the resulting mutant was unable to grow in macrophages and was avirulent in mice (Nano *et al.*, 2004). We now know that the replacement mutant had polar effects on downstream genes, resulting in decreased expression of PdpB and increased attenuation in chicken embryos (this work). It is likely that the *in trans* complementing plasmid recreated the intact operon by integration into the chromosome. In this paper we examine

the expression and characteristics of the PdpA protein as well as the phenotype of a non-polar deletion mutant of *pdpA*.

METHODS

Bacterial strains and growth conditions. The bacterial strains and plasmids used in this study are listed in Table 1. *Escherichia coli* strains were grown using Luria Bertani (LB) broth or agar supplemented with 250 μ g ampicillin ml^{-1} , 30 μ g kanamycin ml^{-1} or 100 μ g erythromycin ml^{-1} , as required. *F. novicida* and *F. tularensis* strains were grown using trypticase soy agar or broth supplemented with 0.1% (w/v) cysteine (TSAC, TSBC). When necessary, 30 μ g erythromycin ml^{-1} or 15 μ g kanamycin ml^{-1} was added. For negative selection in deletion mutagenesis experiments, sucrose was added to TSAC to a final concentration of 10% (w/v).

Mutagenesis and complementation. To create the *pdpA* deletion mutant strain the upstream and downstream regions flanking the *pdpA* gene were joined using fusion overlap PCR and the resulting amplicon was cloned into pWSK29 to create pWSK29:: $\Delta pdpA$. The last 22 bp of *pdpA* were left to preserve the ribosome-binding site of

Table 1. Bacterial strains and plasmids used in this study

Strain or plasmid	Characteristics	Source or reference
Strains		
U112	Wild-type <i>F. novicida</i>	ATCC 15482
JLO	U112 with deletion in FTN_1390, where SKX vector inserts; identical growth and virulence with respect to U112	Ludu <i>et al.</i> (2008b)
GB2	U112 with point mutation in global virulence regulator, <i>mgla</i>	Baron & Nano (1998)
GB6	U112, <i>mgIB::mTn10Km^R</i>	Baron & Nano (1998)
SC92	U112, O-antigen gene::Em ^R	Cowley <i>et al.</i> (2000)
<i>F. tularensis</i> subsp. <i>holarctica</i> LVS	Live vaccine strain, type B biotype	ATCC 29684
<i>F. tularensis</i> subsp. <i>tularensis</i> B38	Attenuated type A biotype strain	ATCC 6223
$\Delta pdpA$	JLO with a deletion of <i>pdpA</i>	This study
$\Delta pdpA$ /SKX:: <i>pdpA</i>	$\Delta pdpA$ complemented with the integrating pJL-SKX:: <i>pdpA</i> construct	This study
$\Delta pdpA$ -LPS	$\Delta pdpA$, O-antigen gene::Em ^R from SC92	This study
NZ9	U112, <i>pdpA::Em^R</i> gene replacement mutant.	Laboratory strain
$\Delta pdpA$ /SKX:: <i>pdpA</i> -LPS	$\Delta pdpA$ /SKX:: <i>pdpA</i> , O-antigen gene::Em ^R from SC92	This study
<i>E. coli</i> DH5 α	F ⁻ <i>supE44</i> Δ (<i>lacZYA-argF</i>)U169 (ϕ 80 <i>lacZ</i> AM15) <i>hsdR17</i> (r _K ⁻ m _K ⁺) <i>recA1</i> <i>endA1</i> <i>gyrA96</i> <i>thi-1</i> <i>relA1</i> λ^-	Invitrogen
<i>E. coli</i> BL21 (DE3) pLysS	B F ⁻ <i>dcm ompT hsdS</i> (r _B ⁻ m _B ⁻) <i>gal</i> λ (DE3) [pLysS Cm ^r]	Stratagene
Plasmids		
pWSK29	Low-copy cloning vector, Ap ^r	Wang & Kushner (1991)
pJL-SKX	pWSK29 with integrating SKX cassette, Km ^r , Amp ^r	Ludu <i>et al.</i> (2008b)
pJL-ES-X	pWSK29 containing an <i>ermCsacB</i> cassette with flanking <i>XhoI</i> restriction sites	Ludu <i>et al.</i> (2008b)
pWSK29:: $\Delta pdpA$	A clone containing the fused flanking regions of <i>pdpA</i> ; used to create the <i>F. novicida</i> $\Delta pdpA$ mutant	This study
pJL-SKX:: <i>pdpA</i>	An integrating complementation construct containing the full <i>pdpA</i> gene and its predicted promoter region (522 bp upstream of <i>pdpA</i> start)	This study
pET 28a	Expression vector with 6 $\times$ His and T7 tags	Invitrogen
pET 28a:: <i>pdpA</i> -C	Clone expressing amino acids 405–783 of PdpA with a carboxyterminal 6 $\times$ His tag. Used to create the monoclonal antibody against PdpA	This study

pdpB (see Fig. 1). The insert from pWSK29:: $\Delta pdpA$ was excised using *XhoI*, ligated to the *XhoI*-*XhoI* *sacB*-Em^R cassette from pJL-ES-X, and the ligation reaction was used to transform *F. novicida* JLO. The integration and excision of the *sacB*-Em^R cassette was used to create a markerless deletion of the *pdpA* gene as previously described (Ludu *et al.*, 2008a). The *F. novicida* JLO strain was used as the wild-type strain in order to maintain an isogenic background for mutants and complemented strains, as the integrating plasmid used for complementation inserts at the non-essential locus FTN_1390 that is deleted in the JLO strain. The $\Delta pdpA$ mutant was complemented using the kanamycin integration vector pJL-SKX (Ludu *et al.*, 2008b) that inserts into the chromosome via a double crossover recombination event. The full-length *pdpA* gene and the promoter region, 522 bp upstream of *pdpA*, were ligated to pJL-SKX. The vector was then linearized and used to transform the *pdpA* deletion mutants, creating *in cis* complementation strains. The sequence of the genomic regions in the *pdpA* deletions has been deposited with GenBank and assigned the number EU810409. The sequences of the primers used in all the genetic constructs will be made available upon request.

SDS-PAGE and immunoblotting. All SDS-PAGE was carried out using standard techniques. The Bradford (Bio-Rad) protein assay was used to determine sample protein concentration and normalize the amount of protein loaded in each lane. Once separated, the proteins were transferred to an Immobilon-FL PVDF membrane (Millipore) and then blocked in 5% skim milk in PBS, pH 7.4. Mouse monoclonal antibody against PdpA was added at a 1:2000 dilution, whereas mouse monoclonal antibodies against PdpB and IgIB were

used at a 1:4000 dilution. Rabbit polyclonal antibody against MglB was used at a 1:5000 dilution. Bound antibody was detected using IRDye800-conjugated goat anti-mouse or IRDye800-conjugated goat anti-rabbit antibody (Rockland Immunochemicals) and visualized, and the integrated intensity of fluorescence quantified, using the LiCor Odyssey imaging software version 2.1. The anti-PdpA monoclonal antibody was produced by immunizing mice with recombinant C-terminal portion (aa 405–783) of PdpA expressed from pET28a. The anti-PdpB and anti-IgIB antibodies were also produced by immunizing mice with recombinant protein expressed from pET28a. The mouse anti-PdpA hybridoma developed in this research, and the hybridomas producing anti-IgIB and anti-PdpB, have been deposited with the American Type Culture Collection's Biodefence and Emerging Infections Resources program.

Iron-limitation studies. *F. novicida* strains were grown overnight in TSBC at 37 °C without shaking. Then 3 ml of overnight culture was added to 30 ml fresh TSBC and allowed to grow for 1 h at 37 °C while shaking at 200 r.p.m. After 1 h the cultures were split into three flasks. One flask was left untreated and deferoxamine mesylate salt (Desferal) was added to the other two flasks to a final concentration of 1 mM. FeSO₄ (1 mM) was also added to one of the Desferal-containing flasks. Bacterial cultures were grown for a further 2 h. Cells were harvested by centrifugation at 1000 g for 15 min and resuspended in PBS, pH 7.4, containing protease inhibitors (Sigma). Cell suspensions were sonicated with five 30 s pulses and then prepared for SDS-PAGE and Western blotting as described above.

Subcellular fractionation of *F. novicida*. *F. novicida* U112 was added to J774A.1 macrophages at an m.o.i. of 1000:1 (bacterium to macrophage). Infected monolayers were incubated for 1 h at 37 °C, 5% CO₂ in complete Dulbecco's Modified Eagle Medium (cDMEM) supplemented with 10% (v/v) fetal bovine serum (FBS) and 4 mM L-glutamine. Macrophages were washed five times in sterile PBS (Invitrogen) and fresh growth medium was added. The infection was allowed to progress for 18 h, at which point the macrophages were lysed by adding deoxycholic acid to a final concentration of 0.1% (w/v). Bacteria were separated from the macrophage lysate by centrifugation at 2000 g for 15 min. The bacterial pellet was resuspended in PBS and sonicated to lyse cells. Any unbroken cells were removed by centrifugation at 10 000 g for 10 min. A sample was taken at this point as the total protein fraction. The cell lysate was ultracentrifuged at 100 000 g, 4 °C (Beckman ultramicrocentrifuge TLA-100.3) for 2 h to pellet the membranes. The supernatant representing the soluble proteins was removed and ultracentrifuged once more for 30 min at 100 000 g, 4 °C to minimize membrane protein contamination. The pellet representing the membrane-associated proteins was washed once in PBS and ultracentrifuged once more for 30 min at 100 000 g, 4 °C. The membrane-associated protein pellet was then treated with 1% (w/v) sodium lauroyl sarcosinate (Sarkosyl) and left at room temperature for 30 min. The inner membrane (Sarkosyl-soluble) and outer membrane (Sarkosyl-insoluble) fractions were separated by ultracentrifugation at 100 000 g at 4 °C for 2 h. The fractions were then processed for SDS-PAGE and immunoblotting as described above. To estimate the amount of inner-membrane protein associated with each fraction, the NADH oxidase activity was determined using the method described by Osborn *et al.* (1972). Fraction purity was determined by comparing relative enzyme activity per milligram of protein per fraction.

Chicken embryo infections. Fertilized White Leghorn chicken embryos were incubated for 7 days prior to infection. After incubation, embryos were infected with the appropriate strains of *F. novicida* in a range of doses as described by Nix *et al.* (2006). After infection, embryo death was monitored over a period of 6 days; the results of one inoculating dose are presented.

Mouse infections. For *in vivo* infections, 6- to 8-week-old male specific-pathogen-free BALB/cByJ mice were purchased from the Jackson Laboratory. Animals were housed in sterile micro-isolator cages in a barrier environment at the Center for Biologics Evaluation and Research. Mice were fed autoclaved food and water *ad libitum*, and all experiments were performed under Institutional Animal Care and Use Committee guidelines. Mice were given 0.1 ml of appropriately diluted bacteria intradermally at the base of the tail; actual doses of inoculated bacteria were simultaneously determined by plate count. All materials used in animals, including bacteria, were diluted in PBS (BioWhittaker) containing <0.01 ng endotoxin ml⁻¹.

Graphing and statistics. The Prism GraphPad v4.03 software was used to create graphs and determine statistical values. For comparison of different survival curves the log-rank test was used to generate *P*-values.

RESULTS

Detection of PdpA with monoclonal antibody

The *pdpA* gene is the first in an AT-rich, 12-cistron operon in the FPI (Fig. 1); it is 2463 bp in length, and encodes a predicted 820 aa (95 kDa) protein. The deduced protein shows greater than 97% amino acid identity across all sequenced strains of *Francisella*. The PdpA protein lacks a

sec-associated signal peptide, and does not show significant similarity to other proteins by standard BLASTP analysis. However, iterated BLAST analysis (PSI-BLAST) suggests the presence of a eukaryotic F-box motif, which is often found in ubiquitin-ligase complex-associated proteins. Analysis of PdpA using HHpredict software also detects several protein domains associated with the ubiquitin-proteasome pathway.

Although we previously used a genetic approach to demonstrate that PdpA plays a role in virulence, there has been no biochemical evidence demonstrating the expression of PdpA. To detect the product of the *pdpA* gene we developed a monoclonal antibody against a recombinant fragment of PdpA, and used it in Western immunoblots of *F. novicida* and *F. tularensis* extracts. As can be seen in Fig. 2(a), the PdpA band in *F. novicida* often runs as a distorted band of relative molecular mass of approximately 90. In the *F. tularensis* subsp. *holarctica* strain LVS or the *F. tularensis* subsp. *tularensis* strain B38 the PdpA band migrates as a tight band at a relative molecular mass of approximately 82. *F. novicida* has an O-antigen structure and banding pattern that is different from that found in *F. tularensis* strains and this may account for the distortion of the PdpA band that occurs in *F. novicida* but not in *F. tularensis* extracts.

To add confidence that the reactive band seen in immunoblots was truly PdpA, we created a deletion mutant that eliminated codons 1–813 of *pdpA* (Fig. 1b, c); the last seven codons were not deleted in order to preserve the ribosome-binding site of *pdpB*. The deletion mutation was verified by PCR analysis (Fig. 1b) and by sequencing of the genomic region of the Δ *pdpA* mutant. We also complemented the Δ *pdpA* mutation by inserting *pdpA* and the region encompassing 522 bp upstream of *pdpA* into an integrating vector, pJL-SKX, which inserts into the chromosome at approximately 507 kb from the *pdpA* locus. As mentioned, we reasoned that the distortion of the PdpA band was due to its co-migration with several LPS bands that are present in *F. novicida* but not in *F. tularensis* at the approximate relative molecular mass where PdpA is found. Hence, we also introduced a genetic lesion that diminishes the amount of O-antigen produced in *F. novicida* into the Δ *pdpA* mutant and the Δ *pdpA*/SKX::*pdpA* complemented strains. Immunoblot analysis of the resulting O-antigen-defective strains (Fig. 2b) showed that PdpA is not made in the Δ *pdpA* strain but is made in the wild-type and the complemented strain (both with O-antigen mutations). These results also show that the reduction in the amount of O-antigen results in PdpA migrating as a sharper band. In order to ascertain if the Δ *pdpA* lesion affected expression of genes downstream of *pdpA*, we assessed the expression levels of PdpB, which is encoded by the cistron immediately downstream of *pdpA*. As Fig. 2(b) shows, the level of PdpB was essentially identical in the wild-type, the Δ *pdpA* and the Δ *pdpA*/SKX::*pdpA* strains. We also assessed the expression of PdpA and PdpB in Δ *pdpA* and Δ *pdpA*/SKX::*pdpA* strains

Fig. 2. Detection and subcellular localization of PdpA in *Francisella*. (a) Reactivity of anti-PdpA monoclonal antibody with cell extracts of *F. novicida* and two *F. tularensis* subspecies. The O-antigen in *F. novicida* is thought to cause aberrant migration of PdpA. The migration of an 83 kDa molecular mass marker is shown on the right. (b) Effect of deletion of *pdpA* and mutation of O-antigen gene. An O-antigen mutation from the previously described mutant SC92 ('WT-LPS') was transferred to the Δ*pdpA* mutant and the complemented strain Δ*pdpA*/SKX::pdpA, and Western blots with anti-PdpA and anti-PdpB monoclonal antibodies are shown. For the top part of (b) the number at the right indicates actual migration of a molecular mass marker, and for the bottom part, the number indicates a calculated relative molecular mass. (c) A Western blot using anti-PdpB antibody shows that the *pdpA*::Em^R gene replacement mutant NZ9 expresses greatly reduced levels of PdpB compared to wild-type (WT). (d) Bacteria isolated from a J774A.1 macrophage infection were fractionated to separate soluble and membrane-associated proteins. PdpA was found to localize with the soluble proteins, as was the transcriptional regulator MglB. PdpB localized to the Sarkosyl-soluble protein fraction, indicating its association with the inner membrane of *F. novicida*. NADH oxidase activity was determined in each fraction as a measure of the relative amount of inner-membrane protein. The number to the right of the top panel of (d) represents the actual migration of the molecular mass marker, whereas the numbers in the two bottom panels represent calculated relative molecular masses. The amount of sample loaded in each lane was normalized by protein content. Results are representative of duplicate experiments.

that had the wild-type form of the O-antigen and found essentially identical results as in Fig. 2(b) except that the PdpA band was usually distorted (data not shown). Unlike the markerless *pdpA* mutant constructed in this study, the *pdpA*::Em^R allelic replacement mutant, NZ9, demonstrated marked polarity effects on PdpB (Fig. 2c).

Solubility of PdpA

Some of the genes in the FPI encode homologues of components of T6SSs, and have been shown to be membrane associated. To test the possible role of PdpA as a membrane-bound component of a possible T6SS, we analysed the localization of PdpA in *F. novicida* cells that had infected macrophages. We reasoned that the localization of PdpA during an infection would provide information that might suggest a function. The data presented in Fig. 2(d) show that PdpA localizes to the soluble bacterial fraction, and co-localizes with the gene regulatory protein MglB. We had previously shown that PdpB localizes to the inner membrane in broth-grown cells (Ludu *et al.*, 2008a), and it localized to the same fraction in macrophage-grown cells (Fig. 2d). Outer-membrane pro-

teins Tul4 and FopA were only detected in the Sarkosyl-insoluble fraction using anti-*F. novicida* antibody (data not shown). Similar results were obtained when we analysed the localization of PdpA in broth-grown *F. novicida* (data not shown), with the majority of PdpA localizing to the soluble fraction.

We previously showed that inactivation of FPI genes *iglA*, *dotU* and *pdpB*, which are homologues of canonical components of T6SSs, abolished transport of IglC to the outer membrane (Ludu *et al.*, 2008a). We found that IglC transport in the Δ*pdpA* background was unaffected (data not shown), suggesting that PdpA does not play a role in T6SS-mediated secretion. However, these results should be regarded cautiously, since we have so little information about possible T6SS-mediated secretion in general, and especially in *Francisella*. Attempts to visualize localization of PdpA in the macrophage cytosol were unproductive (data not shown).

PdpA levels affected by *mglAB* background and iron concentration

Previous work has shown that FPI mRNA expression is depressed when MglA is absent and is increased by low iron concentration (Brotcke *et al.*, 2006; Charity *et al.*, 2007; Deng *et al.*, 2006; Lauriano *et al.*, 2003), but none of these works examined protein expression levels. In order to

assess the effect of these factors on PdpA protein expression, we examined protein production by immunoblot analysis. The expression of PdpA was much reduced in both *mglA* (GB2) and *mglB* (GB6) genetic backgrounds as compared to expression in the wild-type strain of *F. novicida* (Fig. 3a). A similar effect was seen on the expression of PdpB and IgIB in the *mglA* and *mglB* backgrounds (Fig. 3a). Addition of the chelator Desferal to cultures of *F. novicida* resulted in an increase in PdpA expression in *F. novicida* (Fig. 3b), presumably due to the reduction in the amount of available iron. IgIB, IgIC and PdpB expression responded to the addition of Desferal essentially the same as did PdpA (Fig. 3b). The effect of the chelator on protein expression was reversed by the addition of 1 mM FeSO₄, which resulted in PdpA, PdpB, IgIB and IgIC levels that were below those found in untreated cultures (Fig. 3b). Expression levels of a cross-reactive

control protein remained constant regardless of the addition of chelator or excess iron. The levels of PdpB, IgIB and IgIC in the $\Delta pdpA$ strain responded essentially identically to chelator or iron conditions as in the wild-type *F. novicida* strain (data not shown).

Virulence properties of the $\Delta pdpA$ mutant

Our previous work and the studies of others suggest that PdpA plays a role in virulence. However, these previous studies were unable to measure expression of proteins encoded by genes downstream of *pdpA* or failed to carry out genetic complementation; most studies failed to carry out either of these controls. Hence, we felt it was important to test the virulence of a $\Delta pdpA$ mutant that lacked translational polarity and could be complemented with a single copy of *pdpA*. We first performed a series of infections with different inoculating doses in chicken embryos and followed these with smaller experimental infections of mice. The data presented in Fig. 4 show that, at the inoculating dose of 10³ c.f.u., wild-type *F. novicida* killed 100% of the chicken embryos by day 4 post-infection. In contrast, the $\Delta pdpA$ strain failed to kill 50% of the embryos by 6 days after the start of the infection. Complementation of the $\Delta pdpA$ mutation with a single copy of wild-type *pdpA* resulted in a strain, $\Delta pdpA$ /SKX::*pdpA*, that was equally virulent as the wild-type strain in the chicken embryo infection assay. Similar results were found in mouse infection experiments. Infection with 10⁵ wild-type *F. novicida* killed 5/5 BALB/cByJ male mice but infection of 10 mice with 10⁷ $\Delta pdpA$ failed to kill any mice. As we have shown before (Ludu *et al.*, 2008a), infections of chicken embryos with *F. novicida* mutants are largely predictive of the outcome of infection of mice with the same mutants. However, mice appear to be considerably more resistant to killing, especially by attenuated *F. novicida*.

DISCUSSION

Recent study of *F. tularensis* has yielded surprisingly little advancement concerning the characterization and function of virulence-associated proteins. Although many virulence factors have been discovered, particularly within the FPI, several of these proteins have proven to be very difficult to study due to toxicity issues and low expression levels. This research represents the first examples of PdpA protein characterization and has investigated the virulence of a non-polar, markerless, in-frame, *pdpA* deletion mutant.

Immediately downstream of *pdpA* lies the *pdpB* gene, and the *pdpB* ribosome-binding site overlaps the 3' end of *pdpA*. This tight translational coupling of *pdpA* and *pdpB*, and between most other adjacent FPI genes, is consistent with mRNA expression studies that indicate that FPI genes are controlled by the same global regulators, including MglA, SspA, PmrA and FevR, or by the same environmental cues, such as low iron concentration (Brotcke

Fig. 3. Regulation of PdpA expression by iron and transcriptional regulators. (a) Levels of PdpA, PdpB and IgIB protein expression were determined in *mglA* and *mglB* mutant backgrounds. (b) Expression of these proteins, as well as IgIC, was also determined in varying iron conditions. The relative level of fluorescence signal generated by reactivity of anti-PdpA, anti-PdpB, anti-IgIB, or anti-IgIC monoclonal antibodies, calculated as described in Methods, is indicated. The amount of sample loaded to the wells of each polyacrylamide gel was normalized by protein content. The reactivity of a non-specific, cross-reactive protein band is shown in the bottom panel. Results are representative of experiments performed in triplicate.

Fig. 4. Attenuation of the $\Delta pdpA$ mutant in chicken embryos. Seven-day-old chicken embryos were infected with 10^3 c.f.u. of wild-type (WT) (a), $\Delta pdpA$ (b) and $\Delta pdpA/SKX::pdpA$ (c) strains of *F. novicida*. Time to death of the embryos was monitored over a period of 6 days. All experiments were done at three separate inoculating doses, and the experiments included at least seven embryos per infective dose per strain; the survival curves presented are representative of three separate trials. The statistical difference between the WT and $\Delta pdpA$ strain survival curve was measured by the log-rank test and yielded a *P*-value of <0.001 . The *P*-value for a comparison of $\Delta pdpA$ vs $\Delta pdpA/SKX::pdpA$ was <0.001 .

et al., 2006; Brotcke & Monack, 2008; Charity *et al.*, 2007; Deng *et al.*, 2006; Lenco *et al.*, 2007; Mohapatra *et al.*, 2007). Data presented in this work on the effect of MglA and iron concentration on expression of PdpA and other FPI-encoded proteins provide evidence that expression extends to the protein level. The MglB protein has been presumed to be involved in expression, and in this work we provided evidence supporting this idea, as PdpA, PdpB and IglB expression was virtually undetectable in an *mglB* mutant background.

It is also apparent that the allelic replacement of *pdpA* has polar effects on expression of the downstream gene *pdpB*,

as there is considerably less PdpB protein present in strain NZ9 ($\Delta pdpA::Em^R$). This polarity may account for the increased attenuation of NZ9 in chicken embryos compared to $\Delta pdpA$ (data not shown).

One central biological question regarding the FPI-encoded proteins involves their possible role in secretion or as effector proteins that interact with host-cell components. Using macrophage-grown *F. novicida* we found PdpA in the bacterial cytoplasm. Although macrophage-grown *F. novicida* cells are more difficult to analyse than broth-grown cells, we reasoned that they would provide better evidence of the biologically relevant localization of PdpA than analysis of broth-grown *F. novicida*. Localization of PdpA to the bacterial cytoplasm may lead one to believe that the protein is not a component of the T6SS, as most proteins involved in assembling a secretion apparatus are membrane associated. However, this is not always the case; T6SS proteins have been shown to localize to both membrane and cytoplasmic fractions (Wu *et al.*, 2008) so our results have to be interpreted carefully. As a soluble protein, PdpA may function as a chaperone component of the T6SS or may be a secreted effector protein. We believe that PdpA might be a secreted effector protein as it does not appear to play a role in secretion as defined by our recently described criteria (Ludu *et al.*, 2008a), yet its absence profoundly decreases virulence in both chicken embryos and adult mice. Also, the deduced PdpA amino acid sequence shows low-level similarities to fragments of multiple proteins involved in the eukaryotic ubiquitin-proteasome pathway, and such molecular mimicry has been observed in virulence effectors of other intracellular pathogens (Angot *et al.*, 2007). One of the motifs that we identified is a possible F-box motif in the N-terminal region of PdpA. In eukaryotic cells F-box-containing proteins are involved in protein-protein interactions, most commonly in proteins in the ubiquitin-ligase complex (Kipreos & Pagano, 2000). Inspection of the Pfam tree of F-box proteins (PF00646; <http://pfam.sanger.ac.uk/family?acc=PF00646>) reveals several prokaryotic F-box-containing proteins, and F-box regions have been shown to be important in two bacterial plant pathogens (Angot *et al.*, 2006; Tzfira *et al.*, 2004). Attempts to express domains that showed similarity to those found in ubiquitin-ligase complexes have been unsuccessful due to apparent toxicity involved in expressing PdpA protein domains, particularly those located within the amino terminus of the protein. Also, the introduction of alanine substitutions into key residues of the putative F-box failed to affect virulence of the resultant *F. novicida* mutants (data not shown). All efforts to detect PdpA in the host-cell cytoplasm have been unsuccessful. Although we believe PdpA is a secreted effector protein, the current tools and methodologies available have been unable to confirm this. Clearly, extensive analysis of PdpA will be required to surmise its function in virulence. The accompanying paper (Schmerck *et al.*, 2009) reports studies on the intracellular growth phenotype of the $\Delta pdpA$ mutant and the effect of the

mutation on the macrophage gene expression response to *F. novicida* infection.

ACKNOWLEDGEMENTS

This work was supported by grant 5R01 AI056212-02 from the National Institute of Allergy and Infectious Diseases and MOP 89812 from the Canadian Institutes of Health Research to F.E.N. The development of monoclonal antibodies was partially supported by a contract from the Biodefense and Emerging Infections Resources program at the American Type Culture Collection.

REFERENCES

- Angot, A., Peeters, N., Lechner, E., Vaillau, F., Baud, C., Gentzmittel, L., Sartorel, E., Genschik, P., Boucher, C. & Genin, S. (2006). *Ralstonia solanacearum* requires F-box-like domain-containing type III effectors to promote disease on several host plants. *Proc Natl Acad Sci U S A* **103**, 14620–14625.
- Angot, A., Vergunst, A., Genin, S. & Peeters, N. (2007). Exploitation of eukaryotic ubiquitin signaling pathways by effectors translocated by bacterial type III and type IV secretion systems. *PLoS Pathog* **3**, e3.
- Anthony, L. D., Burke, R. D. & Nano, F. E. (1991). Growth of *Francisella* spp. in rodent macrophages. *Infect Immun* **59**, 3291–3296.
- Baron, G. S. & Nano, F. E. (1998). MglA and MglB are required for the intramacrophage growth of *Francisella novicida*. *Mol Microbiol* **29**, 247–259.
- Brotcke, A. & Monack, D. M. (2008). Identification of *fevR*, a novel regulator of virulence gene expression in *Francisella*. *Infect Immun* **76**, 3473–3480.
- Brotcke, A., Weiss, D. S., Kim, C. C., Chain, P., Malfatti, S., Garcia, E. & Monack, D. M. (2006). Identification of MglA-regulated genes reveals novel virulence factors in *Francisella tularensis*. *Infect Immun* **74**, 6642–6655.
- Charity, J. C., Costante-Hamm, M. M., Balon, E. L., Boyd, D. H., Rubin, E. J. & Dove, S. L. (2007). Twin RNA polymerase-associated proteins control virulence gene expression in *Francisella tularensis*. *PLoS Pathog* **3**, e84.
- Clemens, D. L., Lee, B. Y. & Horwitz, M. A. (2004). Virulent and avirulent strains of *Francisella tularensis* prevent acidification and maturation of their phagosomes and escape into the cytoplasm in human macrophages. *Infect Immun* **72**, 3204–3217.
- Cowley, S. C., Gray, C. J. & Nano, F. E. (2000). Isolation and characterization of *Francisella novicida* mutants defective in lipopolysaccharide biosynthesis. *FEMS Microbiol Lett* **182**, 63–67.
- de Bruin, O. M., Ludu, J. S. & Nano, F. E. (2007). The *Francisella* pathogenicity island protein IglA localizes to the bacterial cytoplasm and is needed for intracellular growth. *BMC Microbiol* **7**, 1.
- Deng, K., Blick, R. J., Liu, W. & Hansen, E. J. (2006). Identification of *Francisella tularensis* genes affected by iron limitation. *Infect Immun* **74**, 4224–4236.
- Fortier, A. H., Green, S. J., Polsinelli, T., Jones, T. R., Crawford, R. M., Leiby, D. A., Elkins, K. L., Meltzer, M. S. & Nacy, C. A. (1994). Life and death of an intracellular pathogen: *Francisella tularensis* and the macrophage. *Immunol Ser* **60**, 349–361.
- Golovliov, I., Baranov, V., Krocova, Z., Kovarova, H. & Sjostedt, A. (2003a). An attenuated strain of the facultative intracellular bacterium *Francisella tularensis* can escape the phagosome of monocytic cells. *Infect Immun* **71**, 5940–5950.
- Golovliov, I., Sjostedt, A., Mokrievich, A. & Pavlov, V. (2003b). A method for allelic replacement in *Francisella tularensis*. *FEMS Microbiol Lett* **222**, 273–280.
- Gray, C. G., Cowley, S. C., Cheung, K. K. & Nano, F. E. (2002). The identification of five genetic loci of *Francisella novicida* associated with intracellular growth. *FEMS Microbiol Lett* **215**, 53–56.
- Kipreos, E. T. & Pagano, M. (2000). The F-box protein family. *Genome Biol* **1**, REVIEWS3002.
- Lai, X. H., Golovliov, I. & Sjostedt, A. (2004). Expression of IglC is necessary for intracellular growth and induction of apoptosis in murine macrophages by *Francisella tularensis*. *Microb Pathog* **37**, 225–230.
- Lauriano, C. M., Barker, J. R., Nano, F. E., Arulanandam, B. P. & Klose, K. E. (2003). Allelic exchange in *Francisella tularensis* using PCR products. *FEMS Microbiol Lett* **229**, 195–202.
- Lauriano, C. M., Barker, J. R., Yoon, S.-S., Nano, F. E., Arulanandam, B. P., Hassett, D. J. & Klose, K. E. (2004). MglA regulates transcription of virulence factors necessary for *Francisella tularensis* intraamoebae and intramacrophage survival. *Proc Natl Acad Sci U S A* **101**, 4246–4249.
- Lenco, J., Hubalek, M., Larsson, P., Fucikova, A., Brychta, M., Macela, A. & Stulik, J. (2007). Proteomics analysis of the *Francisella tularensis* LVS response to iron restriction: induction of the *F. tularensis* pathogenicity island proteins IglABC. *FEMS Microbiol Lett* **269**, 11–21.
- Ludu, J. S., de Bruin, O. M., Duplantis, B. N., Schmerk, C. L., Chou, A. Y., Elkins, K. L. & Nano, F. E. (2008a). The *Francisella* pathogenicity island protein PdpD is required for full virulence and associates with homologues of the type VI secretion system. *J Bacteriol* **190**, 4584–4595.
- Ludu, J. S., Nix, E. B., Duplantis, B. N., de Bruin, O. M., Gallagher, L. A., Hawley, L. M. & Nano, F. E. (2008b). Genetic elements for selection, deletion mutagenesis and complementation in *Francisella* spp. *FEMS Microbiol Lett* **278**, 86–93.
- Mohapatra, N. P., Soni, S., Bell, B. L., Warren, R., Ernst, R. K., Muszynski, A., Carlson, R. W. & Gunn, J. S. (2007). Identification of an orphan response regulator required for the virulence of *Francisella* spp. and transcription of pathogenicity island genes. *Infect Immun* **75**, 3305–3314.
- Nano, F. E. & Schmerk, C. (2007). The *Francisella* pathogenicity island. *Ann N Y Acad Sci* **1105**, 122–137.
- Nano, F. E., Zhang, N., Cowley, S. C., Klose, K. E., Cheung, K. K., Roberts, M. J., Ludu, J. S., Letendre, G. W., Meierovics, A. I. & other authors (2004). A *Francisella tularensis* pathogenicity island required for intramacrophage growth. *J Bacteriol* **186**, 6430–6436.
- Nix, E. B., Cheung, K. K., Wang, D., Zhang, N., Burke, R. D. & Nano, F. E. (2006). Virulence of *Francisella* spp. in chicken embryos. *Infect Immun* **74**, 4809–4816.
- Osborn, M. J., Gander, J. E. & Parisi, E. (1972). Mechanism of assembly of the outer membrane of *Salmonella typhimurium*. Site of synthesis of lipopolysaccharide. *J Biol Chem* **247**, 3973–3986.
- Proctor, R. A., White, J. D., Ayala, E. & Canonico, P. G. (1975). Phagocytosis of *Francisella tularensis* by Rhesus monkey peripheral leukocytes. *Infect Immun* **11**, 146–151.
- Santic, M., Molmeret, M., Klose, K. E., Jones, S. & Kwaik, Y. A. (2005). The *Francisella tularensis* pathogenicity island protein IglC and its regulator MglA are essential for modulating phagosome biogenesis and subsequent bacterial escape into the cytoplasm. *Cell Microbiol* **7**, 969–979.
- Schmerk, C. L., Duplantis, B. N., Howard, P. L. & Nano, F. E. (2009). A *Francisella novicida* *pdpA* mutant exhibits limited intracellular replication and remains associated with the lysosomal marker LAMP-1. *Microbiology* **155**, 1498–1504.

- Tempel, R., Lai, X. H., Crosa, L., Kozlowicz, B. & Heffron, F. (2006).** Attenuated *Francisella novicida* transposon mutants protect mice against wild-type challenge. *Infect Immun* **74**, 5095–5105.
- Tzfira, T., Vaidya, M. & Citovsky, V. (2004).** Involvement of targeted proteolysis in plant genetic transformation by *Agrobacterium*. *Nature* **431**, 87–92.
- Wang, R. F. & Kushner, S. R. (1991).** Construction of versatile low-copy-number vectors for cloning, sequencing and gene expression in *Escherichia coli*. *Gene* **100**, 195–199.
- Weiss, D. S., Brotcke, A., Henry, T., Margolis, J. J., Chan, K. & Monack, D. M. (2007).** In vivo negative selection screen identifies genes required for *Francisella* virulence. *Proc Natl Acad Sci U S A* **104**, 6037–6042.
- Wu, H. Y., Chung, P. C., Shih, H. W., Wen, S. R. & Lai, E. M. (2008).** Secretome analysis uncovers an Hcp-family protein secreted via a type VI secretion system in *Agrobacterium tumefaciens*. *J Bacteriol* **190**, 2841–2850.

Edited by: T. P. Hatch