Protective capacities of cell surface-associated proteins of *Streptococcus suis* mutants deficient in divalent cation-uptake regulators

Jesús Aranda,¹ Maria Elena Garrido,^{1,2} Nahuel Fittipaldi,³ Pilar Cortés,^{1,4} Montserrat Llagostera,^{1,2} Marcelo Gottschalk³ and Jordi Barbé^{1,2}

Correspondence
Jordi Barbé
jordi.barbe@uab.cat

¹Department de Genètica i Microbiologia, Universitat Autònoma de Barcelona (UAB), Bellaterra, 08193 – Barcelona, Spain

²Centre de Recerca en Sanitat Animal (CReSA), Bellaterra, 08193 – Barcelona, Spain

³Groupe de Recherche sur les Maladies Infectieuses du Porc and Centre de Recherche en Infectiologie Porcine, Faculté de Médecine Vétérinaire, Université de Montréal, St-Hyacinthe, QC J2S 7C6, Canada

⁴Servei d'Anàlisi i d'Aplicacions Microbiològiques, UAB, Bellaterra, 08193 – Barcelona, Spain

Many cell surface-associated, divalent cation-regulated proteins are immunogenic, and some of them confer protection against the bacterial species from which they are derived. In this work, two *Streptococcus suis* divalent cation uptake regulator genes controlling zinc/manganese and iron uptake (*adcR* and *fur*, respectively) were inactivated in order to study the protective capacities of their cell surface-associated proteins. The results obtained showed overexpression of a set of immunogenic proteins (including members of the pneumococcal histidine triad family previously reported to confer protection against streptococcal pathogens) in *S. suis* *adcR* mutant cell surface extracts. Likewise, genes encoding zinc transporters, putative virulence factors and a ribosomal protein paralogue related to zinc starvation appeared to be derepressed in this mutant strain. Moreover, protection assays in mice showed that although neither *adcR*- nor *fur*-regulated cell surface-associated proteins were sufficient to confer protection in mice, the combination of both *adcR*- and *fur*-regulated cell surface-associated proteins is able to confer significant protection (50 %, $P=0.038$) against a challenge to mice vaccinated with them.

Received 28 November 2008
Revised 22 January 2009
Accepted 23 January 2009

INTRODUCTION

Streptococcus suis is an important pathogen that causes significant economic losses in the swine industry worldwide (Higgins *et al.*, 1995). Infection caused by *S. suis* may vary from asymptomatic bacteraemia to fulminant systemic disease, with meningitis as the most outstanding feature (Higgins *et al.*, 1995). It is also an important zoonotic agent for humans, who may acquire the infection through skin wounds upon contact with pigs and/or their by-products. Recently, during an outbreak in China, 215 human cases of *S. suis* infection were documented, and 39 of those patients died (Yu *et al.*, 2006). More than 30 serotypes of *S. suis*, based on its capsular antigens, are currently known (Higgins *et al.*, 1995). Serotype 2 is the most frequently isolated from diseased animals (Staats *et al.*, 1997). Efforts to control

infection are hampered by the lack of effective vaccines against *S. suis*. Furthermore, mechanisms involved in pathogenesis of this micro-organism are not yet completely understood (Gottschalk & Segura, 2000).

Iron and zinc are essential components of many bacterial proteins either as structural or catalytic cofactors (Andrews *et al.*, 2003; Babich & Stotzky, 1978). The concentration of both elements in the free state in the body fluids of mammals is extremely low in order to prevent bacterial proliferation. On the other hand, metals at high concentrations are toxic to micro-organisms. Toxicity occurs through the displacement of essential metals from their native binding sites or through ligand interactions (Babich & Stotzky, 1978; Chamnongpol *et al.*, 2002). In addition, iron can be particularly toxic under aerobic conditions, interacting with either oxygen or oxygen-reduced species (Touati, 2000). Consequently, micro-organisms require homeostatic mechanisms to control intracellular metal levels.

Fur (ferric uptake regulator) is a transcriptional regulator of genes involved in iron uptake and storage in many

Abbreviations: ABC, ATP-binding cassette; EMSA, electrophoretic mobility shift assay.

Two supplementary tables, listing bacterial strains and plasmids and oligonucleotide primers used in this work, are available with the online version of this paper.

bacterial species (Andrews *et al.*, 2003). In *Escherichia coli* and *Bacillus subtilis*, and in species closely related to them, transcription of the zinc transporter genes is regulated by Zur (zinc uptake regulator), which belongs to the Fur protein family (Patzner & Hantke, 2000). However, in *Streptococcus* species, zinc transporter genes are controlled by AdcR, a transcriptional regulatory factor of the MarR protein family (Loo *et al.*, 2003).

Several cell surface-associated transporters involved in divalent cation uptake, including ATP-binding cassette (ABC) transporters, have been shown to elicit a protective immune response (Garmory & Titball, 2004; Tanabe *et al.*, 2006). It has been shown that in *S. suis*, AdcR controls at least two operons related to Zn^{2+}/Mn^{2+} uptake, encoding a complete ABC transporter (SsuiDRAFT 1237–1240) and a divalent cation-binding lipoprotein (SsuiDRAFT 0103) (Aranda *et al.*, 2008). Interestingly, both cation-binding lipoproteins encoded by these operons (SsuiDRAFT 1237 and SsuiDRAFT 0103) are immunogenic when analysed on Western blots using immunologically reactive serum obtained from mice surviving after inoculation with the wild-type *S. suis* 89/1591 strain (Aranda *et al.*, 2008). Furthermore, a high level of protection against *S. suis* challenge is obtained in mice when vaccinated with the product of the ssuiDRAFT 0103 ORF (Aranda *et al.*, 2008).

Different strategies have been developed to produce vaccines against pathogenic bacteria, including overexpression on the bacterial cell surface of divalent cation-uptake transporters which are induced in the presence of chelating agents, or by the construction of strains deficient in cation-uptake repressors (Garrido *et al.*, 2008). In this context, the aim of this work has been to obtain knockout *S. suis* mutants for the regulator genes of either zinc/manganese- or iron-uptake systems to study their possible use as a protective tool against this pathogen.

METHODS

Bacterial strains, plasmids and growth conditions. Bacterial strains and plasmids used in this work are listed in Supplementary Table S1. *E. coli* cells were grown in Luria–Bertani (LB) medium (Miller, 1992). When necessary, ampicillin ($50 \mu\text{g ml}^{-1}$), kanamycin ($50 \mu\text{g ml}^{-1}$) or spectinomycin ($50 \mu\text{g ml}^{-1}$) was added to LB medium. *S. suis* was grown in Todd–Hewitt medium (TH medium; Difco). When required, spectinomycin ($100 \mu\text{g ml}^{-1}$) was added to TH medium. Unless otherwise indicated, all cultures were incubated at 37°C , with shaking for *E. coli*. DNA extractions, cloning, and other molecular techniques used were performed as described elsewhere (Sambrook & Russell, 2001). All oligonucleotides (Roche) used in this work are listed in Supplementary Table S2.

Construction of *S. suis* mutants. The *adcR* and *fur* genes were inactivated by precise in-frame deletion. Briefly, upstream and downstream regions of both *adcR* and *fur* genes (including partial coding regions) were amplified by overlap-extension PCR using genomic DNA from *S. suis* P1/7 strain as a template. A PCR product of 892 bp with a deletion of 229 bp was obtained for the *adcR* gene, and a PCR product of 848 bp with a deletion of 222 bp was obtained for the *fur* gene. Both PCR fragments were cloned into the pCR4-

TOPO vector following the manufacturer's instructions and propagated in *E. coli* One Shot TOP 10 cells. Inserts were amplified afterwards by PCR using primers M13F and M13R (Supplementary Table S2), amplicons were digested with *EcoRI* and cloned into the *EcoRI*-digested thermosensitive suicide vector pSET4s, which carries the *spc* and *repA*s genes conferring spectinomycin resistance and thermosensitivity, respectively (Takamatsu *et al.*, 2001b), giving rise to plasmids pSET4 Δ adcR and pSET4 Δ fur. Recombinant plasmids were propagated in *E. coli* ABLE-C cells, purified, and used to electroporate *S. suis* electrocompetent P1/7 and UA5000 strains, as described elsewhere (Takamatsu *et al.*, 2001a). Cells were then plated on TH agar containing spectinomycin and incubated for 48 h at 28°C to select for clones that had incorporated the plasmid. Several clones were then selected and purified by single-colony isolation on TH agar containing spectinomycin and incubated at 37°C . Integration of the construct into the chromosome as a result of homologous recombination via a single crossover was verified by PCR in temperature- and spectinomycin-resistant colonies. These colonies were then passaged several times onto non-selective media at 28°C and screened for loss of vector-mediated spectinomycin resistance. Candidate spectinomycin-sensitive, temperature-resistant clones were analysed by PCR, which showed the exchange of the wild-type allele for the genetic segment containing the desired deletion as a consequence of a homologous recombination through a double-crossover event. Allelic replacement in candidate clones was further confirmed by sequencing of the region in the resulting mutants.

Preparation of *S. suis* cell surface-associated proteins.

Extraction of these proteins was performed as described elsewhere (Tavares & Sellstedt, 2000), with some modifications. Briefly, cells from 200 ml overnight culture were pelleted at 4°C by centrifugation at 8000 g for 10 min. The cells were washed with 62.5 mM Tris/HCl (pH 6.8) and the pellet was pipetted up and down for 5 min in 4 ml of the same buffer supplemented with 0.1% (v/v) Triton X-100 (Roche). The suspension was centrifuged at 8000 g for 10 min at 4°C . The supernatant was collected and centrifuged for 2 h at 51 030 g at 4°C in order to concentrate the cell surface-associated protein extract. Finally, the protein pellet was resuspended in 100 μl 62.5 mM Tris/HCl (pH 6.8), visualized by SDS-PAGE (Laemmli, 1970) and stored at 4°C . The SDS-PAGE gel was stained with Brilliant Blue G Colloidal (Sigma) in order to avoid any stain interference with the subsequent procedures. The identification of overexpressed proteins was done by the Servei de Proteòmica i Bioinformàtica (SepBio) of the Universitat Autònoma de Barcelona (UAB). Thus, electrophoretic bands were in-gel digested with trypsin and analysed by MALDI-TOF MS as previously reported (Abella *et al.*, 2007).

RNA techniques. RNA was isolated as described elsewhere (Fontaine *et al.*, 2004), with slight modifications. A 10 ml sample of an *S. suis* culture in the mid-exponential growth phase (OD_{600} 0.6) was collected by centrifugation at 8000 g for 10 min. Cells were then resuspended by vigorous vortexing at 37°C for 10 min in 300 μl Tris-EDTA prelysis buffer containing 10 mg lysozyme ml^{-1} . Total RNA was extracted by using the RNeasy Mini kit (Qiagen) according to the manufacturer's instructions. DNA contamination was removed from the RNA during purification by treatment with RNase-free DNase (Qiagen), followed by digestion with DNase Turbo (Ambion). The concentration and integrity of RNA were determined by A_{260} measurements and 1% agarose gel electrophoresis, respectively. RT-PCR assays were done using a Titan One Tube RT-PCR system (Roche) following the manufacturer's instructions. Real-time RT-PCR analysis of gene expression was performed for all genes as reported previously (Campoy *et al.*, 2002), with specific internal oligonucleotides used for each gene (Supplementary Table S2). In all cases, the absence of DNA in RNA samples was tested by PCR without the addition of reverse transcriptase.

Mobility shift assays. Electrophoretic mobility shift assays (EMSAs) were done as described elsewhere (Bsat & Helmann, 1999) with slight modifications. DNA promoters were PCR-amplified from P1/7 *S. suis* genomic DNA using suitable oligonucleotides (Supplementary Table S2), and the purified PCR fragments were cloned in *E. coli* DH5 α cells via the pGEM-T vector. The presence of the desired promoter was confirmed by sequencing the plasmid DNA using the primers M13F and M13R. DNA probes were prepared by PCR amplification, using primers of which one was labelled with DIG at its 5' end (Supplementary Table S2). DNA–protein reaction mixtures (20 μ l) containing 25 ng DIG-DNA-labelled probe and either 0 or 10 μ g purified *S. suis* AdcR protein were incubated in EMSA buffer [20 mM Tris/HCl (pH 8), 50 mM KCl, 5% (v/v) glycerol, 1 μ g bulk carrier salmon sperm DNA, 0.5 mM DTT, and 0.1 mg BSA ml⁻¹] for 10 min at room temperature. DNA–protein complexes were visualized by separation on a 5% non-denaturing polyacrylamide gel (40 mM Tris/acetate) at 150 V for 1.5 h and then transferred to a Biodine B nylon membrane (Pall Gelman Laboratory). DIG-DNA-labelled protein complexes were detected according to the manufacturer's protocol (Roche). AdcR protein was purified as previously described (Aranda *et al.*, 2008).

Protection assays. Female BALB/cAnNHsd mice (3 weeks old) obtained from Harlan Iberica were used for these studies. All animals underwent a quarantine period of 1 week. A maximum of four animals per cage were housed under specific-pathogen-free conditions at 19–21 °C with an artificial 12 h light/12 h dark cycle; the relative humidity was 50–60%. Antibiotic-free pelleted food and autoclaved water were provided *ad libitum* during the experiments. Animals were monitored daily for morbidity and mortality. All animal experiments were approved by the UAB Animal Ethics Committee. For protection assays, groups of eight female 4-week-old BALB/cAnNHsd mice were given intraperitoneal injections (0.1 ml) of 50 μ g cell surface-associated proteins adsorbed with Imject Alum adjuvant (aluminium hydroxide; Pierce) according to the manufacturer's protocol. One additional group of eight mice was given injections of 0.1 ml solvent including Imject Alum adjuvant as a negative control. After 2 weeks, a second immunization was carried out, followed 3 weeks later by a challenge with an intraperitoneal injection of a 0.1 ml suspension of TH broth supplemented with 10% inactivated bovine serum containing 10⁸ c.f.u. *S. suis* 89/1591. The survival of the mice was subsequently monitored for 7 days. Serum for Western blot assays was obtained as previously reported (Aranda *et al.*, 2008).

Statistical analysis. Fisher's exact test was used to analyse the statistical significance of the lethal challenge data. A statistical test with a *P* value less than 0.05 was considered to be significant.

RESULTS AND DISCUSSION

Analysis of the *S. suis* *adcR* regulon composition

It has been predicted *in silico* and demonstrated *in vitro* that the TTAACNRGTTAA motif is the AdcR DNA-binding sequence (Aranda *et al.*, 2008; Panina *et al.*, 2003). A search in the complete sequenced genome of *S. suis* strain P1/7 (http://www.sanger.ac.uk/Projects/S_suis/) with the EditSeq program (DNASTAR) revealed four ORFs with a motif identical to the AdcR-binding sequence in their promoter regions.

Two of the ORFs are *ssu1390* and *ssu1352*, whose predicted products are homologous, respectively, to internalin A (InlA) of *Listeria monocytogenes* [which allows this bacterium to invade cells (Wollert *et al.*, 2007)] and to the 30S ribosomal protein S14, which lacks the typical zinc-ribbon motif present in most ribosomal proteins. Furthermore, several EMSAs confirmed that the purified AdcR protein binds specifically to the promoters of both genes (Fig. 1).

The other two ORFs of strain P1/7 presenting an AdcR-binding sequence correspond to the *S. suis* 89/1591 homologues previously identified as *ssu1240* ORF (*adcR*) and *ssu1013* ORF (encoding a putative zinc-binding streptococcal lipoprotein) (Aranda *et al.*, 2008). These ORFs were annotated as *ssu0112* and *ssu0308*, respectively, in the *S. suis* P1/7 strain (Table 1). As in strain 89/1591, in P1/7 *adcR* is the first gene of a transcriptional unit, which also contains three components of an ABC transporter (*Ssu0113–0115*) involved in zinc/manganese uptake (Table 1). Furthermore, it is worth noting that both *adcR*-regulated *ssu1240* and *ssu1013* lipoproteins (*Ssu0308* and *Ssu0115* P1/7 homologues) have been reported to be immunogenic and able to confer protection against *S. suis* in mice (Aranda *et al.*, 2008). In this regard, it is known that lipoproteins are the ABC transporter components most exposed outside the Gram-positive cell surface (Hantke, 2001; Sutcliffe & Russell, 1995). Taken together, these data suggest that *adcR* mutant cell surface-associated proteins could be candidates for effective vaccines against *S. suis* infection.

Fig. 1. (a) Schematic representation of *ssu1352* and *ssu1390* ORFs with their promoter regions (P1352 and P1390, respectively). Small arrows indicate the primer pairs used to amplify the promoters for EMSA experiments. (b) Electrophoretic mobility of the DNA fragments containing the indicated *S. suis* promoters in the absence (–) or presence (+) of purified *S. suis* AdcR protein.

Table 1. Nomenclature of *adcR*-regulated genes in P1/7 and 89/1591 *S. suis* strains

ORF (P1/7)*	ORF (89/1591)†	Putative function	AdcR-binding sequence and position‡
ssu0112§	ssuiDRAFT 1240	Regulatory protein (AdcR)	TTAACAAGTTAA (–28)
ssu0113§	ssuiDRAFT 1239	ABC transporter component; ATPase	
ssu0114§	ssuiDRAFT 1238	ABC transporter component; permease	
ssu0115§	ssuiDRAFT 1237	ABC transporter component; zinc-binding lipoprotein	
ssu0308	ssuiDRAFT 0103	ABC transporter component; zinc-binding lipoprotein	TTAACTGGTTAA (–26)
ssu0309	ssuiDRAFT 1195	Pht protein	
ssu1390	ssuiDRAFT 0326	Internalin A	TTAACTGGTTAA (–31)
ssu1352	ssuiDRAFT 0240	30S ribosomal protein S14	TTAACTGGTTAA (–36)
ssu1103	ssuiDRAFT 1606	Pht protein	TTAACTAGTAAA (–230)

*ORF name in *S. suis* strain P1/7 (http://www.sanger.ac.uk/Projects/S_suis/).

†ORF name in *S. suis* strain 89/1591 (<http://www.ncbi.nlm.nih.gov>).

‡Position with respect to its predicted translational starting point.

§||Genes constituting a single transcriptional unit. The AdcR motif is located upstream of the first gene listed.

In order to evaluate the role of the AdcR protein in the regulation of genes whose products could present immunogenic and/or protective abilities, the construction and characterization of an *adcR* in-frame deletion mutant was carried out as described in Methods. After electrotransformation with plasmid pSET4Δ*adcR* to replace the wild-type *adcR* gene of *S. suis* strain P1/7 with an *adcR*-deleted gene, the *S. suis* UA5000 strain (AdcR[–]) was obtained, with the deletion confirmed by both PCR and sequencing (data not shown). The repressor function of the AdcR protein upon these genes was confirmed by the fact that the null-*adcR* strain showed derepression of all AdcR-regulated ORFs (Fig. 2). Moreover, none of the non-*adcR* regulated ssu0626, ssu1246 and 0118 genes (lacking an AdcR-binding site in their promoter regions) was induced in the *adcR* mutant strain when compared with the wild-type strain (results not shown).

The AdcR protein regulates an *S. suis* gene that encodes a ribosomal protein paralogue

It has been proposed that genes that encode ribosomal protein paralogues are regulated by AdcR and thus contribute to survival during zinc starvation (Panina *et al.*, 2003). This hypothesis is based on the fact that original copies of ribosome proteins present zinc-ribbon motifs (C-X₂-C-X_n-C-X₂-C) and are thus likely to bind zinc, while paralogous copies lack these motifs. However, the presence of zinc in the ribosomes seems to be required to stabilize them at high temperature only in thermophilic bacteria and Archaea (Makarova *et al.*, 2001). Therefore, it has been suggested that under conditions of zinc starvation, the induction of paralogous genes could lead to the incorporation of these paralogous proteins into the ribosomes in place of the original ribosomal proteins. After degradation of the original zinc-containing ribosomal proteins, the metal becomes available to be used by other zinc-

Fig. 2. Expression factors of several ORFs (Table 1) in the *S. suis* *adcR* mutant (UA5000). The expression factor is the ratio of the mRNA concentration of each gene from the UA5000 strain with respect to the wild-type strain (P1/7). The amount of mRNA of each gene was determined by using a standard curve generated by the amplification of an internal fragment of the *S. suis* P1/7 ORF encoding cysteine synthase (see Supplementary Table S2 for primer sequences). The results are the mean of two independent experiments, each carried out in duplicate. Error bars, SD.

dependent proteins. Scenarios in concordance with this hypothesis have been reported during zinc starvation for ribosomal proteins of *B. subtilis* and *Streptococcus pyogenes* (Brenot *et al.*, 2007; Makarova *et al.*, 2001). In accordance with this, our results showed that expression of ssu1352 ORF in the *adcR*-knockout strain, encoding ribosomal protein S14, is approximately 100-fold higher than in the wild-type strain (Fig. 2). In this respect, an ORF (ssu0083) encoding the putative original 30S ribosomal protein S14 ORF is also present in the complete genome of *S. suis* strain P1/7. As expected, the predicted product of the ssu0083 ORF includes a zinc-ribbon motif (CEKCGRPHSVYRKFKLCRV) which is absent in the *AdcR*-regulated ssu1352 ORF (data not shown). Moreover, expression of the ssu0083 ORF, which lacks the *AdcR*-binding site in its promoter region, is not increased in the *adcR* mutant strain (data not shown). Although further experiments are needed to fully evaluate its validity, our data support the hypothesis that genes encoding ribosomal protein paralogues regulated by *AdcR* contribute to zinc homeostasis in streptococci.

Two Pht proteins are under *AdcR* regulation in *S. suis*

SDS-PAGE analysis of cell surface-associated proteins of both the *adcR* mutant and wild-type strains revealed the overexpression of two proteins of approximately 120 and 100 kDa in the *adcR*-knockout strain (Fig. 3). To identify them, both protein gel bands were in-gel digested with trypsin and the peptides obtained were identified by MALDI-TOF MS (data not shown). The results indicated that the two proteins corresponded to ssu0309 and ssu1103 ORF products, which have both been annotated as Pht (pneumococcal histidine triad) proteins, showing scores of

Fig. 3. SDS-PAGE profile of cell wall-associated proteins of both *S. suis* P1/7 (wt) and UA5000 (*AdcR*⁻) strains. Two overexpressed proteins of UA5000 cell extracts are indicated by arrows with their apparent molecular masses. The molecular mass marker (M) and its values are shown on the left of the figure.

165 and 73, respectively (using the MASCOT search engine, version 2.1). The predicted products of these two genes are proteins of 116.5 and 95.4 kDa molecular mass, respectively, in agreement with the apparent molecular mass seen on SDS-PAGE (Fig. 3). Pht proteins have only been found in pathogenic streptococcal species and are involved in the invasion process (Adamou *et al.*, 2001; Panina *et al.*, 2003). This family of proteins shows a conserved and repeated motif HxxHxH (histidine triad motif) which is implicated in zinc binding (Adamou *et al.*, 2001; Panina *et al.*, 2003; Riboldi-Tunncliffe *et al.*, 2005).

Overexpression of Ssu1103 and Ssu0309 proteins in the *adcR* mutant strain suggested that they are under the control of *AdcR*. In this respect, it must be noted that an *AdcR*-like motif (TTAACTAGTAAA) is found at position -230 with respect to the ssu1103 predicted translational starting point. For this reason, the ability of *S. suis* *AdcR* protein to bind the promoter region of ssu1103 ORF was analysed by EMSA. Data showed that the *AdcR* protein specifically binds to the promoter of this gene (Fig. 4).

Moreover, and since ssu0308 and ssu0309 ORFs were demonstrated to be co-transcribed by RT-PCR analysis (data not shown), a search for putative *AdcR* boxes upstream of this transcriptional unit was carried out. As expected, one motif with these characteristics (TTAACTGGTTAA) was found at position -26 with respect to the predicted translational starting point of the ssu0308 ORF, whose product is homologous (99% identity) to the protein encoded by the *AdcR*-regulated ssu1DRAFT 0103 ORF of the 89/1591 *S. suis* strain (Table 1). Furthermore, EMSA experiments demonstrated that the *S. suis* *AdcR* protein is able to bind this *AdcR*-like sequence present upstream of the ssu0308-ssu0309 transcriptional unit (data not shown). In accordance with all these data, derepression of both ssu1103 and ssu0309 ORFs was observed in the *adcR* mutant strain (Fig. 5).

Furthermore, Pht proteins are involved in the host-cell-pathogen interaction, and it has been hypothesized that they have evolved to become sensitive to *AdcR* regulation in order to exploit zinc as a compartment-specific regulatory cue (Brenot *et al.*, 2007). It is also noteworthy that Pht proteins present hydrophobic leader sequences, which presumably target these proteins to the bacterial cell surface and induce specific antibodies against pneumococcal sepsis, suggesting that Pht proteins could serve as effective vaccines against streptococcal infections (Adamou *et al.*, 2001; Kunitomo *et al.*, 2008). Our data effectively demonstrate the presence of Pht proteins at the *S. suis* cell surface and that *AdcR* protein is not only involved in cation homeostasis but also controls putative virulence factors.

Protective capacities of cell surface-associated proteins of *adcR*, *fur* and *adcR fur* *S. suis* mutants

As mentioned above, the *adcR* mutant strain overexpresses genes encoding predicted immunogenic proteins. For that

Fig. 4. (a) Schematic representation of the *ssu1103* ORF with its promoter region (P1103). Small arrows indicate the primer pairs used to amplify the promoters for EMSA experiments. (b) Electrophoretic mobility of the DNA fragment containing the indicated *S. suis* promoter in the absence (–) or presence (+) of purified *S. suis* AdcR protein.

reason, the cell surface-associated proteins of both wild-type and *adcR*-null strains were analysed on Western blots by using immunologically reactive serum obtained from surviving mice after inoculation with the *S. suis* 89/1591 wild-type strain. Results showed that the *adcR* mutant strain extracts presented more immunological proteins than the wild-type strain (Fig. 6). However, no significant protective effect was obtained when the cell surface-associated proteins of the *adcR* mutant strain were inoculated into mice prior to a challenge with the wild-type *S. suis* 89/1591 strain (data not shown). This result suggested that *adcR*-regulated proteins associated with the cell surface are not sufficient to prevent the infective process in mice. This could be due to the amount of immunogenic proteins not being sufficient or their exposure to the external medium not being as extensive

as necessary to confer good protection. In fact, previous data have demonstrated that a recombinant *adcR*-regulated protein (SsuiDRAFT 0103) is sufficient to confer significant protection against this pathogen in mice (Aranda *et al.*, 2008).

A search in the NCBI database (<http://www.ncbi.nlm.nih.gov/>) allowed the identification of the *Streptococcus thermophilus fur* gene (gene ID: 3166852). Then, this gene was used as a probe against the Sanger BLAST server (http://www.sanger.ac.uk/Projects/S_suis/), enabling the identification of the *S. suis* P1/7 *fur* gene, which has a high degree of identity (64 %) to the *S. thermophilus fur* gene. As mentioned above, the Fur protein binds iron in the cytoplasm of many bacterial cells and blocks the transcription of iron-regulated genes, several of which express strongly immunogenic proteins (Escobar *et al.*, 1999; Hantke, 1981; Ratledge & Dover, 2000). After electrotransformation with plasmid pSET4Δ*fur* to replace the wild-type *fur* gene of *S. suis* P1/7 strain with a *fur*-deleted gene, a *fur*-defective *S. suis* strain (UA5001) was obtained, with the deletion confirmed by both PCR and sequencing (data not shown). However, no significant protective effect was obtained when the cell surface-associated proteins of the *fur* mutant strain were used to inoculate mice prior to a challenge with the wild-type *S. suis* 89/1591 strain (data not shown). Then, to increase the putative immunogenic capacity of both *S. suis* mutants, the construction of a double mutant strain (UA5002, *AdcR*[–] *Fur*[–]) was carried out. The presence of both the *adcR* and *fur* gene deletions was confirmed by both PCR and sequencing (data not shown).

Analysis of the SDS-PAGE profile of *S. suis* UA5002 (*AdcR*[–] *Fur*[–]) cell surface-associated proteins revealed that the double mutant strain presented a higher number of immunogenic proteins than both *S. suis* single mutants (Fig. 6). In order to determine whether the induction of this immunogenic response was enough to protect mice against *S. suis* infection, three groups of eight mice were immunized twice with 50 µg cell surface-associated proteins of the double mutant strain, 50 µg cell surface-associated proteins of the wild-type strain, or the solvent including the adjuvant (as a control). When the animals were challenged with 10⁸ c.f.u. *S. suis* 89/1591, all the mice belonging to either the control group or the group

Fig. 5. Expression factors of Pht-encoding *ssu0309* and *ssu1103* ORFs in the *S. suis* *adcR* mutant (UA5000). The expression factor is the ratio of the mRNA concentration of each gene from the UA5000 strain with respect to the wild-type strain (P1/7). The amount of mRNA of each gene was determined by using a standard curve generated by the amplification of an internal fragment of the *S. suis* P1/7 ORF encoding cysteine synthase (see Supplementary Table S2 for primer sequences). The results are the means of two independent experiments, each carried out in duplicate. Error bars, SD.

Fig. 6. (a) SDS-PAGE profile of *S. suis* P1/7 wild-type (wt), UA5000 (AdcR⁻), UA5001 (Fur⁻) and UA5002 (AdcR⁻ Fur⁻) strains. (b) Western blot analysis with serum recovered from mice infected with *S. suis* strain 89/1591. The analysis was carried out with cell surface-associated proteins obtained from *S. suis* P1/7 wild-type (wt), UA5000 (AdcR⁻), UA5001 (Fur⁻) and UA5002 (AdcR⁻ Fur⁻) strains.

vaccinated with the wild type extract died within 5 days of infection (Fig. 7). In contrast, 50 % of mice in the group vaccinated with the double mutant extract inoculated with an identical dose of the virulent strain remained alive throughout the experiment (Fig. 7). In addition, both wild-type and *adcR fur* double mutant cell surface-associated proteins were analysed on Western blots by using immunologically reactive serum obtained from mice surviving after inoculation with the *S. suis* 89/1591 wild-

type strain. The results provided clear evidence that the protective effect of the double mutant extract was correlated with specific antibodies against the overexpressed antigens of this strain (Fig. 6). It is worth noting that negative results (no reactivity) were obtained when all the *S. suis* extracts used in this work were tested with serum from non-infected mice (data not shown). All these data indicated that the combination of cell surface-associated proteins under the control of either zinc/manganese- or iron-uptake regulators (AdcR and Fur, respectively) not only is immunogenic but also confers a statistically significant level of protection ($P=0.038$) against *S. suis* challenge to mice immunized with these proteins.

Fig. 7. Survival curve of mice vaccinated with cell surface-associated proteins of wild-type P1/7 strain (●), cell surface-associated proteins of *adcR fur* UA5002 strain (■), or solvent including the adjuvant as a negative control (▲). After the last immunization, an intraperitoneal challenge with 10^8 c.f.u. wild-type *S. suis* 89/1591 cells was administered to each animal ($n=8$ mice per group).

To sum up, the data reported in this work demonstrate that the use of cell surface extracts of *S. suis* double mutants defective in genes regulating either Zn^{2+}/Mn^{2+} or Fe^{2+} cations is able to protect mice against challenge with *S. suis*. This double mutant-based strategy has not to our knowledge been used to date and can be applied to other virulent bacterial species. Thus, mutagenesis of this kind of regulator makes it possible to obtain effects in the overexpression of divalent cation-uptake transporters on the bacterial cell surface similar to those induced by chelating agents, overcoming problems associated with the use of these chemicals, such as poor bacterial cell growth, and simplifying the bulk production of this antigen source. In addition, the data reported in this work indicate that the use of purified immunogenic proteins such as Pht or SsuiDRAFT 0103 is a more promising vaccination strategy than the use of protein extracts.

ACKNOWLEDGEMENTS

We are very grateful to Dr T. Sekizaki, National Institute of Animal Health, Ibaraki, Japan, for providing the pSET4s plasmid used in this work. We thank Joan Ruiz and Sonia Lacouture for their excellent

technical assistance. This work was funded by grants AGL2005-03574 from the Ministerio de Educación y Ciencia (MEC) de España and 2005SGR-533 from the Departament d'Universitats, Recerca i Societat de la Informació (DURSI) de la Generalitat de Catalunya. J.A. is the recipient of a pre-doctoral fellowship from the Universitat Autònoma de Barcelona. N.F. is the recipient of a Natural Sciences and Engineering Research Council of Canada postgraduate scholarship.

REFERENCES

- Abella, M., Rodríguez, S., Paytubi, S., Campoy, S., White, M. F. & Barbé, J. (2007). The *Sulfolobus solfataricus* *radA* paralogue sso0777 is DNA damage inducible and positively regulated by the Sta1 protein. *Nucleic Acids Res* **35**, 6788–6797.
- Adamou, J. E., Heinrichs, J. H., Erwin, A. L., Walsh, W., Gayle, T., Dormitzer, M., Dagan, R., Brewah, Y. A., Barren, P. & other authors (2001). Identification and characterization of a novel family of pneumococcal proteins that are protective against sepsis. *Infect Immun* **69**, 949–958.
- Andrews, S. C., Robinson, A. K. & Rodríguez-Quinones, F. (2003). Bacterial iron homeostasis. *FEMS Microbiol Rev* **27**, 215–237.
- Aranda, J., Garrido, M. E., Cortes, P., Llagostera, M. & Barbé, J. (2008). Analysis of the protective capacity of three *Streptococcus suis* proteins induced under divalent-cation-limited conditions. *Infect Immun* **76**, 1590–1598.
- Babich, H. & Stotzky, G. (1978). Toxicity of zinc to fungi, bacteria, and coliphages: influence of chloride ions. *Appl Environ Microbiol* **36**, 906–914.
- Brenot, A., Weston, B. F. & Caparon, M. G. (2007). A PerR-regulated metal transporter (PmtA) is an interface between oxidative stress and metal homeostasis in *Streptococcus pyogenes*. *Mol Microbiol* **63**, 1185–1196.
- Bsat, N. & Helmann, J. D. (1999). Interaction of *Bacillus subtilis* Fur (ferric uptake repressor) with the *dhb* operator in vitro and in vivo. *J Bacteriol* **181**, 4299–4307.
- Campoy, S., Mazon, G., Fernandez de Henestrosa, A. R., Llagostera, M., Monteiro, P. B. & Barbe, J. (2002). A new regulatory DNA motif of the gamma subclass *Proteobacteria*: identification of the LexA protein binding site of the plant pathogen *Xylella fastidiosa*. *Microbiology* **148**, 3583–3597.
- Chamngongpol, S., Dodson, W., Cromie, M. J., Harris, Z. L. & Groisman, E. A. (2002). Fe(III)-mediated cellular toxicity. *Mol Microbiol* **45**, 711–719.
- Escolar, L., Perez-Martin, J. & de Lorenzo, V. (1999). Opening the iron box: transcriptional metalloregulation by the Fur protein. *J Bacteriol* **181**, 6223–6229.
- Fontaine, M. C., Perez-Casal, J. & Willson, P. J. (2004). Investigation of a novel DNase of *Streptococcus suis* serotype 2. *Infect Immun* **72**, 774–781.
- Garmory, H. S. & Titball, R. W. (2004). ATP-binding cassette transporters are targets for the development of antibacterial vaccines and therapies. *Infect Immun* **72**, 6757–6763.
- Garrido, M. E., Bosch, M., Bigas, A., Badiola, I., Barbé, J. & Llagostera, M. (2008). Heterologous protective immunization elicited in mice by *Pasteurella multocida* fur *ompH*. *Int Microbiol* **11**, 17–24.
- Gottschalk, M. & Segura, M. (2000). The pathogenesis of the meningitis caused by *Streptococcus suis*: the unresolved questions. *Vet Microbiol* **76**, 259–272.
- Hantke, K. (1981). Regulation of ferric iron transport in *Escherichia coli* K12: isolation of a constitutive mutant. *Mol Gen Genet* **182**, 288–292.
- Hantke, K. (2001). Bacterial zinc transporters and regulators. *Biometals* **14**, 239–249.
- Higgins, R., Gottschalk, M., Boudreau, M., Lebrun, A. & Henrichsen, J. (1995). Description of six new capsular types (29–34) of *Streptococcus suis*. *J Vet Diagn Invest* **7**, 405–406.
- Kunitomo, E., Terao, Y., Okamoto, S., Rikimaru, T., Hamada, S. & Kawabata, S. (2008). Molecular and biological characterization of histidine triad protein in group A streptococci. *Microbes Infect* **10**, 414–423.
- Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680–685.
- Loo, C. Y., Mittrakul, K., Voss, I. B., Hughes, C. V. & Ganeshkumar, N. (2003). Involvement of the *adc* operon and manganese homeostasis in *Streptococcus gordonii* biofilm formation. *J Bacteriol* **185**, 2887–2900.
- Makarova, K. S., Ponomarev, V. A. & Koonin, E. V. (2001). Two C or not two C: recurrent disruption of Zn-ribbons, gene duplication, lineage-specific gene loss, and horizontal gene transfer in evolution of bacterial ribosomal proteins. *Genome Biol* **2**, RESEARCH 0033.
- Miller, J. H. (1992). *A Short Course in Bacterial Genetics*. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory.
- Panina, E. M., Mironov, A. A. & Gelfand, M. S. (2003). Comparative genomics of bacterial zinc regulons: enhanced ion transport, pathogenesis, and rearrangement of ribosomal proteins. *Proc Natl Acad Sci U S A* **100**, 9912–9917.
- Patzner, S. I. & Hantke, K. (2000). The zinc-responsive regulator Zur and its control of the *znu* gene cluster encoding the ZnuABC zinc uptake system in *Escherichia coli*. *J Biol Chem* **275**, 24321–24332.
- Ratledge, C. & Dover, L. G. (2000). Iron metabolism in pathogenic bacteria. *Annu Rev Microbiol* **54**, 881–941.
- Riboldi-Tunnicliffe, A., Isaacs, N. W. & Mitchell, T. J. (2005). 1.2 Å crystal structure of the *S. pneumoniae* PhtA histidine triad domain a novel zinc binding fold. *FEBS Lett* **579**, 5353–5360.
- Sambrook, J. & Russell, D. W. (2001). *Molecular Cloning: a Laboratory Manual*, 3rd edn. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory.
- Staats, J. J., Feder, I., Okwumabua, O. & Chengappa, M. M. (1997). *Streptococcus suis*: past and present. *Vet Res Commun* **21**, 381–407.
- Sutcliffe, I. C. & Russell, R. R. (1995). Lipoproteins of Gram-positive bacteria. *J Bacteriol* **177**, 1123–1128.
- Takamatsu, D., Osaki, M. & Sekizaki, T. (2001a). Construction and characterization of *Streptococcus suis*–*Escherichia coli* shuttle cloning vectors. *Plasmid* **45**, 101–113.
- Takamatsu, D., Osaki, M. & Sekizaki, T. (2001b). Thermosensitive suicide vectors for gene replacement in *Streptococcus suis*. *Plasmid* **46**, 140–148.
- Tanabe, M., Atkins, H. S., Harland, D. N., Elvin, S. J., Stagg, A. J., Mirza, O., Titball, R. W., Byrne, B. & Brown, K. A. (2006). The ABC transporter protein OppA provides protection against experimental *Yersinia pestis* infection. *Infect Immun* **74**, 3687–3691.
- Tavares, F. & Sellstedt, A. (2000). A simple, rapid and non-destructive procedure to extract cell wall-associated proteins from *Frankia*. *J Microbiol Methods* **39**, 171–178.
- Touati, D. (2000). Iron and oxidative stress in bacteria. *Arch Biochem Biophys* **373**, 1–6.
- Wollert, T., Heinz, D. W. & Schubert, W. D. (2007). Thermodynamically reengineering the listerial invasion complex InlA/E-cadherin. *Proc Natl Acad Sci U S A* **104**, 13960–13965.
- Yu, H., Jing, H., Chen, Z., Zheng, H., Zhu, X., Wang, H., Wang, S., Liu, L., Zu, R. & other authors (2006). Human *Streptococcus suis* outbreak, Sichuan, China. *Emerg Infect Dis* **12**, 914–920.

Edited by: T. J. Mitchell