Protein phosphatase CaPpz1 is involved in cation homeostasis, cell wall integrity and virulence of *Candida albicans*

Csaba Ádám,¹ Éva Erdei,² Carlos Casado,³ László Kovács,¹ Asier González,³ László Majoros,⁴ Katalin Petrényi,¹ Péter Bagossi,^{5†} Ilona Farkas,¹ Monika Molnar,^{2‡} István Pócsi,² Joaquín Ariño³ and Viktor Dombrádi^{1,6}

Correspondence

Viktor Dombrádi
dombradi@med.unideb.hu

¹Department of Medical Chemistry, Faculty of Medicine, Research Centre for Molecular Medicine, University of Debrecen, Debrecen, Hungary

²Department of Microbial Biotechnology and Cell Biology, Faculty of Science, University of Debrecen, Debrecen, Hungary

³Institut de Biotecnologia i Biomedicina, Departament de Bioquímica i Biologia Molecular, Universitat Autònoma Barcelona, Cerdanyola del Vallès, Spain

⁴Department of Medical Microbiology, Faculty of Medicine, University of Debrecen, Debrecen, Hungary

⁵Department of Biochemistry and Molecular Biology, Faculty of Medicine, University of Debrecen, Debrecen, Hungary

⁶Cell Biology and Signalling Research Group of the Hungarian Academy of Sciences, University of Debrecen, Debrecen, Hungary

The opportunistic pathogen *Candida albicans* has a single protein phosphatase Z (PPZ) candidate gene termed *CaPPZ1*, which shows significant allele variability. We demonstrate here that bacterially expressed CaPpz1 protein exhibits phosphatase activity which can be inhibited by recombinant Hal3, a known inhibitor of *Saccharomyces cerevisiae* Ppz1. Site-directed mutagenesis experiments based on natural polymorphisms allowed the identification of three amino acid residues that affect enzyme activity or stability. The expression of *CaPPZ1* in *ppz1* *S. cerevisiae* and *pzh1* *Schizosaccharomyces pombe* cells partially rescued the salt and caffeine phenotypes of the deletion mutants. CaPpz1 also complemented the *slt2* *S. cerevisiae* mutant, which is crippled in the mitogen-activated protein (MAP) kinase that mediates the cell wall integrity signalling pathway. Collectively, our results suggest that the orthologous PPZ enzymes have similar but not identical functions in different fungi. The deletion of the *CaPPZ1* gene in *C. albicans* resulted in a mutant that was sensitive to salts such as LiCl and KCl, to caffeine, and to agents that affect cell wall biogenesis such as Calcofluor White and Congo red, but was tolerant to spermine and hygromycin B. Reintegration of the *CaPPZ1* gene into the deletion mutant alleviated all of the mutant phenotypes tested. Thus CaPpz1 is involved in cation homeostasis, cell wall integrity and the regulation of the membrane potential of *C. albicans*. In addition, the germ tube growth rate, and virulence in the BALB/c mouse model, were reduced in the null mutant, suggesting a novel function for CaPpz1 in the yeast to hypha transition that may have medical relevance.

Received 21 December 2011

Revised 7 February 2012

Accepted 9 February 2012

[†]With sadness we note that our colleague and co-author passed away on 22 July 2011.

[‡]Present address: Institute of Environmental Sciences, Faculty of Science and Informatics, College of Nyiregyhaza, Nyiregyhaza, Hungary.

Abbreviations: CFW, Calcofluor White; CWI pathway, cell wall integrity pathway; MAP kinase, mitogen-activated protein kinase; PPZ, protein phosphatase Z. Three supplementary figures and a supplementary table are available with the online version of this paper.

INTRODUCTION

The Ppz Ser/Thr protein phosphatases constitute a family of enzymes that are structurally related to the type-1 protein phosphatases, and are restricted to fungal species (Ariño, 2002). In *Saccharomyces cerevisiae*, where these proteins were discovered (Lee *et al.*, 1993; Posas *et al.*, 1992, 1993), there exist two genes, namely *PPZ1* and *PPZ2*. To avoid confusion, these two phosphatase genes will be referred to as *ScPPZ1* and *ScPPZ2* in the present paper. The encoded proteins contain C-terminal catalytic domains, which are highly similar to each other, while their N-terminal moieties are much less related (Fig. 1a). Deletion of *ScPPZ2* does not result in a detectable phenotype. However, strains lacking *ScPpz1* exhibit increased tolerance to NaCl and LiCl (Posas *et al.*, 1995), altered K⁺ influx, and hypersensitivity to caffeine (Posas *et al.*, 1993; Yenush *et al.*, 2002). On the other hand, overexpression of *ScPPZ1* results in slow growth and cell cycle blockage at the G1–S transition (Clotet *et al.*, 1999). The effect of the *Scppz1* mutation on cation homeostasis is due to the elevated expression of the Na⁺-ATPase gene *ENA1* (Posas *et al.*, 1995; Ruiz *et al.*, 2003) and to the increased influx of potassium ions mediated by the high-affinity K⁺-transporter Trk1 (Yenush *et al.*, 2002). *ScPPZ1* exhibits genetic interactions with *SLT2*, which encodes a mitogen-activated protein (MAP) kinase required for signalling in the cell wall integrity (CWI) pathway (reviewed by Levin, 2005). *slt2* cells are prone to lysis and are very sensitive to high temperature, caffeine, or cell wall-damaging compounds. Overexpression of *ScPPZ1* suppresses the lytic phenotype of an *slt2* mutant, whereas deletion of the *ScPPZ1* gene or inhibition of its phosphatase activity results in a phenotype additive to that of the *slt2* strain (de Nadal *et al.*, 1998; Lee *et al.*, 1993). The functional interaction between *ScPpz1* and the CWI pathway has been explained on the basis of the combination of increased internal turgor pressure in Ppz-deficient strains and cell wall instability observed in strains lacking *Slr2* (Merchan *et al.*, 2004). Therefore, in budding yeast, the Ppz proteins play key roles in cation homeostasis, which is likely to affect cell cycle regulation. The Hal3 and Vhs3 regulatory subunits bind to the catalytic domain of *ScPpz1* and inhibit its activity (de Nadal *et al.*, 1998; Ruiz *et al.*, 2004). The overexpression of these inhibitors mimics the effects of the *scppz1* mutation.

The protein phosphatase Z (PPZ) phosphatases have been only partially characterized in other yeasts or fungi. *Schizosaccharomyces pombe* contains the *pzh1* gene, encoding a protein that is similar to budding yeast Ppz (Fig. 1a), but has a shorter and rather divergent N-terminal domain (Balcels *et al.*, 1997). In *Sch. pombe*, the deletion of *pzh1*⁺ results in cells hypertolerant to Na⁺ and hypersensitive to K⁺ (Balcels *et al.*, 1997). However, the mechanisms of action of Pzh1 in fission yeast are probably different from those observed for *ScPpz1* in budding yeast, since cells lacking Pzh1 have no altered sodium or lithium efflux; rather, they display decreased influx for these cations, together with a reduced K⁺ efflux (Balcels *et al.*, 1999). A

Fig. 1. Structure of the CaPpz1 protein. (a) Scheme representing the amino acid sequence of CaPpz1 as compared with those of the *S. cerevisiae*, *N. crassa* and *Sch. pombe* orthologues. The N-terminal moiety, including a myristoylation site (asterisk), is green, the conserved catalytic domain is red and the variable, disordered segments are represented by white boxes. (b) Homology model of the CaPpz1 catalytic domain. α -Helices are red, β -sheets are blue and loops are yellow. Two essential metal ions in the catalytic centre are grey and the amino acid side chains involved in catalysis are in violet. Three amino acid residues that are affected by natural polymorphisms are highlighted in light brown and one replacement caused by the specific codon usage of *C. albicans* is yellow. The amino acid exchanges analysed in the present study are labelled with arrows.

PPZ phosphatase gene from the filamentous fungus *Neurospora crassa*, named *Pzl-1*, has been identified by Szöör *et al.* (1998). The PZL-1 protein (Fig. 1a) was expressed in *S. cerevisiae* and was shown to fulfil every known function carried out by its *S. cerevisiae* counterpart, despite the marked sequence divergence within their N-terminal moieties (Vissi *et al.*, 2001). The expression of PZL-1 in the *pzh1* mutant *Sch. pombe* resulted in a less

efficient complementation. Recently, a PPZ orthologue, termed DhPpz1, has been characterized in the extremely halotolerant yeast species *Debaryomyces hansenii* (Minhas *et al.*, 2012). This phosphatase also has a disordered N-terminal segment that includes a short conserved Ser/Arg-rich motif which is important in salt tolerance but not in cell wall integrity. Interestingly, this fungus utilizes a Na^+/H^+ antiporter to evade the toxic effects of cations. The comparison of known PPZ enzymes suggests that although their major functions are retained across fungi, the underlying mechanisms can be different.

Candida albicans is an opportunistic pathogen with considerable medical significance. This organism contains a single PPZ candidate gene, termed *CaPPZ1*, which has at least four distinct alleles (Kovács *et al.*, 2010). The allele combinations in the diploid organism together with individual point mutations result in a great genetic variability. The *CaPPZ1* gene encodes a protein whose primary structure is similar to those of its better-characterized fungal counterparts (Fig. 1a and Supplementary Fig. S1 available with the online version of this paper). Homology modelling suggests that the 3D structure of the CaPpz1 catalytic domain is reminiscent of that of the protein phosphatase 1 catalytic subunit (Fig. 1b). The physiological significance of the *C. albicans* PPZ phosphatase has not yet been discovered. The only available information extracted from large-scale genetic screens tells us that the disruption of either one (Xu *et al.*, 2007) or both (Hanaoka *et al.*, 2008) of the alleles is not detrimental. In the present work we characterize the function of this enzyme by biochemical assays of the recombinant protein, by expressing the *CaPPZ1* gene in Ppz-deficient *S. cerevisiae* and *Sch. pombe* strains, as well as by studying the relevant *C. albicans* mutant. Our work reveals that despite the structural similarities, CaPpz1 only partially complements the lack of its orthologues in other fungi and has a novel function in controlling germ tube formation in *C. albicans*.

METHODS

DNA cloning. The cloning of *C. albicans* *CaPPZ1* gene has been described previously (Kovács *et al.*, 2010). The *CaPPZ1-3* allele (Genbank accession no. GQ487308) was used to insert the ORF into a pET28a+ expression vector (Novagen) in two steps. The two halves of the coding region were amplified by PCR with the UPPZ-NdeI/LPPZ-NdeI and UPPZ-PstI/LPPZ-XhoI primer pairs (Table S1a), and were ligated together, resulting in the *CaPPZ1*-pET28a+ plasmid. In order to confirm the predicted gene structure, we also cloned the corresponding cDNA from the ATCC 10231 reference strain. Total RNA was prepared with the SV Total RNA Isolation System (Promega), and the mRNA was reverse-transcribed with the 5'/3' RACE kit (Roche). The cDNA was amplified by PCR using the 2U and 2L primers (Table S1a), and was placed into a pGEM-T Easy vector (Promega) to yield the *CaPPZ1*-pGEM-T plasmid, whose sequence has been deposited in the NCBI database under accession number JF330253.

For expression in *S. cerevisiae* under the control of the *ScPPZ1* promoter, first the -525/-10 segment of the promoter region was

amplified by PCR from *S. cerevisiae* FY1679 genomic DNA with ScPromSacI and ScPromXbaI primers (Table S1a) according to Vissi *et al.* (2001) and the amplicon was cloned into plasmid pGEM-T Easy. Next, the promoter region was released with SacI and XbaI and cloned into plasmids YCplac111 and YEplac181. Then, the coding region of the *CaPPZ1-3* allele was amplified from the *CaPPZ1*-pGEM-T cDNA clone by PCR with C1XbaI and C2HindIII primers (Table S1a) and placed after the promoter to generate plasmids YCp-*CaPPZ1* and YE-*CaPPZ1*. The construction of the control plasmid YCp111-*ScPPZ1* has been described elsewhere (Clotet *et al.*, 1996).

For expression in *Sch. pombe*, the coding region of *CaPPZ1-3* was amplified from the *CaPPZ1*-pGEM-T construct by PCR with C3BamHI and C4BamHI primers (Table S1a) and inserted into the expression vector pREP41 (Basi *et al.*, 1993; Maundrell, 1993), producing the pREP41-*CaPPZ1* plasmid. All of the vector constructs were confirmed by DNA sequencing.

In vitro mutagenesis. The positions of the mutated amino acids are shown in Figs 1(b) and S1. The homology model of the CaPpz1 catalytic domain was built on the crystal structure of rabbit muscle PP1 catalytic subunit fragment (amino acid residues 7–300; Goldberg *et al.*, 1995; PDB accession no. 1FJM) using the Modeller 7 program (Sali & Blundell, 1993). Sequence alignment was done by CLUSTAL W (Thompson *et al.*, 1994). Mutations resulting in single or double amino acid exchanges were introduced into the *CaPPZ1-3* sequence with the aid of the QuikChange Lightning Site-Directed Mutagenesis kit (Agilent Technologies). PCR of the *CaPPZ1*-pET28a+ plasmid target with the oligonucleotide primers described in Table S1(b) resulted in the R262L, D261N, G333E and C337R mutations, while the double mutant encompassing both G333E and C337R replacements was generated from the G333E mutant by a second round of mutagenesis.

Expression of *CaPPZ1* in *Escherichia coli*. Wild-type and mutated CaPpz1-3 was expressed in *E. coli* BL21(DE3)-RIL (Stratagene) after the addition of 0.4 mM IPTG (Sigma-Aldrich) followed by 16 h incubation at 18 °C (Fig. S2a). The recombinant proteins were purified from the supernatant of the bacterial extract by Ni-NTA Agarose (Qiagen) affinity chromatography. The fractions eluted from the columns were analysed by SDS-PAGE (Fig. S2b). The protein concentration of the fractions was assayed by the Bradford method (Bradford, 1976). The CaPpz1 content in the peak fractions was estimated by scanning of the electrophoretograms. The phosphatase activity of the fractions was assayed in triplicate samples containing 1–2 µg recombinant phosphatase with *p*-nitrophenylphosphate (Sigma-Aldrich) substrate in the presence or absence of Hal3, as reported previously (Muñoz *et al.*, 2004). Recombinant *S. cerevisiae* Hal3 was expressed in *E. coli* and purified according to Garcia-Gimeno *et al.* (2003).

Expression of *CaPPZ1* in *S. cerevisiae*. The genotypes and origin of the *S. cerevisiae* strains used in this work are listed in Table 1. Single *kanMX* deletion mutants in the BY4741 background were generated in the context of the *Saccharomyces* Genome Deletion Project (Winzeler *et al.*, 1999). The mutants were transformed with different rescue plasmids as in Vissi *et al.* (2001). *S. cerevisiae* cells were grown at 28 °C in YPD medium (10 g yeast extract l⁻¹, 20 g peptone l⁻¹ and 20 g glucose l⁻¹) or, when carrying plasmids, in synthetic complete drop-out medium (lacking leucine). Sensitivity of yeast cells to LiCl, NaCl, caffeine (Merck) or Calcofluor White (CFW; Sigma) was evaluated by growth on agar plates (drop tests), as previously described (Posas *et al.*, 1995).

Expression of *CaPPZ1* in *Sch. pombe*. The genotypes and origin of the *Sch. pombe* strains are summarized in Table 1. The *pzh1*⁺ control strain was isolated from a cross between the *Sch. pombe* *h*⁺

Table 1. Fungal strains used in the present study

Strain	Genotype	Reference or source
<i>S. cerevisiae</i> FY1679	MATa/α <i>ura3-52/ura3-52 trp1Δ63/TRP1 leu2Δ1/LEU2 his3Δ200/HIS3 GAL2/GAL2</i>	Winston <i>et al.</i> (1995)
<i>S. cerevisiae</i> BY4741	MATa <i>his3Δ1 leu2Δ met15Δ ura3Δ</i>	Winzeler <i>et al.</i> (1999)
<i>S. cerevisiae</i> BY4741 <i>ppz1::KanMX4</i>	MATa <i>his3Δ1 leu2Δ met15Δ ura3Δ ppz1::kanMX4</i>	Winzeler <i>et al.</i> (1999)
<i>S. cerevisiae</i> BY4741 <i>slt2::KanMX4</i>	MATa <i>his3Δ1 leu2Δ met15Δ ura3Δ slt2::kanMX4</i>	Winzeler <i>et al.</i> (1999)
<i>Sch. pombe</i> LB2	<i>h⁻ ade6-M210 leu1-32 pzh1::ura4⁺ ura-D18</i>	Balcells <i>et al.</i> (1997)
<i>Sch. pombe</i> pzh1 ⁺	<i>h⁻ ade6-M210 leu1-32</i>	This study
<i>C. albicans</i> SN87	<i>ura3Δ-iro1Δ::imm⁴³⁴/URA3-IRO1, his1Δ/his1Δ, leu2Δ/leu2Δ</i>	Noble & Johnson (2005)
<i>C. albicans</i> <i>cappz1</i>	<i>ura3Δ-iro1Δ::imm⁴³⁴/URA3-IRO1, his1Δ/his1Δ, leu2Δ/leu2Δ ppz1Δ::HIS1/ppz1Δ::LEU2</i>	This study
<i>C. albicans</i> <i>CaPPZ1-HIS1</i>	<i>ura3Δ-iro1Δ::imm⁴³⁴/URA3-IRO1, his1Δ/his1Δ, leu2Δ/leu2Δ ppz1Δ::HIS1/ppz1Δ::LEU2::PPZ1</i>	This study
<i>C. albicans</i> <i>CaPPZ1-LEU2</i>	<i>ura3Δ-iro1Δ::imm⁴³⁴/URA3-IRO1, his1Δ/his1Δ, leu2Δ/leu2Δ ppz1Δ::LEU2/ppz1Δ::HIS1::PPZ1</i>	This study

wild-type strain and *Sch. pombe* 117 strain (*h⁻ ade6-M210 leu 1-32 ura4-D18*). For cultivation of *Sch. pombe*, a standard complete medium, yeast extract agar (YEA), yeast extract liquid (YEL) (Gutz *et al.*, 1974) or Edinburgh minimal medium 2 (EMM2; US Biological), as plates or liquid cultures, was used with appropriate supplements (100 mg l⁻¹ adenine, uracil, leucine). Fission yeast strains were transformed by the lithium acetate method (Ito *et al.*, 1983). Transformed cells grew on EMM2+ade medium, which contained 400 μmol thiamine l⁻¹ to repress the *nmt1* (no message in thiamine) promoter of the vectors. In our assays, first the transformants were grown in EMM2+ade (without thiamine) overnight (for the expression of the pREP construct), and these pre-cultures were used to inoculate EMM2+ade liquid media supplemented with different salts or caffeine at the following concentrations: 300, 350 and 400 mM KCl; 150, 200 and 250 mM NaCl, 5, 8 and 10 mM LiCl, as well as 5, 10 and 15 mM caffeine. The starting OD₅₉₅ values were set to 0.2, and cell growth was characterized by measuring the optical density of the cultures after 24 h incubations.

Disruption of the *CaPPZ1* gene in *C. albicans*. The SN87 strain was used for gene disruption (Table 1). By cloning and sequencing of the *CaPPZ1* gene according to Kovács *et al.* (2010) we proved that this strain was homozygous for the *CaPPZ1-1* allele. *C. albicans* strains were cultivated in YPD medium at 37 °C, while the transformants were selected and grown on SD solid medium (6.7 g l⁻¹ yeast nitrogen base with ammonium sulphate, 20 g glucose l⁻¹ and 15 g agar l⁻¹, with or without the amino acids 100 mg leucine l⁻¹ and 20 mg histidine l⁻¹, as needed).

The *CaPPZ1-1* gene was deleted by using a PCR-based procedure with primers harbouring approximately 100 bp regions from the 5'- and 3'-flanking sequences of the target gene (Gola *et al.*, 2003). The *cappz1* strain was generated by successive transformations with two disruption cassettes containing the *HIS1* and *LEU2* auxotrophic markers. The cassettes were amplified from the pFA-CdHIS1 and pFA-CmLeu2 plasmids (Schaub *et al.*, 2006) with the primers described in Table S1(c), and were transformed into the SN87 strain by electroporation in two steps (Noble & Johnson, 2005).

Characterization of *C. albicans* strains. For the validation of gene disruption, genomic DNA was extracted from *C. albicans* as described by Lee *et al.* (1988). The integration of the disruption cassettes was tested by PCR using 'diagnostic' primers (Table S1d, Fig. S3a, b). Southern hybridizations were performed with cassette-specific probes

generated by PCR (Table S1e) to verify the deletions and to exclude the possibility of the ectopic integration of disruption cassettes (Fig. S3c). The absence of *CaPPZ1* transcripts from the *cappz1* strain was confirmed by RT-PCR with the primers given in Table S1(f) (Fig. S3b).

One functional *CaPPZ1* gene copy was reintegrated into the *cappz1* mutant with the aid of the *SAT1* flipper containing pSFS2A vector (Reuss *et al.*, 2004) as described in Fig. S3(d). The presence of the *CaPPZ1* gene and either the *HIS1* or the *LEU2* cassette in the heterozygous transformants was demonstrated by PCR. It was also shown that the *CaPPZ1-HIS1* strain exhibited a His⁺ and Leu⁻ phenotype, while the *CaPPZ1-LEU2* strain was His⁻ and Leu⁺. The genotypes of the *C. albicans* strains investigated in the present study are described in Table 1.

The sensitivity of the *C. albicans* cells to salts (LiCl, NaCl and KCl), toxic cations (spermine and hygromycin B), cell wall-damaging agents (CFW, Congo red) and caffeine was evaluated in YPD medium. The relative growth of the liquid cultures was measured in 96-well plates using a Multiskan RC ELISA reader (Thermo Labsystems) at OD₆₂₀ after 18 h incubations at 37 °C. The starting OD₆₂₀ was 0.0005 in all experiments.

Hyphal growth was induced by the addition of 90 % sheep serum, and the germination capacity of the cells was determined microscopically (Gyetvai *et al.*, 2007). More than 100 cells were counted in each experiment. For better visualization of hyphae the fungal samples were stained with CFW according to the manufacturer's recommendations (<http://www.sigmaaldrich.com/etc/medialib/docs/Fluka/Datasheet/18909dat.Par.0001.File.tmp/18909dat.pdf>), and analysed under an Olympus BX40 immersion microscope.

The virulence of the *C. albicans* strains was analysed in immuno-competent female BALB/c mice essentially as in Noble & Johnson (2005), with the exception that lower doses of pathogen were used for inoculation. Higher doses of fungi led to 80–90 % lethality within 2–3 days. To prevent bacterial infections all mice received ceftazidime (5 mg day⁻¹ subcutaneously) every day after infection. Ten mice per *Candida* strain were inoculated through the lateral tail vein with 1.8–2.4 × 10⁵ c.f.u. per mouse in three independent experiments. Mice were followed up for 14 days and the survival rate was analysed by Kaplan–Meyer test (GraphPad Prism v4.03 software). The animal experiments were approved by the Animal Care Committee of the University of Debrecen, Debrecen, Hungary (permission no. 12/2008).

Fig. 2. Biochemical properties of *C. albicans* CaPpz1. The specific activity of wild-type and mutated CaPpz1 phosphatases was determined with *p*-nitrophenylphosphate. *S. cerevisiae* Hal3 protein was added to the wild-type enzyme in a 1 : 1 molar ratio (+ Hal3). The means $\pm$ SD of three to five independent experiments performed with two independent preparations are shown.

RESULTS

Biochemical characterization of the CaPpz1 protein

First we tested the phosphatase activity of the *CaPPZ1* gene product. We demonstrated that the bacterially expressed and purified recombinant CaPpz1-3 dephosphorylated *p*-nitrophenylphosphate, and that this phosphatase activity was inhibited by recombinant *S. cerevisiae* Hal3 (Fig. 2). Thus the enzymic properties of the wild-type CaPpz1 are similar to those of ScPpz1.

Next the functional significance of four selected amino acid residues in the catalytic domain of CaPpz1 (Fig. 1b) was analysed by site-directed mutagenesis. Expression in *E. coli* and purification of the mutated proteins were accomplished with the same efficiency as the control wild-type enzyme (Fig. S1), suggesting that the mutations did not affect either the production or the solubility of the phosphatase. As a proof of the principle we replaced an essential amino acid in the catalytic cleft and found that the R262L mutation resulted in an inactive protein (Fig. 2), as expected from a previous publication (Clotet *et al.*, 1996) that reported that an analogous point mutation in *S. cerevisiae* ScPpz1 eliminates phosphatase activity. Then the effects of naturally occurring polymorphisms (Kovács *et al.*, 2010) were investigated in the same way. The D261N mutation caused a moderate reduction in the phosphatase activity. When G333 was modified to E the phosphatase reaction rate doubled. On the other hand, the C337R replacement completely eradicated the catalytic reaction. Likewise, the G333E/C337R double mutation rendered the recombinant phosphatase inactive.

Complementation of the phenotypes of mutant yeasts by expression CaPpz1

Based on the structural and biochemical similarities between CaPpz1 and the ScPpz1 (Figs 1a, 2 and S1), we examined whether the *C. albicans* protein was able to complement the phenotypes of the *S. cerevisiae* *scppz1* deletion mutant. CaPpz1-3 was expressed in the mutant budding yeast cells under the control of the *ScPPZ1* promoter from either a low-copy-number centromeric YCp-*CaPPZ1* or a high-copy-number episomal YE-*CaPPZ1* plasmid (Fig. 3a). The heterologous *C. albicans* phosphatase partially normalized the tolerance to LiCl of the *S. cerevisiae* *scppz1* mutant, but was somewhat ineffective when cells were challenged with 1 M NaCl. In contrast, *CaPPZ1* markedly rescued the growth defect of the *ppz1* mutant in the presence of caffeine and, when in high copy, resulted in a near wild-type phenotype.

A strain lacking the *SLT2* MAP kinase gene shows strong hypersensitivity to compounds that affect cell wall synthesis, such as caffeine or CFW. Overexpression of *ScPPZ1* in the *S. cerevisiae* *slt2* mutant is known to attenuate these defects. As shown in Fig. 3(b), high-copy-number expression of *CaPPZ1* was able to markedly improve the tolerance of the *slt2* mutant when cells were grown in the presence of caffeine or CFW. Therefore, CaPpz1 is able to reproduce the genetic interaction with the *SLT2* MAP kinase.

Since the primary structures of CaPpz1 and Pzh1 proteins are similar to one another (Figs 1a and S1), the functional competence of CaPpz1-3 was analysed in the genetically distant fission yeast. The *C. albicans* phosphatase was expressed from a pREP41 plasmid in *Sch. pombe* *pzh1* deletion mutant cells (Fig. 4). The *C. albicans* protein rescued, at least in part, the salt and caffeine sensitivity of the mutant. Similar results were obtained at three different LiCl, NaCl, KCl and caffeine concentrations (data not shown). Thus, CaPpz1 can perform similar but not completely identical physiological functions to its *S. cerevisiae* and *Sch. pombe* counterparts.

Dissection of the functions of CaPpz1 in *C. albicans* by gene inactivation

In order to identify the physiological functions of CaPpz1 in *C. albicans*, we disrupted both copies of the corresponding gene in the diploid organism. The efficiency and specificity of the gene deletion were demonstrated by PCR, RT-PCR and Southern blot experiments (Fig. S3a–c). The mutant *C. albicans* did not exhibit any easily recognizable phenotype when cultivated in YPD medium. Therefore, the viability of the Δ *CaPPZ1* strain was compared with that of the parental SN87 strain under different stress conditions. There was almost no difference between the two strains in the presence of NaCl; however, the disrupted strain was more tolerant to LiCl, spermine and hygromycin B, and was more sensitive to KCl, caffeine, CFW and Congo red

Fig. 3. Expression of *C. albicans* CaPpz1 protein in *S. cerevisiae*. (a) Wild-type BY4741 (*ScPPZ1*) strain and its isogenic derivative BY4741 *Scppz1* were transformed with the indicated plasmids. Cells (two dilutions of approx. 3×10^3 and 3×10^2 cells) were spotted on YPD plates containing the indicated concentrations of LiCl, NaCl or caffeine and growth was monitored after 48 h of incubation at 28 °C (except for cells growing in 15 mM caffeine, which were scored for growth after 60 h). (b) Strains BY4741 (*SLT2*) and BY4741 *slt2* were transformed with the indicated plasmids and cultures were spotted on YPD plates containing the indicated concentrations of the drugs. Growth was monitored after 72 h, while the plate containing 20 mM caffeine was incubated for 96 h.

treatments (Fig. 5). In order to verify that these phenotypes were indeed related to the absence of the target gene, we reintegrated one functional *CaPPZ1* copy into its original locus. Both of the heterozygous transformants that carried a single copy of *CaPPZ1* in a different genetic background behaved more similarly to the parental SN87 strain under selected stress conditions (Fig. 5e–h). Reintegration partially compensated sensitivity to CFW and Congo red, and complemented spermine as well as hygromycin B tolerance. Together, our biochemical and genetic data suggest that CaPpz1 has functions similar but not identical to the well-characterized ScPpz1.

Beside the stress treatments, we also analysed germ tube formation, a biological process that is thought to be related to the virulence of this pathogenic fungus. The germination frequency was determined for the parental as well as for the homozygous and heterozygous mutant strains (Table 2). At 30 min after the addition of sheep serum a significant delay was detected in the germ tube formation of the *C. albicans* that had no functional *CaPPZ1*. The difference between the null mutant and the other strains gradually diminished with the time of incubation and disappeared after 90 min. CFW staining of the cells demonstrated that all of the tested strains produced regular hyphae (Sudbery *et al.*, 2004) during the incubation period.

The virulence of the disrupted strain was directly tested in BALB/c mice (Fig. 6). It was found that the *cappz1* mutant was a less effective pathogen than the parental SN87 strain,

while both of the heterozygous strains were more virulent than the null mutant. Kaplan–Meyer analysis of the combined survival data proved that the virulence of the four strains analysed was significantly different ($P=0.0115$). Pairwise comparisons revealed that the SN87 and the *CaPPZ1-LEU2* strains were significantly more pathogenic than the null mutant ($P=0.078$ and 0.0343 , respectively), although the difference between the *CaPPZ1-HIS1* and *cappz1* cells did not reach the level of significance. In fact, *CaPPZ1-LEU2* was more pathogenic than SN87, probably because it grew somewhat faster than the other strains. The reversion to the more pathogenic phenotype by gene reintegration suggests that the disruption of the *CaPPZ1* was responsible for the reduced virulence.

DISCUSSION

The heterogeneity of the *CaPPZ1* gene is an inherent property of *C. albicans* (Kovács *et al.*, 2010) that has some impact on structure–function investigations. The natural alleles used in the present study (*CaPPZ1-1* and *CaPPZ1-3*) encode the same amino acid sequence in the catalytic domain (Fig. S1) and are expected to have the same catalytic properties. In addition, due to the different codon usage of *C. albicans* with respect to other organisms (Ohama *et al.*, 1993), five CUG triplets were translated as Leu instead of Ser in *S. cerevisiae*, *Sch. pombe* and *E. coli* (Fig. S1). Only one of these (L452) is in the catalytic

Fig. 4. Expression of *C. albicans* CaPpz1 protein in *Sch. pombe*. The relative growth rate of the *pzh1*⁺ control cells transformed with the empty pREP1 vector (black bars), the *pzh1*-disrupted LB2 cells transformed with the empty pREP1 vector (white bars), and with CaPPZ1-pREP41 (grey bars) was compared in the presence of (a) 8 mM LiCl, (b) 150 mM NaCl, (c) 300 mM KCl and (d) 5 mM caffeine. The relative growth rate of the cells was determined after 24 h incubation in liquid cultures, and was expressed as a percentage with respect to the growth of the same strain without additions. The means and SDs of nine experiments are shown. The significances of the changes relative to the LB2 control cells are given according to *t* test analysis (triple asterisks indicate $P < 0.0005$).

domain, where it is found in an external loop (Fig. 1b). We suppose that these surplus mutations (that are present in all of the recombinant enzymes tested) have no significant effect on enzyme activity. We demonstrated in biochemical assays that, in agreement with the structural conservation of its catalytic domain, the CaPpz1 protein exhibits phosphatase activity. This activity was inhibited by the ScHal3 protein, which is a specific inhibitor of PPZ phosphatases and, at the same time, one of the putative subunits of phosphopantothenoylcysteine decarboxylase (Ruiz *et al.*, 2009). It should be noted that in *C. albicans* there are two distantly related Hal3 orthologues (orf19.7378 and orf19.3260). Interestingly, in orf19.7378, all of the known amino acid residues required for Ppz1 binding and phosphatase inhibition (Muñoz *et al.*, 2004) have been conserved. Therefore, it is likely that orf19.7378 encodes the inhibitory component of the *C. albicans* Ppz1/Hal3 system.

By inactivating the protein via the R262L mutation we demonstrated the validity of the phosphatase assay. According to our structural model, the conserved R262 residue is essential for activity because it coordinates the metal ions which are indispensable for the catalytic reaction (Fig. 1b). After proving the principle we investigated the effects of allele-specific amino acid polymorphisms on enzymic activity (Figs 1b and 2). Despite its proximity to the essential R262, the D261N mutation, characteristic of the CaPPZ1-4 allele of the *C. albicans* WO-1 strain, had only a moderate effect. Clearly, N fits well in place of D and the loss of a negative charge does not have a dramatic effect, as the D side chain is pointing outwards from the active site.

Fig. 5. Comparison of relative growth of the parental SN87 (●), homozygous null mutant *cappz1* (△), as well as the heterozygous *CaPPZ1-LEU2* (■) and *CaPPZ1-HIS1* (◇) reintegrand *C. albicans* strains under various stress conditions. The effects of (a) LiCl, (b) NaCl, (c) KCl, (d) caffeine, (e) CFW, (f) Congo red (CR), (g) spermine and (h) hygromycin B (Hyg. B) were tested after 18 h incubation at 37 °C. The means ± SDs of three independent experiments are shown.

Table 2. Role of CaPpz1 in germination of *C. albicans*

The proportion of cells producing germ tubes in sheep serum is given as a percentage of the total number of cells. The means $\pm$ SDs of three independent experiments are shown. The significance of the changes relative to SN87 were calculated by *t* test (**, $P < 0.01$; ***, $P < 0.001$).

Strain	Incubation time		
	30 min	60 min	90 min
SN87	23.5 $\pm$ 3.1	63.3 $\pm$ 4.4	96.1 $\pm$ 3.2
<i>cappz1</i>	6.0 $\pm$ 2.1***	42.6 $\pm$ 3.6**	95.7 $\pm$ 2.5
<i>CaPPZ1-LEU2</i>	25.6 $\pm$ 6.2	64.0 $\pm$ 2.1	97.9 $\pm$ 1.2
<i>CaPPZ1-HIS1</i>	23.8 $\pm$ 5.4	59.2 $\pm$ 5.6	97.2 $\pm$ 1.7

Unexpectedly, the G333E mutation (which is present in the *CaPPZ1-2* allele) significantly activated the phosphatase. Homology modelling of the catalytic domain predicts that G333 is at the surface of the protein and is not thought to influence the structure of the catalytic cleft. The C337R polymorphism (that is also found in the *CaPPZ1-2* allele) caused the inactivation of the enzyme. C337 is in the middle of the central β -sheet and may have important functions in the stability of the catalytic domain. Its replacement with a bulky, charged residue may interfere with the proper folding of the tertiary structure. Indeed, the C337R mutation eliminated the activating effect of the G333E exchange and resulted in an inactive double mutant (Fig. 2). Previously we identified these two amino acid exchanges together in the heterozygous ATCC 10231 strain that harbours alleles *CaPPZ1-2* and *CaPPZ1-3* (Kovács *et al.*, 2010). In addition, we isolated several clinical samples that were homozygous for the *CaPPZ1-2* allele according to RFLP of the PCR fragment that encompasses the hypervariable 3' non-coding region (Kovács *et al.*, 2010). We tested one of them (number 10934) under several stress conditions and found that it did not show the characteristic phenotypes of the null mutant. In order to determine the molecular basis of this unexpected behaviour we amplified and sequenced the *CaPPZ1* gene

from the clinical sample. We found that both copies of the gene exhibited a DNA sequence characteristic of the *CaPPZ1-2* allele, but in both alleles the triplets GAG and TGT, encoding E333 and R337, respectively, reverted to GGG and CGU, encoding G333 and C337, that is the amino acids of the active CaPpz1 isoenzymes! Our findings indicate that the destabilizing C337R replacement is not tolerated in a homozygous organism, and indirectly support the notion that the phosphatase activity is important for the survival of this pathogenic fungus in its natural habitat.

The functions of CaPpz1 were first tested in complementation experiments. If CaPpz1 was expressed in *S. cerevisiae* *ppz1* cells it partially complemented the salt sensitivity and rescued the caffeine sensitivity of the mutant. In addition, the caffeine and CFW sensitivities of a strain lacking the Slr2 MAP kinase were also alleviated. The observation that the presence of the CaPpz1 protein can, at least in part, reverse the effects of the absence of ScPpz1 or mimic its overexpression with respect to both cation homeostasis and cell wall integrity is coherent, since it has been shown that these two phenotypes are interrelated (Merchan *et al.*, 2004). The partial complementation of the *Sch. pombe* *pzh1* deletion mutant by the expression of *CaPPZ1* revealed that, although the heterologous protein can replace the authentic *Sch. pombe* enzyme, it is clearly less efficient. Similar conclusions were previously drawn from the functional study of *N. crassa* Pz1-1 (Vissi *et al.*, 2001).

The functions of CaPpz1 were directly assessed by the disruption of the gene in *C. albicans* strain SN87. In accordance with earlier reports (Hanaoka *et al.*, 2008; Xu *et al.*, 2007), the deletion of the gene was not lethal; thus, it has no essential role under common cultivation conditions. However, when the *C. albicans* cells were challenged by various stress treatments, the lack of the phosphatase became detectable. We also demonstrated that the re-integration of *CaPPZ1* into the deletion mutant alleviated the typical mutant phenotypes. We found that the null mutant was tolerant to LiCl, and was sensitive to KCl. According to these properties it behaves like the *S. cerevisiae* *ppz1* mutant (Posas *et al.*, 1995; Ruiz *et al.*, 2003; Yenush *et al.*, 2002). However, there was a clear-cut difference between the two mutants when sensitivity to NaCl was tested. It is known that *S. cerevisiae* *ppz1* tolerates saline treatment, but we found no significant differences between the parental and mutated *C. albicans* strains in the presence of NaCl. It was previously suggested that the tolerance to NaCl (and LiCl) of the *S. cerevisiae* mutant can be explained by the overexpression of the *ENA1* sodium transporter (Posas *et al.*, 1995; Ruiz *et al.*, 2003). Sequence comparisons revealed that in *C. albicans* the orf19.6070 protein is an *ENA1* orthologue; however, its function and regulation are not known at the moment. On the other hand, in the highly salt-tolerant fungus *D. hansenii*, the hypertolerance of toxic cations caused by deletion of *DhPPZ1* is due not to the increased expression of the *DhENA1* Na⁺-ATPase, but to increased expression of the *DhEHA1* Na⁺/H⁺ antiporter, which appears as the likely

Fig. 6. Role of CaPpz1 in the virulence of *C. albicans*. The survival of BALB/c mice was tested after the administration of $1.8\text{--}2.0 \times 10^5$ SN87 (●), *cappz1* (△), *CaPPZ1-LEU2* (■) and *CaPPZ1-HIS1* (◇) *C. albicans* cells. Representative results of one out of three experiments are shown.

functional target for the phosphatase in this organism (Minhas *et al.*, 2012). Consequently, different fungi may utilize distinct molecular mechanisms to elicit similar physiological responses.

As the *cappz1* mutant was sensitive to cell wall-damaging agents such as caffeine, CFW and Congo red, we suggest that CaPpz1 interacts with the CWI pathway in a similar manner to its *S. cerevisiae* counterpart. Furthermore, the *C. albicans* mutant was tolerant to toxic cations (spermine and hygromycin B), indicating that the absence of CaPpz1 resulted in a depolarization of cell membrane potential that decreased uptake of the toxic agents, as previously found for budding yeast *ppz* mutants (Yenush *et al.*, 2002). These experiments underline again the similarity between the *C. albicans* and *S. cerevisiae* PPZ phosphatases. In conclusion, our data collectively indicate that PPZ phosphatases have similar but not identical functions in different yeasts.

In addition, we identified a novel function for this enzyme that operates in the filamentous form of *C. albicans*. We observed that after serum stimulation the onset of germ tube growth was significantly reduced in the *cappz1* mutant relative to the parental SN87 strain or to the heterozygous mutants that had one reintegrated *CaPPZ1* copy. The size and the shape of the tubes were similar in all of the strains tested (Fig. S4), suggesting that the phosphatase affected the initiation and not the rate of hyphal outgrowth in liquid cultures. Previously, we reported that two phosphatase inhibitors, cantharidin and calyculin A, hindered hyphal growth in the filamentous fungus *N. crassa* (Yatzkan *et al.*, 1998). Genetic evidence indicated that protein phosphatase 2A (*pph-1*) was involved in the process; however, the role of additional phosphatases was not excluded. Now we have found that 220 µM cantharidin completely blocked, while 250 nM calyculin A significantly reduced, the phosphatase activity of recombinant CaPpz1 (data not shown). The genetic and biochemical data together support the hypothesis that (besides PP2A) CaPpz1 can also contribute to the regulation of hypha formation. Since filament formation is considered to be a critical element of *C. albicans* pathogenesis, we tested the virulence of the mutated *C. albicans* in BALB/c mice, and we found that the *cappz1* mutant was somewhat less virulent than its parental strain. This result differs from those reported in another paper (Hanaoka *et al.*, 2008), which indicated that a *cappz1* strain displays wild-type virulence in a silkworm infection model. It is conceivable that the difference in the model employed could explain the contradictory results. The rescue of the less virulent phenotype by the reintegration of a single copy of the *CaPPZ1* gene indicates that the phosphatase gene is involved in the virulence of the pathogen. The finding that the absence of CaPpz1 function moderately reduces virulence in a mammalian model may be of importance, as Ppz phosphatases are fungus-specific. Therefore, CaPpz1 could be considered as a possible target for antifungal treatments, and CaPpz1-specific inhibitors may act as antifungal drugs.

ACKNOWLEDGEMENTS

Thanks are due to Dr Susanne Noble (University of California at San Francisco) for the SN87 *C. albicans* strain. We thank Dr Jürgen Wendland (Carlsberg Laboratory, Denmark) for providing the pFA-CdHIS1 and pFA-CmLeu2 plasmids, and Dr Jesús Pla (Universidad Complutense de Madrid) for the pSFS2A plasmid. The authors are grateful to Mr Imre Pócsi, Mrs Ágota Kelemenné Szántó, Mrs Andrea Tankáné Farkas, Ms Enikő Boros and Ms Montse Robledo for their technical assistance. The help of Dr Rudolf Gesztelyi (Department of Pharmacology, University of Debrecen) in the statistical calculations is acknowledged. We are grateful to Dr Ida Miklós and Mr László Papp (Department of Genetics, University of Debrecen) for the microscopic images. This work was supported by a Hungarian Research Fund (OTKA K 68765) grant and the TÁMOP 4.2.1/B-09/1/KONV-2010-0007 project to V.D., by the Ministry of Science and Innovation, Spain and FEDER grants BFU2008-04188-C03-01 and BFU2011-30197-C3-01 to J.A., and by bilateral Hungarian-Spanish research grants HH2008-0026 and ES-22/2008 to J.A. and V.D. J.A. is the recipient of Generalitat de Catalunya 'Ajut 2009SGR-1091' and an ICREA Academia Award.

REFERENCES

- Ariño, J. (2002). Novel protein phosphatases in yeast. *Eur J Biochem* **269**, 1072–1077.
- Balcells, L., Gómez, N., Casamayor, A., Clotet, J. & Ariño, J. (1997). Regulation of salt tolerance in fission yeast by a protein-phosphatase-Z-like Ser/Thr protein phosphatase. *Eur J Biochem* **250**, 476–483.
- Balcells, L., Calero, F., Gómez, N., Ramos, J. & Ariño, J. (1999). The *Schizosaccharomyces pombe* Pzh1 protein phosphatase regulates Na⁺ ion influx in a Trk1-independent fashion. *Eur J Biochem* **260**, 31–37.
- Basi, G., Schmid, E. & Maundrell, K. (1993). TATA box mutations in the *Schizosaccharomyces pombe nmt1* promoter affect transcription efficiency but not the transcription start point or thiamine repressibility. *Gene* **123**, 131–136.
- Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* **72**, 248–254.
- Clotet, J., Posas, F., de Nadal, E. & Ariño, J. (1996). The NH₂-terminal extension of protein phosphatase PP2A has an essential functional role. *J Biol Chem* **271**, 26349–26355.
- Clotet, J., Garí, E., Aldea, M. & Ariño, J. (1999). The yeast Ser/Thr phosphatases Sit4 and Ppz1 play opposite roles in regulation of the cell cycle. *Mol Cell Biol* **19**, 2408–2415.
- de Nadal, E., Clotet, J., Posas, F., Serrano, R., Gomez, N. & Ariño, J. (1998). The yeast halotolerance determinant Hal3p is an inhibitory subunit of the Ppz1p Ser/Thr protein phosphatase. *Proc Natl Acad Sci U S A* **95**, 7357–7362.
- García-Gimeno, M. A., Muñoz, I., Ariño, J. & Sanz, P. (2003). Molecular characterization of Ypi1, a novel *Saccharomyces cerevisiae* type 1 protein phosphatase inhibitor. *J Biol Chem* **278**, 47744–47752.
- Gola, S., Martin, R., Walther, A., Dünkler, A. & Wendland, J. (2003). New modules for PCR-based gene targeting in *Candida albicans*: rapid and efficient gene targeting using 100 bp of flanking homology region. *Yeast* **20**, 1339–1347.
- Goldberg, J., Huang, H. B., Kwon, Y. G., Greengard, P., Nairn, A. C. & Kuriyan, J. (1995). Three-dimensional structure of the catalytic subunit of protein serine/threonine phosphatase-1. *Nature* **376**, 745–753.

- Gutz, H., Heslot, H., Leupold, U. & Loprieno, N. (1974). *Schizosaccharomyces pombe*. In *Handbook of Genetics*, vol. 1, pp. 395–446. Edited by R. C. King. New York: Plenum.
- Gyetvai, Á., Emri, T., Fekete, A., Varga, Z., Gazdag, Z., Pesti, M., Belágyi, J., Emödy, L., Pócsi, I. & Lenkey, B. (2007). High-dose methylprednisolone influences the physiology and virulence of *Candida albicans* ambiguously and enhances the candidacidal activity of the polyene antibiotic amphotericin B and the superoxide-generating agent menadione. *FEMS Yeast Res* 7, 265–275.
- Hanaoka, N., Takano, Y., Shibuya, K., Fugo, H., Uehara, Y. & Niimi, M. (2008). Identification of the putative protein phosphatase gene *PTC1* as a virulence-related gene using a silkworm model of *Candida albicans* infection. *Eukaryot Cell* 7, 1640–1648.
- Ito, H., Fukuda, Y., Murata, K. & Kimura, A. (1983). Transformation of intact yeast cells treated with alkali cations. *J Bacteriol* 153, 163–168.
- Kovács, L., Farkas, I., Majoros, L., Miskei, M., Pócsi, I. & Dombrádi, V. (2010). The polymorphism of protein phosphatase Z1 gene in *Candida albicans*. *J Basic Microbiol* 50 (Suppl. 1), S74–S82.
- Lee, S. B., Milgroom, M. G. & Taylor, J. W. (1988). A rapid, high yield mini-prep method for isolation of total genomic DNA from fungi. *Fungal Genet Newsl* 35, 23.
- Lee, K. S., Hines, L. K. & Levin, D. E. (1993). A pair of functionally redundant yeast genes (*PPZ1* and *PPZ2*) encoding type 1-related protein phosphatases function within the *PKC1*-mediated pathway. *Mol Cell Biol* 13, 5843–5853.
- Levin, D. E. (2005). Cell wall integrity signaling in *Saccharomyces cerevisiae*. *Microbiol Mol Biol Rev* 69, 262–291.
- Maundrell, K. (1993). Thiamine-repressible expression vectors pREP and pRIP for fission yeast. *Gene* 123, 127–130.
- Merchan, S., Bernal, D., Serrano, R. & Yenush, L. (2004). Response of the *Saccharomyces cerevisiae* Mpk1 mitogen-activated protein kinase pathway to increases in internal turgor pressure caused by loss of Ppz protein phosphatases. *Eukaryot Cell* 3, 100–107.
- Minhas, A., Sharma, A., Kaur, H., Rawal, Y., Ganesan, K. & Mondal, A. K. (2012). A conserved Ser/Arg rich motif in PPZ orthologs from fungi is important for its role in cation tolerance. *J Biol Chem* 287, 7301–7312.
- Muñoz, I., Ruiz, A., Marquina, M., Barceló, A., Albert, A. & Ariño, J. (2004). Functional characterization of the yeast Ppz1 phosphatase inhibitory subunit Hal3: a mutagenesis study. *J Biol Chem* 279, 42619–42627.
- Noble, S. M. & Johnson, A. D. (2005). Strains and strategies for large-scale gene deletion studies of the diploid human fungal pathogen *Candida albicans*. *Eukaryot Cell* 4, 298–309.
- Ohama, T., Suzuki, T., Mori, M., Osawa, S., Ueda, T., Watanabe, K. & Nakase, T. (1993). Non-universal decoding of the leucine codon CUG in several *Candida* species. *Nucleic Acids Res* 21, 4039–4045.
- Posas, F., Casamayor, A., Morral, N. & Ariño, J. (1992). Molecular cloning and analysis of a yeast protein phosphatase with an unusual amino-terminal region. *J Biol Chem* 267, 11734–11740.
- Posas, F., Casamayor, A. & Ariño, J. (1993). The PPZ protein phosphatases are involved in the maintenance of osmotic stability of yeast cells. *FEBS Lett* 318, 282–286.
- Posas, F., Camps, M. & Ariño, J. (1995). The PPZ protein phosphatases are important determinants of salt tolerance in yeast cells. *J Biol Chem* 270, 13036–13041.
- Reuss, O., Vik, A., Kolter, R. & Morschhauser, J. (2004). The *SAT1* flipper, an optimized tool for gene disruption in *Candida albicans*. *Gene* 341, 119–127.
- Ruiz, A., Yenush, L. & Ariño, J. (2003). Regulation of *ENA1* Na⁺-ATPase gene expression by the Ppz1 protein phosphatase is mediated by the calcineurin pathway. *Eukaryot Cell* 2, 937–948.
- Ruiz, A., Muñoz, I., Serrano, R., González, A., Simón, E. & Ariño, J. (2004). Functional characterization of the *Saccharomyces cerevisiae* *VHS3* gene: a regulatory subunit of the Ppz1 protein phosphatase with novel, phosphatase-unrelated functions. *J Biol Chem* 279, 34421–34430.
- Ruiz, A., González, A., Muñoz, I., Serrano, R., Abrie, J. A., Strauss, E. & Ariño, J. (2009). Moonlighting proteins Hal3 and Vhs3 form a heteromeric PPCDC with Ykl088w in yeast CoA biosynthesis. *Nat Chem Biol* 5, 920–928.
- Sali, A. & Blundell, T. L. (1993). Comparative protein modelling by satisfaction of spatial restraints. *J Mol Biol* 234, 779–815.
- Schaub, Y., Dünkler, A., Walther, A. & Wendland, J. (2006). New pFA-cassettes for PCR-based gene manipulation in *Candida albicans*. *J Basic Microbiol* 46, 416–429.
- Sudbery, P., Gow, N. & Berman, J. (2004). The distinct morphogenic states of *Candida albicans*. *Trends Microbiol* 12, 317–324.
- Szőör, B., Fehér, Z., Zeke, T., Gergely, P., Yatzkan, E., Yarden, O. & Dombrádi, V. (1998). *pzl-1* encodes a novel protein phosphatase-Z-like Ser/Thr protein phosphatase in *Neurospora crassa*. *Biochim Biophys Acta* 1388, 260–266.
- Thompson, J. D., Higgins, D. G. & Gibson, T. J. (1994). CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res* 22, 4673–4680.
- Vissi, E., Clotet, J., de Nadal, E., Barceló, A., Bakó, E., Gergely, P., Dombrádi, V. & Ariño, J. (2001). Functional analysis of the *Neurospora crassa* PZL-1 protein phosphatase by expression in budding and fission yeast. *Yeast* 18, 115–124.
- Winston, F., Dollard, C. & Ricupero-Hovasse, S. L. (1995). Construction of a set of convenient *Saccharomyces cerevisiae* strains that are isogenic to S288C. *Yeast* 11, 53–55.
- Winzeler, E. A., Shoemaker, D. D., Astromoff, A., Liang, H., Anderson, K., Andre, B., Bangham, R., Benito, R., Boeke, J. D. & other authors (1999). Functional characterization of the *S. cerevisiae* genome by gene deletion and parallel analysis. *Science* 285, 901–906.
- Xu, D., Jiang, B., Ketela, T., Lemieux, S., Veillette, K., Martel, N., Davison, J., Sillaots, S., Trosok, S. & other authors (2007). Genome-wide fitness test and mechanism-of-action studies of inhibitory compounds in *Candida albicans*. *PLoS Pathog* 3, e92.
- Yatzkan, E., Szőör, B., Fehér, Z., Dombrádi, V. & Yarden, O. (1998). Protein phosphatase 2A is involved in hyphal growth of *Neurospora crassa*. *Mol Gen Genet* 259, 523–531.
- Yenush, L., Mulet, J. M., Ariño, J. & Serrano, R. (2002). The Ppz protein phosphatases are key regulators of K⁺ and pH homeostasis: implications for salt tolerance, cell wall integrity and cell cycle progression. *EMBO J* 21, 920–929.

Edited by: J. Pla