The Aerobic Pseudomonads: a Taxonomic Study

By R. Y. STANIER, N. J. PALLERONI AND M. DOUDOROFF

*Department of Bacteriology and Immunology, University of
California, Berkeley, California, U.S.A.*

Dedicated to L. E. DEN DOOREN DE JONG, who laid the foundations
upon which this work was built

(Received 2 October 1965)

CONTENTS

	PAGE		PAGE
Summary	160	The constitutive synthesis of arginine dihydrolase and its relation to the utilization of arginine as a carbon source	183
Introduction	160	Certain aromatic ring cleavage mechanisms	185
Origin and plan of the present study	164	Some universally negative nutritional characters of the strains examined	186
The choice of biological material	167	Nutritional versatility of the strains examined	186
Methods	168	The fluorescent group	188
Standard media	168	Origins of the strains	188
Screening of organic compounds as sources of carbon and energy for growth	168	General group characters	194
Tests for chemolithotrophic growth with H ₂	171	Internal subdivision of the fluorescent group	198
Tests for denitrification	171	Flagellar number	201
Determination of pigment production	172	Extracellular enzyme production	201
Biochemical methods	173	Temperature relations	203
Screening for ring fission mechanisms	173	Denitrification	203
Assay of the arginine dihydrolase system	174	Phenazine pigment production	203
Oxidase test	174	Nutritional patterns	204
Determination of cytochrome difference spectra	174	<i>Pseudomonas aeruginosa</i>	208
Production of extracellular enzymes	174	<i>Pseudomonas fluorescens</i>	208
Cytological observations	175	Biotype A	210
Temperature relationships	176	Biotype B	210
Nitrogen sources and growth factor requirements	176	Biotype C	213
Determination of base composition of DNA	176	Biotype D	216
Miscellaneous tests	177	Biotype E	218
Biological materials	177	Biotype F	221
Results	178	Biotype G	223
Primary differentiation of species and species groups	178	<i>Pseudomonas putida</i>	225
Taxonomic significance of poly- β -hydroxybutyrate metabolism in aerobic pseudomonads	179	Biotype A	225
Significance of denitrifying ability	182	Biotype B	226
The oxidase reaction and its relation to the cytochrome system	182	Unclassified strains	227
		Neotype strains in the fluorescent group	227
		Nomenclatural problems in the fluorescent group	229
		The acidovorans group	236
		Origins of the strains	236
		General group characters	237

CONTENTS (*cont.*)

	PAGE		PAGE
Results (<i>cont.</i>)		Ecological considerations and enrichment	253
The acidovorans group (<i>cont.</i>)			
Differentiation of two species:	241	<i>Pseudomonas stutzeri</i>	254
<i>P. acidovorans</i> and <i>P. testosteroni</i>		Origins of the strains	254
Nomenclatural problems and designation of type strains	241	General properties	254
Ecological considerations and enrichment	242	DNA base composition and internal subdivision of <i>P. stutzeri</i>	258
Properties of unclassified strains	243	<i>Pseudomonas maltophilia</i>	259
The alcaligenes group	243	Origins of the strains	259
Origins of the strains	244	General properties	260
General group characters	244	Ecological considerations	262
Differentiation of two species:	247	Discussion	263
<i>P. alcaligenes</i> and <i>P. pseudoalcaligenes</i> sp.nov.		Nutritional spectra: their taxonomic utility, physiological interpretation, and limitations	263
<i>Pseudomonas multivorans</i> sp.nov.	247	Prospective developments: towards a biochemical taxonomy of the aerobic pseudomonads	264
Origins of the strains	248		
General properties	249	Acknowledgements	267
Phenotypic comparison between <i>P. multivorans</i> and other species of aerobic pseudomonads	252	References	268
Proposal of type strain	253		

SUMMARY

A collection of 267 strains, representing many of the principal biotypes among aerobic pseudomonads, has been subjected to detailed study, with particular emphasis on biochemical, physiological and nutritional characters. A total of 146 different organic compounds were tested for their ability to serve as sources of carbon and energy. Other characters that were studied included: production of extracellular hydrolases; nitrogen sources and growth factor requirements; H_2 -chemolithotrophy; denitrifying ability; pigment production; ability to accumulate poly- β -hydroxybutyrate as a cellular reserve material; biochemical mechanisms of aromatic ring cleavage; and nature of the aerobic electron transport system. The resultant data have revealed many hitherto unrecognized characters of taxonomic significance. As a consequence, it has become possible to recognize among the biotypes examined a limited number of species which can be readily and clearly distinguished from one another by multiple, unrelated phenotypic differences.

INTRODUCTION

The aerobic pseudomonads are an important assemblage of eubacteria which has so far proved resistant to satisfactory taxonomic analysis. We shall describe in this paper a methodology that provides taxonomically significant information about these organisms, as shown by the fact that it permits the division of a large collection of strains into a relatively small number of clusters, distinguishable from one another by multiple, unrelated phenotypic differences. Most of these clusters appear to merit recognition as bacterial species.

We offer the following definition of the aerobic pseudomonads:

Unicellular rods, with the long axis straight or curved, but not helical. Motile by means of one or more polar flagella. Gram-negative. Do not form spores, stalks, or

sheaths. The energy-yielding metabolism is respiratory, never fermentative or photosynthetic. All use molecular oxygen as a terminal oxidant; and some can use denitrification as an alternative, anaerobic respiratory mechanism. All are chemo-organotrophs; some are facultative chemolithotrophs that use H_2 as an energy source.

One additional group property which appears significant, but which cannot yet be fully evaluated, is the base content of the deoxyribonucleic acid. In the species that we have examined, the DNA contains between 58 and 69 moles % of guanine and cytosine (Mandel, 1966). However, it seems unwise to set absolute limits for the base content of the DNA until other biotypes, which possibly deserve to be included, have been subjected to analysis.

This definition excludes many other groups of Gram-negative, polarly flagellated bacteria. The chemoheterotrophic spirilla, the caulobacters, and the *Leptothrix-Sphaerotilus* group are excluded on structural grounds. The specifications with respect to energy-yielding metabolism exclude all pseudomonads and vibrios capable of fermentative metabolism (*Aeromonas*, *Zymomonas*, *Photobacterium*, *Vibrio* in the strict sense); *Desulfovibrio*; and photosynthetic bacteria. The statement concerning chemolithotrophic nutrition excludes chemolithotrophic pseudomonads such as *Nitrosomonas*, *Ferrobacillus* and *Thiobacillus*. The only Gram-negative, polarly flagellated groups which it seems desirable to separate from the aerobic pseudomonads, but which are not rigorously excluded by the proposed definition, are the small-celled parasitic vibrioid bacteria for which the genera *Campylobacter* (Sebald & Véron, 1963) and *Bdellovibrio* (Stolp & Starr, 1963) have been recently proposed. Since the DNA of *Campylobacter* spp. contains about 40 moles % of guanine and cytosine (Sebald & Véron, 1963), a separation may prove feasible on this character alone.

The aerobic pseudomonads constitute a large and fairly diverse array of bacteria. There is general agreement that some of them belong to the genus *Pseudomonas*. However, the circumscription of this genus at present is far from clear, and many of the bacteria that we include among the aerobic pseudomonads are now most commonly classified in other genera: *Xanthomonas*, *Methanomonas*, *Hydrogenomonas*, *Comamonas*, *Vibrio sensu lato*, *Acetomonas* (*Gluconobacter*), and possibly also *Alginomonas*, *Cellulomonas* and *Cellvibrio*. We doubt the usefulness of this generic hypertrophy, and would assign all these organisms provisionally to the single genus *Pseudomonas*, using the group definition of aerobic pseudomonads already presented as a generic definition. Eventually, generic subdivision may prove useful; but, if and when it is undertaken, it should be based on criteria more substantial, and much more thoroughly evaluated, than those which now purportedly separate such genera as *Xanthomonas*, *Hydrogenomonas* and *Comamonas* from *Pseudomonas*.

It is exceedingly difficult to give a satisfactory account of the nature and extent of the aerobic pseudomonads, because the existing descriptions of the constituent subgroups are so fragmentary that the boundaries between them are in many cases by no means clearly defined; some subgroups are recognized primarily on pigmentation, others on pathogenicity, other on physiological attributes that have been specifically selected by the methods used to isolate them from nature. With due recognition of the hazardous nature of the enterprise, it nevertheless seems desirable to attempt an initial survey of the group.

The most extensively studied subgroup of aerobic pseudomonads consists of the

fluorescent pseudomonads, primarily characterized by their ability to produce water-soluble, yellow-green fluorescent pigments of unknown chemical nature. Simple fluorescent pseudomonads are common inhabitants of soil and water, and were first described by Flügge (1886). Flügge recognized two biotypes, distinguishable on the character of gelatin liquefaction, which now bear the names of *Pseudomonas fluorescens* (liquefying) and *P. putida* (non-liquefying). Since then, many other species of simple fluorescent pseudomonads from soil and water have been named on the basis of fragmentary descriptions; their differences from the two classical species, if any, are not clear. As first shown by den Dooren de Jong (1926), simple fluorescent pseudomonads are nutritionally versatile, and can use a very wide range of simple organic compounds as sole sources of carbon and energy.

The bacteria pathogenic for higher plants include many simple fluorescent pseudomonads, and numerous species have been defined largely or solely on the basis of their ability to infect specific plant hosts. This proliferation of specific names for phytopathogenic fluorescent pseudomonads is probably not justifiable (Starr, 1959); and it is still not really clear in what respects, if any, phytopathogenic strains differ from the simple fluorescent pseudomonads commonly found in soil and water.

Several other species have been recognized by virtue of their ability to synthesize, in addition to the group-specific fluorescent pigment, a type-specific phenazine pigment. From the available descriptions, some of these phenazine producers (*Pseudomonas chlororaphis*, *P. aureofaciens*, *P. lemonnierii*) appear to resemble closely in most phenotypic respects the simple fluorescent pseudomonads of the type of *P. fluorescens*. However, *P. aeruginosa*, the type species of the genus *Pseudomonas*, has several properties in addition to the production of a specific phenazine pigment that set it apart from the rest of the group. It is the only fluorescent pseudomonad pathogenic for mammals; it has a temperature maximum higher than that of the other fluorescent pseudomonads (Haynes, 1951); and its flagellation is invariably monotrichous, while nearly all other fluorescent strains are multitrichous (Lautrop & Jessen, 1964).

The ability to produce phenazine pigments also occurs in a group of non-fluorescent pseudomonads, originally isolated from soil in Trinidad (Morris & Roberts, 1959). These bacteria accumulate poly- β -hydroxybutyrate as a cellular reserve material, a property not found in the fluorescent group (Morris & Roberts, 1959). Although this group has been studied by several workers (Forsyth, Hayward & Roberts, 1958; Sneath, 1960), it has not received a formal taxonomic designation. As we shall show, it is a very distinctive, nutritionally versatile biotype which we propose to recognize as a new species, *Pseudomonas multivorans*.

Apart from the fluorescent group and *Pseudomonas multivorans*, the aerobic pseudomonads are either non-pigmented, or produce yellow or red cellular pigments. The character of cellular pigmentation has major diagnostic importance, since the genus *Xanthomonas* was created by Dowson (1939) for certain yellow phytopathogenic pseudomonads. Starr & Stephens (1964) have recently characterized the pigments of the *Xanthomonas* group as new and unusual carotenoids. However, the production of yellow cellular pigments is also characteristic of some hydrogenomonads, and not uncommon in chemoheterotrophic pseudomonads isolated by other enrichment methods or by non-selective methods from water and soil. The relation of these pigmented pseudomonads to the xanthomonads is at present obscure.

The group of hydrogenomonads is now in practice defined by the specific enrichment method used to isolate them from natural sources: selection for ability to grow chemolithotrophically with H_2 . Most of the named species are very poorly described. The detailed nutritional study by Doudoroff (1940) on one species, *Pseudomonas saccharophila*, showed that it is a versatile chemo-organotroph, and the less detailed nutritional information about other biotypes suggests that this may be true of all hydrogenomonads. The relations of these organisms to other yellow and non-pigmented aerobic pseudomonads are unknown.

Another group (many of which produce yellow or red cellular pigments) consists of bacteria isolated by specific enrichment at the expense of methane, methanol, formate, oxalate and lower amines. These compounds do not seem to be attacked by other kinds of pseudomonads (or, indeed, by most other aerobic bacteria); hence the capacity to utilize them is probably a character of taxonomic significance. The best-characterized member of this group is *Pseudomonas methanica*, for which methane and methanol are the only organic compounds which support growth (Dworkin & Foster, 1956). Strains isolated by enrichment with methanol, oxalate and lower amines have a somewhat wider nutritional spectrum (den Dooren de Jong, 1926; Peel & Quayle, 1961; Stocks & McCleskey, 1964).

The older literature on bacterial denitrification (e.g. Burri & Stutzer, 1895; Künnemann, 1898; van Iterson, 1902) contains fragmentary descriptions of many species of aerobic pseudomonads, both fluorescent and non-fluorescent, which were enriched from nature by virtue of their denitrifying capacity. The one well-characterized non-fluorescent denitrifier is *Pseudomonas stutzeri* (van Niel & Allen, 1952).

Apart from *P. aeruginosa*, the only *Pseudomonas* species pathogenic for mammals is *P. pseudomallei*, the causative agent of melioidosis, a disease confined to warm, humid tropical regions.

Many bacteria that seem to conform to our definition of aerobic pseudomonads have been isolated from soil, fresh water and sea water by selection for the ability to grow at the expense of complex polysaccharides (cellulose, agar, chitin, alginic acid) and a number of specific, physiologically defined genera (e.g. *Cellulomonas*, *Cellvibrio*, *Alginomonas*) have been proposed for them. We shall not attempt to cite here the voluminous literature on these organisms; but it should be kept in mind that they may well constitute a large and varied group of aerobic pseudomonads, which has so far never been subjected to systematic comparative study.

Aerobic enrichments with a variety of simple organic compounds may yield non-pigmented aerobic pseudomonads in addition to, or in place of, fluorescent pseudomonads (den Dooren de Jong, 1926; Gray & Thornton, 1928; Marcus & Talalay, 1956). Several species have been named among such isolates; but only one, *Pseudomonas acidovorans* (den Dooren de Jong, 1926), has been adequately described.

Many non-pigmented aerobic pseudomonads have been directly isolated on complex media from soil, water, milk, or specimens submitted to the clinical laboratory. Among the better-characterized species in this category are *Pseudomonas maltophilia* (Hugh & Ryschenkow, 1960), *P. alcaligenes* (Hugh & Ikari, 1964) and *P. putrefaciens* (Derby & Hammer, 1931).

Lastly, it should be noted that the acetic acid bacteria with polar flagellation (genus *Acetomonas* or *Gluconobacter*) conform to our definition of aerobic pseudomonads.

Origin and plan of the present study

In attempting to characterize some of our own isolates of aerobic pseudomonads, we reached the conclusion that the diagnostic tests in current use for this group of bacteria do not yield a sufficiently broad picture of the phenotypic traits to permit a satisfactory definition of species. These tests largely originate from the array of diagnostic tests devised for the identification of coliform bacteria, a fact which explains the heavy dependence on acid production in carbohydrate broth cultures incubated under semi-aerobic conditions. Some years ago, Hugh & Leifson (1953) discovered that the basal medium used in making such tests with coliform bacteria is unreliable for detecting acid production from carbohydrates by aerobic pseudomonads; these organisms often produce little acid from carbohydrates, and are almost all powerful ammonifiers of the complex nitrogenous materials in the medium, so that cultures may become strongly alkaline even though the carbohydrate that they contain has been extensively decomposed. Hugh & Leifson accordingly modified the test medium by greatly decreasing the concentration of peptone, and their modified medium has now become the standard one for determining sugar utilization by aerobic pseudomonads. Even this modified medium does not necessarily give readily interpretable results. For example, many fluorescent pseudomonads can oxidize maltose and lactose to the corresponding bionic acids, but cannot grow at the expense of these sugars (Kluyver, de Ley & Rijven, 1951). Furthermore, it is by no means certain that all these strictly aerobic bacteria will in fact produce a detectable amount of acid during growth at the expense of carbohydrates. However, a much more general criticism of this heavy dependence on determinations of carbohydrate utilization can be made. For most nutritionally versatile aerobic pseudomonads, sugars and sugar alcohols represent at best a very small fraction of the potentially utilizable range of carbon sources, so that, if tests are confined to this chemical class, other diagnostically more valuable nutritional characters may be completely ignored. Indeed, as we shall show, several nutritionally versatile species are virtually or completely unable to use sugars and sugar alcohols as carbon sources. In recent taxonomic studies, the ability to utilize malonate and citrate have been commonly determined; these again are tests which obviously derive from work on coliform bacteria, and are not necessarily informative when applied to aerobic pseudomonads. Some workers (Lysenko, 1961; Davis & Park, 1962; Park, 1962; Iizuka & Komagata, 1963*a, b*) have tested growth on a limited range of additional compounds (aromatic substances, organic acids, alcohols), but these attempts to extend the nutritional spectrum have been at best half-hearted.

Despite superficial similarities, the aerobic pseudomonads differ very greatly from coliform bacteria in most biochemical and physiological respects. A good methodology for the study of aerobic pseudomonads is likely to be one which probes deeply their most striking and ecologically significant group character, not shared to any degree by coliform bacteria: namely, the ability to use a wide variety of organic compounds as carbon and energy sources for aerobic growth. The foundations for this approach were laid by den Dooren de Jong (1926) in a study of many strains, both fluorescent and non-fluorescent. Unfortunately, the major part of this study was published in Dutch in a thesis of limited distribution, with the result that it has

been completely ignored in all later taxonomic studies dealing with the aerobic pseudomonads. In view of its importance, a brief summary appears warranted.

The problem which den Dooren de Jong set out to study was the nature of the bacterial flora responsible for the aerobic mineralization of simple organic compounds. In order to obtain some idea of the nutritional versatility of known bacterial types, he first examined the ability of 13 different strains of aerobic bacteria to grow at the expense of several hundred different organic compounds, supplied individually as sole sources of carbon and energy in an otherwise mineral medium. The strains included three bacilli, five coliform bacteria, *Mycobacterium phlei*, a *Micrococcus* species, a *Sarcina* species, a *Spirillum* species, and *Pseudomonas fluorescens*. Only *M. phlei* and *P. fluorescens* proved capable of using a varied and extensive range of organic compounds. In the light of present knowledge, this series of experiments cannot be considered satisfactory, since some of the species that were tested (e.g. *Bacillus polymyxa* and *Proteus vulgaris*) are now known to require growth factors, not supplied in the basal medium of den Dooren de Jong. The apparently restricted range of compounds able to support their growth may have reflected the nutritional inadequacy of the basal medium.

Many of the organic compounds included in this initial survey were unable to support growth of any of the bacteria tested. Fifty of these compounds—representing a chemically very heterogeneous assortment—were then used individually as carbon and energy sources in aerobic enrichment cultures, inoculated with soil. After inoculation, the cultures were incubated without agitation at 30°. When surface growth occurred, a loopful of material was transferred to a second flask containing a medium of the same composition, and from this flask a pure culture of the predominant organism was isolated by streaking on the homologous agar medium. The procedure of isolation was rigorously selective; den Dooren de Jong reported that each enrichment medium which gave rise to bacterial growth yielded upon plating a virtually uniform population. The bacterial strains obtained from this set of enrichments were compared on the basis of morphology, colony form and pigmentation. The great majority of successful enrichments yielded pseudomonads of two types, designated as group A (15 enrichments) and group C (15 enrichments). Three enrichments yielded mycobacteria. Five, all furnished with methyl- or ethylamines, yielded representatives of a distinctive group of amine-oxidizing bacteria, which den Dooren de Jong subsequently described elsewhere in considerable detail (den Dooren de Jong, 1927). Six enrichments yielded Gram-negative rods which, despite the absence of flagella, were classified by den Dooren de Jong as *Pseudomonas* strains ('group B'). The organisms of group B were probably nutritionally non-exacting representatives of the genus *Moraxella* (Audureau, 1940; Lwoff & Audureau, 1941), to judge from the one authentic surviving strain that we have been able to examine.

Every strain isolated from this series of enrichments was subjected to very detailed nutritional study, during which ability to grow on over 200 organic compounds, furnished as sole sources of carbon and energy, was determined. This analysis revealed that the *Pseudomonas* strains of groups A and C, initially distinguished on the basis of colony appearance and pigmentation, were likewise different from one another with respect to nutritional spectra. The strains of group A were all typical fluorescent pseudomonads which did not liquefy gelatin, and were therefore

assigned by den Dooren de Jong to the species *P. putida*. The strains of group C were non-pigmented. Their nutritional homogeneity, coupled with their manifold and clear-cut nutritional differences from the group A strains, led den Dooren de Jong to create for them a new species, which he named *P. acidovorans*.

Many other strains of fluorescent pseudomonads were isolated by den Dooren de Jong by less selective methods. Eight were obtained from cultures furnished with complex organic materials (e.g. peptone, yeast autolysate, fibrin) and inoculated with soil, and 34 by the direct plating of tap water or river water on yeast agar. These additional strains were also screened, though less extensively, in order to establish their nutritional spectra.

The analysis of this great body of nutritional information reveals many facts of both ecological and taxonomic interest. In all, some 90 organic compounds could be utilized as carbon and energy sources by one or more of the strains examined. Considering only data for *Pseudomonas acidovorans* and the fluorescent group, these compounds can be divided into three categories: 'universal' substrates, attacked by nearly all strains tested; 'species-specific' substrates; and, lastly, substrates utilized by occasional strains of both groups.

The central core of universal substrates included the acids of the tricarboxylic acid cycle; lactic, pyruvic and gluconic acids; several lower fatty acids; and several amino acids, notably alanine, aspartic and glutamic acids, leucine, isoleucine, phenylalanine and tyrosine. The most unusual universal substrates were saccharic acid, mucic acid, quinic acid, uric acid and *p*-hydroxybenzoic acid.

The utilization of other compounds proved to be species-specific. The higher dicarboxylic acids (adipic to suberic) could not be attacked by any of the fluorescent pseudomonads tested, but were excellent substrates for *Pseudomonas acidovorans*. Conversely, six amines (including putrescine, spermine and benzylamine) could be attacked by nearly all fluorescent pseudomonads, but by no strain of *P. acidovorans*. The two groups also differed markedly with respect to utilization of carbohydrates. Apart from one strain of *P. acidovorans* reported to grow on glucose, strains of this species could not attack any of the five common aldopentoses and aldohexoses tested. Glucose was a universal substrate for the fluorescent pseudomonads, and many of them could attack other aldoses as well. Two amino acids served as excellent differential substrates. Arginine could be attacked by 14 out of 15 fluorescent pseudomonads, and by no strain of *P. acidovorans*; α -aminocaproic acid was used by 12 strains out of 15 of *P. acidovorans*, and by not one strain of fluorescent pseudomonad out of 57 tested. Suggestive, but less clear-cut, differences existed with respect to the utilization of aromatic substrates. Benzoic acid was a universal substrate for fluorescent pseudomonads, but was used by only a small minority of strains of *P. acidovorans*; the converse pattern occurred with respect to utilization of *m*-hydroxybenzoic acid.

This analysis accordingly permitted a distinction on many different nutritional criteria between the simple fluorescent pseudomonads and a second, nutritionally versatile group of non-pigmented pseudomonads, the bacteria for which den Dooren de Jong created *Pseudomonas acidovorans*. The total neglect which befell this work is strikingly shown by the fact that no later paper on the taxonomy of the aerobic pseudomonads so much as mentions *P. acidovorans*, even though it is without doubt one of the best-characterized species in the genus *Pseudomonas*. The frequency

of its occurrence in the enrichment cultures of den Dooren de Jong suggests that it is probably a very common soil bacterium; and, as we shall show, it has been repeatedly isolated by later workers, and described under a variety of other names.

Other interesting facts were revealed by den Dooren de Jong's data on the fluorescent pseudomonads. Of the 57 strains belonging to this group which he studied, 23 were isolated from soil, and 34 from water. The source was almost perfectly correlated with the property of gelatin liquefaction: all 34 strains of aquatic origin were liquefiers, whereas only one of the 23 strains from soil possessed this property. The ability to liquefy gelatin is the single criterion that was classically used to distinguish between *Pseudomonas fluorescens* (liquefying) and *P. putida* (non-liquefying). The data of den Dooren de Jong accordingly indicate that the distinction may be ecologically significant. Some of the compounds tested in the nutritional screening of these strains showed promise as differential substrates. Hippuric acid, creatine and α -amylamine were used by virtually all the strains from soil (*P. putida* types), and by no strain of aquatic origin (*P. fluorescens* types). In the light of this work, the later failure of Rhodes (1959) to find any conventional diagnostic criteria of use for the subdivision of a large collection of simple fluorescent pseudomonads from soil and water is interesting. Rhodes concluded that the simple fluorescent pseudomonads constitute a group in which phenotypic variation is essentially continuous; but den Dooren de Jong's study, which was not cited by Rhodes, provides evidence that her phenotypic scanning of these bacteria was of such a nature that taxonomically significant traits had been overlooked. Accordingly, the taxonomic conclusions that she reached, even though buttressed by computer analysis (Rhodes, 1961) cannot be accepted as valid.

Our primary aim in the present study has been to extend the methodology of den Dooren de Jong, by determining detailed nutritional spectra for a wider range of biotypes among the aerobic pseudomonads than those which he examined. We have coupled this with an examination of other characters which seemed to us in the light of previous work to be of potential taxonomic value.

The choice of biological material

Ideally, the definition of a bacterial species should rest on a comparative study of many strains of different origin. Such comparison is necessary to establish whether a given character is species-specific, or peculiar to one or a minority of the strains that compose the species. The minimal number of strains necessary to define a species is certainly arguable, and depends to a considerable extent on the degree of phenotypic isolation between the species in question and its closest congeners. As a rough working rule, we believe that the number of strains examined in the taxonomic analysis of a bacterial group should be at least 10–20 times as great as the anticipated number of species. In order to keep our study within reasonable bounds, we arbitrarily decided to exclude from the analysis certain subgroups. The excluded groups were: the phytopathogens; the aerobic pseudomonads able to use methane, methanol, oxalate, and lower amines; the polysaccharide decomposers; the acetic acid bacteria; and aerobic pseudomonads of marine origin. A few fluorescent strains isolated from sea water were examined, but they were not indigenous marine organisms by the criteria of MacLeod (1965).

Even with these exclusions, the collection eventually assembled consisted of

about 350 strains. In this paper, we shall discuss a total of 267 strains representing the fluorescent group (175 strains), *Pseudomonas acidovorans* and related types (26 strains), *P. multivorans* sp.nov. (19 strains), *P. stutzeri* (17 strains), *P. maltophilia* (23 strains), and *P. alcaligenes* and related types (7 strains). Characterizations of *P. pseudomallei* (26 strains) and *P. mallei* (15 strains) are being published separately (Redfearn, Palleroni & Stanier, 1966); and *P. lemoignei* sp.nov. (1 strain) has been recently characterized by our methods (Delafield *et al.* 1965). We shall draw on the data of these two publications for purposes of comparative analysis. A study of the hydrogenomonads (about 50 strains) is now in progress, and will be the subject of a later report (Davis, Doudoroff, Palleroni & Stanier, in preparation).

METHODS

Standard media

Most media were prepared from a *standard mineral base*, which contained per litre: 40 ml. of $\text{Na}_2\text{HPO}_4 + \text{KH}_2\text{PO}_4$ buffer (M; pH 6.8); 20 ml. of Hutner's vitamin-free mineral base (Cohen-Bazire, Sistrom & Stanier, 1957); and 1.0 g. of $(\text{NH}_4)_2\text{SO}_4$. This basal medium is easy to prepare, and provides all necessary minerals, including trace elements. It is heavily chelated with nitrilotriacetic acid and EDTA, and forms a copious precipitate upon autoclaving. The precipitate redissolves as the medium cools, to form a water-clear solution. Yeast extract (YE) medium was prepared by adding Difco yeast extract 5 g./l. to the standard mineral base, and yeast agar (YA) medium by the further addition of Difco agar 20 g./l. These complex media were used for the routine cultivation, purification and maintenance of stocks.

Chemically defined media were prepared by adding an appropriate organic carbon and energy source to the standard mineral base; the concentration was generally 1–3 g./l.

Unless otherwise stated, cultures were incubated at 30°. Stock cultures were transferred at monthly intervals, and stored at 5° after growth had occurred.

Screening of organic compounds as sources of carbon and energy for growth

Every strain was tested for ability to grow at the expense of 146 different organic compounds. These tests were performed by replica plating (Lederberg & Lederberg, 1952). The test media were prepared by adding each organic compound at the appropriate concentration to the standard mineral base, solidified by the addition of 1% (w/v) of Ionagar. In order to avoid browning of the agar, it was sterilized separately from the mineral base (both at double strength), and mixed with it after removal from the autoclave. Each test series included a control plate without added organic compound. The use of Ionagar in place of less highly purified agars almost completely eliminated background growth on the control plates, which greatly facilitated the reading of the results.

The choice of substrate concentration in a nutritional survey of this nature is a delicate problem. If the concentration is too high, false results can occur. False negative results may be caused by toxicity; and false positive ones by the presence of utilizable organic contaminants in an intrinsically non-utilizable chemical. For this

reason, it is advisable to keep the substrate concentration as low as possible. As a general rule, we used a concentration of 0.1 % (w/v). For organic acids and bases, this concentration expressed the weight of the organic anion or cation, even though they were added to the medium as salts. The only exceptions were sugars used at a concentration of 0.2 % (w/v), and phenol 0.025 % (w/v). This concentration of phenol, although low, still allows unambiguous growth of positive strains. A concentration of 0.05 % is already toxic to some strains which are capable of using phenol.

The above-mentioned concentrations support good growth by strains that are able to use the compound under test, provided that the plate is not too crowded. In order to avoid early substrate exhaustion, the plates were furnished with a relatively large volume of medium (35 ml.), and the number of patches placed on each plate was limited to a maximum of 23.

Many of the organic compounds used were of unknown thermostability. When thermostability was known, the compounds were sterilized by autoclaving with the mineral base, except for volatile compounds, which were sterilized by filtration. When thermostability was in doubt, the compounds were always sterilized by filtration. Geraniol and naphthalene, which are volatile hydrophobic substances, were not sterilized or added to the medium, but placed in the inverted lid of the Petri dish after its inoculation; their effective concentrations in the medium are accordingly unknown.

For successful replication of the highly motile aerobic pseudomonads, it is essential to use well-dried agar plates. To ensure this, the uninoculated plates were placed in an incubator at 37° for 2–3 days after their preparation. Once dry, they were stored at room temperature in plastic bags, which minimized both contamination and further water loss. They could be kept for 4–5 weeks before use.

When a group of strains was to be screened, a set of master plates was prepared by hand patching on yeast agar. After growth had occurred, each plate could be used to prepare as many as ten submasters by replication on the same medium. When the submaster plates had grown, each one could be used to print nine different test plates, followed by a terminal yeast agar plate to control the successful replication of all patches. Accordingly, a single master plate could be used to test a maximum of 23 different strains on a maximum of 90 different substrates. A few strains produced growth on solid media which could not be successfully replicated, or else did not grow well on yeast agar; they were screened by inoculating the test plates with a small drop of a liquid culture, grown to a carbon limit.

The inoculated test plates were first scored after 48 hr, and examined again after a further 48 hr. Geraniol plates, on which growth was very slow, were scored only after 7 days of incubation. Scoring was visual. The growth was recorded as: 0 (growth no greater than on the control plate without carbon source); + (good growth); ± (scanty growth, but significantly more than on the control plate); and +M (late growth of a few colonies, in the area of the patch, presumed to arise from mutants in the inoculum).

When strains of markedly different nutritional character were tested together on a plate, the results were sometimes difficult to interpret, either because of cross-feeding (particularly marked with sucrose and acetamide) or of starvation of slowly growing patches surrounded by rapidly growing ones. In view of these problems,

the initial random screening was followed by a second series of screenings, in which the strains were re-assorted so that only those of one nutritional type were printed on one set of plates.

Instances of marginal growth on plates (dubious $\pm$ values) were further examined by tests of growth in liquid media, where the yield could be determined quantitatively by turbidimetry. This control proved important in some cases, since relatively little total growth on the plates suffices to produce a $\pm$ reading. Our experience with sucrose utilization provides a good example, and also illustrates the danger of assuming too lightly that a particular compound is thermostable. We at first sterilized this non-reducing disaccharide either by autoclaving a concentrated aqueous solution (25 %, w/v) and mixing with the medium after sterilization, or by autoclaving the medium containing the sucrose in the final concentration. In the plate screenings, almost every fluorescent pseudomonad was scored as either + or $\pm$ on sucrose, despite the fact that some members of this group (notably *Pseudomonas aeruginosa*) have been reported by many workers not to attack sucrose, as judged by tests with the medium of Hugh & Leifson (1953). Tested in liquid sucrose media, most of the strains scored as $\pm$ showed very low growth yields. The whole collection was then retested on plates prepared with filter-sterilized sucrose; practically all the strains originally scored as $\pm$ now failed to grow. Evidently, sucrose does decompose to a slight extent on autoclaving in aqueous solution, and some of the decomposition products are utilizable for growth by nutritionally versatile aerobic pseudomonads. This slight thermal instability does not interfere with tests for sucrose utilization assessed in terms of acid production.

The organic compounds tested as substrates were:

(a) Carbohydrates and sugar derivatives: D-ribose, D-xylose, D-arabinose, L-arabinose, D-fucose, L-rhamnose, D-glucose, D-mannose, D-galactose, D-fructose, sucrose, trehalose, maltose, cellobiose, lactose, starch, inulin, gluconate, 2-ketogluconate, saccharate, mucate, salicin.

(b) Fatty acids: acetate, propionate, butyrate, isobutyrate, valerate, isovalerate, caproate, heptanoate, caprylate, pelargonate, caprate.

(c) Dicarboxylic acids: oxalate, malonate, succinate, maleate, fumarate, glutarate, adipate, pimelate, suberate, azelate, sebacate, eicosanedioate.

(d) Hydroxyacids: D-malate, L-malate, D-(−)-tartrate, L-(+)-tartrate, meso-tartrate, DL- β -hydroxybutyrate, DL-lactate, glycollate, DL-glycerate, hydroxymethylglutarate, poly- β -hydroxybutyrate.

(e) Miscellaneous organic acids: citrate, α -ketoglutarate, pyruvate, aconitate, laevulinate, citraconate, itaconate, mesaconate.

(f) Polyalcohols and glycols: erythritol, mannitol, sorbitol, meso-inositol, adonitol, glycerol, ethyleneglycol, propyleneglycol, 2,3-butyleneglycol.

(g) Alcohols: methanol, ethanol, *n*-propanol, isopropanol, *n*-butanol, isobutanol, geraniol.

(h) Non-nitrogenous aromatic and other cyclic compounds: D-mandelate, L-mandelate, benzoylformate, benzoate, *o*-hydroxybenzoate, *m*-hydroxybenzoate, *p*-hydroxybenzoate, phthalate, isophthalate, terephthalate, phenylacetate, phenylethanediole, naphthalene, phenol, quinate, testosterone.

(i) Aliphatic amino acids: glycine, L- α -alanine, D- α -alanine, β -alanine, L-serine, L-threonine, L-leucine, L-isoleucine, DL-norleucine, L-valine, L-aspartate, L-gluta-

mate, L-lysine, DL-arginine, DL-ornithine, DL-citrulline, DL- α -aminobutyrate, γ -aminobutyrate, DL- α -aminovalerate, δ -aminovalerate.

(j) Amino acids and related compounds containing a ring structure: L-histidine, L-proline, L-tyrosine, L-phenylalanine, L-tryptophan, D-tryptophan, L-kynurenine, kynurenate, anthranilate, *m*-aminobenzoate, *p*-aminobenzoate.

(k) Amines: methylamine, ethanolamine, benzylamine, putrescine, spermine, histamine, tryptamine, butylamine, α -amylamine.

(l) Miscellaneous nitrogenous compounds: betaine, sarcosine, creatine, hippurate, pantothenate, acetamide, nicotinate, trigonelline.

(m) Paraffin hydrocarbons: *n*-dodecane, *n*-hexadecane.

In all cases, the purest commercially available preparations were used.

Hydrocarbon utilization. The utilization of hydrocarbons as sole sources of carbon and energy cannot easily be determined on solid media because of the difficulty of obtaining a properly emulsified preparation of these water-insoluble compounds. This property was therefore determined in tubes containing 4–5 ml. liquid mineral medium to which a few drops of the compound had been added under aseptic conditions from a filter-sterilized sample. In our tests we have only used highly purified *n*-dodecane and *n*-hexadecane. The tubes were incubated at 30° on a rotary shaker and the results recorded daily for a period of 7 days.

Tests for chemolithotrophic growth with H₂

Our standard mineral base was not suitable for growth at the expense of H₂; nearly all hydrogenomonads tested grew poorly or not at all chemolithotrophically when tested in it, although they grew excellently at the expense of organic compounds in the same basal medium. Consequently, the simple non-chelated mineral base of Doudoroff (1940) was used in all tests for chemolithotrophy. It contains (per litre): 33 ml. Na₂HPO₄ + KH₂PO₄ buffer (M, pH 6·8); 1 g. NH₄Cl; 0·5 g. MgSO₄·7H₂O; 0·05 g. FeCl₃·6H₂O; 0·005 g. CaCl₂. The last two ingredients were autoclaved separately at tenfold strength, to avoid formation of a precipitate. Solid medium was prepared by addition of Ionagar to 10 g./l. (separately sterilized).

Tests were performed by patching strains on plates of this medium and incubating them at 30° in a desiccator containing an atmosphere of (% v/v) 65, hydrogen; 5, CO₂; 6, oxygen; 24, nitrogen. Control plates were incubated in air. When growth on the experimental plate was no greater than on the control after incubation for 2 weeks, the strain in question was presumed to be incapable of growing chemolithotrophically at the expense of H₂. All strains which appeared positive by the plate test were examined for their ability to grow in the homologous liquid medium under the same conditions of incubation, and only when growth also occurred under these conditions were they deemed to be hydrogen chemolithotrophs.

Tests for denitrification

Tests were conducted in yeast extract (YE) medium, supplemented with 10 g. glycerol, 10 g. KNO₃ and 1 g. Ionagar/l. Each culture was first grown for 24 hr in an unsealed tube containing 5 ml. of this medium incubated without mechanical agitation. A loopful of this semi-aerobic culture was transferred to a tube containing 10 ml. of the same medium, molten but cooled to about 40°. The contents were mixed, and the tube was chilled briefly to allow the medium to become semi-solid. An

overlay of 2–3 ml. of 1.0 % (w/v) Ionagar in water provided an anaerobic seal and a trap for gas production. Vigorous denitrifiers like *Pseudomonas aeruginosa* produce dense turbidity in this medium after 18–24 hr of incubation at 30°, soon followed by abundant gas production. Other denitrifying pseudomonads develop more slowly, and the tubes should be incubated 5 days before recording a negative result. With some strains, heavy anaerobic growth occurred without visible gas production; such strains were reported as positive. Pre-cultivation in the same medium under semi-aerobic conditions is important, since in some species prolonged aerobic cultivation in a nitrate-free medium seems to result in a loss or weakening of denitrifying ability, presumably as a result of mutation and selection. The pre-cultivation step selects once more for denitrifying ability, and this ensures prompt growth of potential denitrifying strains when placed under rigorously anaerobic conditions.

We have also examined growth under denitrifying conditions in defined media, prepared and used as described above. These media were devoid of yeast extract, and furnished with either glycerol or Na lactate (0.5 %, w/v) as carbon and energy source. The results were closely comparable, but some organisms (notably *P. stutzeri*) grew and produced gas more promptly in the defined media.

Determination of pigment production

Ability to produce specific pigments is a very useful determinative character among aerobic pseudomonads. However, synthesis of the two commonest types of pigments found in these bacteria (water-soluble fluorescent pigments and phenazine pigments) is highly dependent on the composition of the medium, and hence not manifested under all conditions of growth. King, Ward & Raney (1954) developed empirically two complex media, with different mineral additions (media A and B, hereafter termed 'King A' and 'King B') which are very helpful in detecting these two classes of pigments. King A favours the production of phenazine pigments, though some strains will also produce a slight amount of fluorescent pigment on it; King B favours the production of fluorescent pigment, and in most cases completely suppresses production of phenazine pigments. Although these media were devised specifically for the differentiation of *Pseudomonas aeruginosa* from non-phenazine-producing fluorescent pseudomonads, we found them to be satisfactory for the recognition of two other phenazine producers, *P. chlororaphis* and *P. aureofaciens*. They are prepared as slopes, and inoculated by streaking. Cultures were examined for pigmentation after 24, 48, and 72 hr (incubation at 30°). Véron (1961) recommended longer periods of incubation, but we have not found this to be necessary.

The phenazine pigments characteristic of *Pseudomonas aeruginosa* and *P. aureofaciens* (pyocyanine, phenazine α -carboxylic acid, respectively) are water-soluble, and can be recognized by the characteristic colours that they impart to slopes of King A. One phenazine pigment of *P. chlororaphis*, chlororaphin, is very sparingly soluble in water, and accumulates as isolated green crystals in the growth at the butt of a King A slope; the other pigment, oxychlororaphin, is relatively soluble, and imparts a pale yellow colour to the slope.

One other fluorescent pseudomonad, *Pseudomonas lemonnieri*, is characterized by the production of a water-insoluble chloroform-soluble blue pigment which is probably

a phenazine (Starr, Blan & Cosens, 1960). The blue pigment could not be elicited by growing the organisms in the King media. Production of the blue pigment also failed in both the complex and the chemically defined media used throughout our work. In investigating this point further, we found that the pigment could be formed on a solid medium of the following composition (% w/v): Bacto peptone, 0.5; glucose, 1.0; Bacto agar, 2; in tap water. The pigment was not produced when the medium contained in addition phosphate buffer (M/30, pH 6.8). Two strains of our collection (83, 84), which had the same origin but were received from different laboratories, produced the blue pigment. The pigment appeared after 24 hr of incubation in the colonies of strain 84, while the colonies of strain 83 showed after a few days several blue-pigmented sectors. Strain 143 totally failed to produce the blue colour even after long incubation, although the pigment had been observed in the colonies of the original isolate. The reappearance of the blue variants in strain 83 and the apparent total loss of the character in strain 143 suggest that the production of the blue pigment is a property that is counter-selected under conditions of laboratory maintenance.

Among the non-fluorescent pseudomonads, many strains of the species that we designate as *Pseudomonas multivorans* produce phenazine pigments of undetermined structure. Apparently two such pigments may be synthesized, both of which are water-soluble under neutral or alkaline conditions; one is yellow and the other is purple (Morris & Roberts, 1959). Strains belonging to this species may produce both pigments, yellow pigment, or none. Both pigments are formed abundantly on King A medium, which is accordingly a useful medium for detection. Both the organisms and the slope appeared coloured.

The majority of the strains of *Pseudomonas maltophilia* produce a pale yellow cellular pigment which we have not studied in detail. This pigment is soluble in water and is not extracted by any of the organic solvents used for the carotenoid pigments. We have also noted that the strains of *P. stutzeri* are capable of producing a faintly yellow diffusible pigment on the King B medium. This pigment is not fluorescent and has not been further characterized.

Biochemical methods

Screening for ring fission mechanisms. In recent years it has become evident that aerobic pseudomonads can de-aromatize two of the central intermediates in the metabolism of aromatic compounds, (catechol, protocatechuic acid), by two different mechanisms, *ortho* and *meta* cleavage (Dagley, Evans & Ribbons, 1960). These two modes of cleavage channel the degradation of a given aromatic compound after ring fission into two completely different metabolic pathways. Hence it seemed probable *a priori* that the mode of cleavage would be a taxonomically significant character. Two strains may both utilize *p*-hydroxybenzoate as a growth substrate; on this nutritional character, they appear identical. But, if it can be shown that they use different cleavage mechanisms for attack on the benzenoid nucleus, a whole series of enzymic differences is at once revealed. The problem was to devise a simple way of testing for the difference. This was solved by Dr Keiichi Hosokawa, whose elegant unpublished method we systematically applied to the collection.

Strains are grown as patches on chemically defined media containing an aromatic substrate as sole source of carbon and energy. We have most commonly used

p-hydroxybenzoate as a substrate; this compound is decomposed via protocatechuate. Each patch is scraped off and suspended in 0.02 M-tris buffer (pH 8.0). The suspension is toluenized and 2 ml. samples are furnished with 20 μ moles of either catechol or sodium protocatechuate. When *meta* cleavage occurs, the mixture turns bright yellow within a few minutes. The tube is shaken for 1 hr at 30°, and then tested for the presence of β -ketoadipate (indicative of an *ortho* cleavage) by the Rothera reaction (Kilby, 1948). An unusual deep purple Rothera reaction is given by this β -keto acid. The development of this colour is strong presumptive evidence of *ortho* cleavage. A parallel control experiment, using patches of organism grown on yeast agar plates, was always done to determine whether the observed activity had been specifically induced by growth on the aromatic substrate.

Assay of the arginine dihydrolase system. We assayed this dihydrolase system as follows. The organisms, grown on mineral+lactate medium, were suspended in M/30 phosphate buffer (pH 6.8) to an optical density of 200 Klett units (green filter 54). Four ml. of bacterial suspension in a test tube were gassed with N₂, and 1 ml. of 10⁻³ M-arginine was added. The mixture was gassed again with N₂ and the tube stoppered and incubated at 30° for 2 hr. At the end of the incubation, the tube was immersed in a boiling water bath for 15 min., and arginine determined in a 1 ml. sample. As control, we used an identical suspension which was heated immediately after the addition of arginine, and thereafter subjected to the same treatment. Arginine was determined by the quantitative method of Rosenberg, Ennor & Morrison (1956).

Oxidase test. The oxidase test was performed by soaking a strip of filter paper with 2-3 drops of a 1% (w/v) aqueous solution of *N,N'*-dimethyl-*p*-phenylenediamine, and immediately smearing a loopful of bacteria from a slope culture on the moist area. Oxidase-positive strains turn dark red within a few seconds. The test can also be performed with 0.2% (w/v) aqueous 2,6-dichlorophenolindophenol, first reduced with dithionite by careful addition of the reducing agent until the blue colour just disappears. With this reagent, oxidase-positive strains turn blue.

Determination of cytochrome difference spectra. Bacteria were grown overnight in shaken flasks of yeast extract medium, harvested, washed, and resuspended in 0.03 M-phosphate buffer (pH 6.8) to a concentration equiv. 2.0-2.5 mg. dry-wt./ml. Two samples of this suspension were placed in cuvettes (1 cm. light path), and gassed with N₂ (O₂-free) for at least 2 min. The absorbances were compared over the range from 390 to 650 m μ , by using a Cary recording spectrophotometer with a sensitive slide wire (total optical density span: 0.2 unit). In one cuvette, the components of the electron transport system were then oxidized by the addition of 3 drops of 3% H₂O₂, and this suspension was immediately used as a blank to record absorbance over the same spectral range in the reduced suspension. The difference spectrum so obtained (reduced minus oxidized), corrected for minor differences of absorbance (reduced minus reduced) caused by the fact that absorbances of the two cuvettes are never exactly comparable at this degree of sensitivity, shows the peaks of the reduced components of the cytochrome system. To identify precisely the peaks in the Soret (γ) region, the determination must be repeated with a more dilute suspension.

Production of extracellular enzymes. The capacity to produce extracellular proteases was determined with gelatin as substrate, by the plate method described by

Skerman (1959). Patches of growth on plates of Skerman's medium were allowed to grow for 48 hr, and the plates then flooded with an acidic solution of mercuric chloride, which precipitated the unhydrolysed gelatin in the agar. When the clear zone around the patch was not more than 2 mm. in extent, gelatinase production was recorded as weak ($\pm$). Wider zones of clearing were recorded as +. Some strains showed partial clearing under the patch, not extending beyond its edge. These were recorded as -, since the clearing might simply reflect partial autolysis of the bacteria with liberation of proteolytic endoenzymes.

The egg-yolk reaction, useful for internal subdivision of fluorescent pseudomonads, was done on the medium recommended by the *Manual of Microbiological Methods* (1957), patched with the strains to be tested. A positive reaction is shown by the formation of a white precipitate surrounding the patch of growth, which generally appears after 24 hr, although readings should be continued for 6 days.

Starch hydrolysis was tested by patching strains on yeast agar supplemented with 0.2% (w/v) soluble starch, and flooding the plates with Lugol's solution after incubation for 48 hr.

The production of extracellular lipases was tested by patching strains on the Tween 80 medium recommended by Sierra (1957). Hydrolysis of the detergent is shown by formation in the medium of a precipitate of calcium oleate. The time of appearance of a turbid halo surrounding the patch of growth is very variable; we arbitrarily scored as positive those strains which showed such a halo after not more than 6 days of incubation. A weak reaction occurring after a longer period may simply reflect autolysis of the bacteria and liberation of intracellular esterases.

The production of an extracellular esterase capable of depolymerizing poly- β -hydroxybutyric acid was determined by patching strains on mineral agar plates overlaid with a suspension of poly- β -hydroxybutyric acid (0.25%, w/v) in mineral agar. Positive strains showed growth of the patch and clearing of the surrounding medium. This is a test of great taxonomic value among aerobic pseudomonads, but unfortunately the substrate, a natural product not commercially available, is tedious to prepare. We used polymer granules extracted from *Bacillus megaterium* and prepared as described by Delafield *et al.* (1965).

Cytological observations

We examined sizes of organisms and were unable to find consistent differences among the strains of different species; therefore, no mention will be made of this character.

Flagellar staining was done by the method of Leifson (1951) on suspensions of bacteria prepared from young slope cultures on yeast agar. In evaluating flagellation as 'monotrichous' or 'multitrichous', we used where necessary the statistical method of analysis recommended by Lautrop & Jessen (1964).

The accumulation of poly- β -hydroxybutyric acid as an intracellular reserve material is an extremely valuable taxonomic character among aerobic pseudomonads. It was detected by microscopic examination of bacteria under phase contrast, and confirmed where necessary by staining smears with Sudan Black or by chemical extraction (Williamson & Wilkinson, 1958). The accumulation of cellular organic reserve materials is favoured under conditions of nitrogen starvation (Doudoroff & Stanier, 1959); and in some purple bacteria, which can form both

glycogen and poly- β -hydroxybutyrate reserves, the formation of poly- β -hydroxybutyrate is dependent on the provision of an organic substrate metabolically closely related to it. These considerations led us to determine poly- β -hydroxybutyrate accumulation on bacteria grown in a special chemically defined medium, consisting of standard mineral base with $(\text{NH}_4)_2\text{SO}_4$ at a low concentration (0.2 g./l.), furnished with DL- β -hydroxybutyrate (5 g./l.) as carbon and energy source. The medium was used in liquid form (5 ml./tube); cultures were mechanically agitated, and examined after 24–48 hr, when growth had become nitrogen limited. Under these circumstances, polymer-accumulating species regularly contained extensive intracellular deposits of the polymer, readily observable by microscopic examination of wet mounts. Although β -hydroxybutyrate is a carbon and energy source for most aerobic pseudomonads, a few cannot use it. For such organisms, either Na acetate or Na succinate (5 g./l.) was used instead.

Temperature relationships

Ability to grow at 4, 37, 39 and 41° was determined in tubes of yeast extract medium, inoculated with a loopful of culture grown for 12–18 hr in the same medium at 30°. The tubes incubated at 4° were read after 10 days, the others after 24 hr. Growth was recorded as abundant (+) or slight ($\pm$). Water baths were used for all determinations; the cultures were not shaken.

Nitrogen sources and growth factor requirements

With the exception of *Pseudomonas maltophilia*, all species described in this paper grow well in standard chemically defined medium, furnished with NH_4^+ as nitrogen source and any utilizable organic carbon and energy source. No strain of *P. maltophilia* will grow under these conditions. It was therefore necessary to determine the minimal nutritional requirements of this species, as a preliminary to examining its carbon and energy requirements. Tests with supplements of amino acids, vitamins, purines and pyrimidines led to the discovery that all strains of this species require methionine, but no other organic growth factor. (In the process of screening amino acids, we observed that leucine was also stimulatory; but this effect is probably attributable to the fact, reported by Demain (1965), that many commercial preparations of leucine are contaminated with methionine.)

All chemically defined media used for the growth and nutritional screening of *P. maltophilia* accordingly had to be supplemented with methionine, which was added to 20 mg./l.

All strains of the collection were tested for their ability to use nitrate as a nitrogen source, and to fix atmospheric nitrogen. Such tests were conducted in tubes of the standard mineral base, in which NH_4^+ was either replaced by KNO_3 (1 g./l.) or omitted. Lactate (1 g./l.) was provided as source of carbon and energy.

Determination of base composition of DNA

The moles % of guanine + cytosine (GC) in the DNA of many of our strains was determined by Dr Manley Mandel, from measurements of buoyant density in CsCl gradients (Mandel, 1966).

Miscellaneous tests

Certain *Pseudomonas* species are reported to produce indole (Haynes, 1957). However, during an extensive survey of the mechanisms of tryptophan dissimilation by pseudomonads (Stanier & Hayaishi, 1951) the one strain found to produce indole (Tsuchida, Hayaishi & Stanier, 1952) was later shown to have been mis-identified, and to be in reality a coliform bacterium (Stanier, 1952). There is, accordingly, no reliable biochemical evidence for the occurrence of tryptophanase among aerobic pseudomonads. To resolve this question, we tested the entire collection for indole production by the customary procedures (Skerman, 1959). The only organisms which gave a pink colour upon addition of the Ehrlich reagent to the culture were certain strains of *P. aeruginosa*, which had formed pyocyanine during growth. Since pyocyanine is red in the acid form, these strains were retested for indole production by the Goré (1921) modification of the usual reaction, in which detection of indole is dependent upon its volatility. The results were then totally negative. We conclude that no aerobic pseudomonad so far studied is able to produce indole, and that this test is diagnostically valueless in the taxonomic analysis of the group. False positives can result from the presence of pyocyanine in the medium.

The oxidation of indole to the insoluble blue compound, indigotin, originally described by Gray (1928) as the salient property of *P. indoloxidans*, was tested on plates of the medium recommended by Gray, except that Na acetate was substituted for glycerol as the carbon source for strains unable to use glycerol. In positive strains, the patch and the adjacent agar turn blue, as a result of the deposition of crystals of indigotin.

Tests for the capacity to grow anaerobically with glucose as a fermentable substrate were conducted in tubes of yeast extract medium supplemented with 1 % (w/v) glucose and 0.1 % (w/v) Ionagar, and overlaid after inoculation with a plain agar seal.

Biological materials

The nucleus of the collection consisted of unidentified *Pseudomonas* strains isolated in our laboratory in Berkeley by enrichment methods over the past 20 years. It was supplemented by many additional strains received from other sources. Before inclusion, each strain was streaked on yeast agar to check its purity and homogeneity. Many of the strains proved to be heterogeneous in colony form. The most common type of colonial heterogeneity was represented by a mixture of relatively translucent, flat colonies and more opaque, raised, compact colonies. This was particularly frequent among fluorescent pseudomonads, and, since our past experience with this group indicated that the translucent flat colony is the wild-type form, we attempted as far as possible to select this form for the establishment of stock cultures. In a few cases where both colony forms were isolated from a single strain and subjected to detailed analysis, no other phenotypic differences between them were observed.

Each strain was also examined for Gram reaction, motility, and ability to grow anaerobically with glucose. Only strains which were Gram-negative, actively motile straight or curved rods and incapable of anaerobic growth with glucose were included.

At the time of its addition to the collection, each strain was given a serial number ($1 \rightarrow n$), and we shall designate strains by these numbers in the discussion of the results. Every strain has now been placed in the American Type Culture Collection, and therefore also carries an ATCC accession number. Specific information on sources, past histories, and ATCC or other collection numbers will be given in the strain descriptions, which are included with the descriptions of the species groups or the individual species.

RESULTS

Primary differentiation of species and species groups

Nearly all the 267 individual strains that we have examined can be assigned to one of 10 species (Table 1). Fortunately, eight of these species have been named by earlier workers. Some can be placed in subgeneric categories which share many common properties, and which we term 'groups'. Taxonomically, the most complex

Table 1. *Specific assignments of 267 individual strains examined in the present study*

Subgeneric group	Species	No. of strains
Fluorescent group	<i>Pseudomonas aeruginosa</i>	29
	<i>P. fluorescens</i>	94
	<i>P. putida</i>	41
	Unassigned	11
Acidovorans group	<i>P. acidovorans</i>	15
	<i>P. testosteroni</i>	9
	Unassigned	2
Alcaligenes group	<i>P. alcaligenes</i>	1
	<i>P. pseudoalcaligenes</i> sp.nov.	6
	<i>P. multivorans</i> sp.nov.	19
	<i>P. stutzeri</i>	17
	<i>P. maltophilia</i>	23
Total		267

of these is the 'fluorescent group', in which we recognize three species: *Pseudomonas aeruginosa*, *P. fluorescens* and *P. putida*. *P. aeruginosa* shows a high degree of internal phenotypic uniformity, but both *P. fluorescens* and *P. putida* can in turn be subdivided into a number of distinct 'biotypes', to which we assign identifying letters. Some of these fluorescent biotypes may eventually prove worthy of specific recognition; but, in our opinion, such recognition is not now justified. The 'acidovorans group' contains two species, *P. acidovorans* and *P. testosteroni*. The 'alcaligenes group' contains two species, *P. alcaligenes* and *P. pseudoalcaligenes* sp.nov. The remaining three species that we recognize are all phenotypically somewhat isolated, both from the three groups so far described, and from one another. They are *P. maltophilia*, *P. stutzeri* and *P. multivorans* sp.nov.

Tables 2 and 3 list the phenotypic characters which are most useful for distinguishing these 10 species: in effect, these tables constitute a diagnostic key. In order to make these tables as complete as possible, we have also included data for the 'pseudomallei group' (*Pseudomonas pseudomallei* and *P. mallei* comb.nov.), taken from the paper of Redfearn *et al.* (1966); and for *P. lemoignei*, taken from the paper of Delafield *et al.* (1965). These are the only other members of the genus *Pseudomonas* so far fully characterized by our methods. Other described species which are

probably valid ones (e.g. *P. putrefaciens* and *P. diminuta*) cannot for this reason be included in the tables.

Tables 2 and 3 by no means include all the characters that we have found to be of some diagnostic value; in particular, many other diagnostically significant nutritional properties could have been added. In these tables, we have emphasized the characters which in our experience are of greatest *general* value for presumptive specific identification; the lists are minimal. If need be, the differentiation between closely related species can be strengthened by more extensive tests. For example, the difficult differentiation between *Pseudomonas fluorescens* and *P. putida*, which depends on only three of the characters listed in Tables 2 and 3, can be strengthened by determining the reaction on egg yolk, and the utilization of other carbohydrates and higher amines. Similarly, the differentiation between *P. acidovorans* and *P. testosteroni*, which depends on six of the characters listed in Table 3, can be strengthened by testing utilization of fructose, malonate, L-tartrate and D-tryptophan. These are all nutritional characters that we have omitted from Table 3 because they are of relatively restricted diagnostic value.

In Tables 2 and 3, the notation 'v' indicates that the character in question is variable for a particular species. Such characters may still of course be of diagnostic value for certain strains, if related species are uniformly either + or - for the same character. It should be emphasized that many of the physiological characters to which we have assigned '+' values may be absent from an occasional strain, but are overwhelmingly positive for the group in question. For example, we have listed fluorescent pigment production as '+' for all species of the fluorescent group, although non-fluorescent strains occur. Even such a characteristic trait of the fluorescent pseudomonads as the ability to grow on glucose is absent from one of the 175 strains that we examined. In contrast, the physiological characters to which we have assigned '-' values are truly absent from every strain of the species. Where this is not so, we have used the symbol '-(+)' to show that rare strains may possess the character in question. One example is the utilization of *m*-hydroxybenzoate by *Pseudomonas fluorescens* and *P. putida*, positive for a total of seven out of 135 strains assigned to these two species. Despite these sporadic exceptions, failure to grow on *m*-hydroxybenzoate is a useful nutritional character for distinguishing fluorescent pseudomonads as a group from *P. multivorans* and the acidovorans group, where nearly all strains can use it.

Rare exceptions to the predominant specific phenotype do not seriously interfere with the correct specific assignments, since a strain that is aberrant in one or two characters is in almost every instance still identifiable on the basis of the other characters listed in Table 2, 3. Hence these thirty-nine selected characters should suffice to permit a correct presumptive identification of most strains that belong to one of the listed species.

Taxonomic significance of poly- β -hydroxybutyrate metabolism in aerobic pseudomonads

The suggestion of Forsyth *et al.* (1958) that poly- β -hydroxybutyrate accumulation might be a useful differential character among aerobic pseudomonads has been confirmed by our work. In fact, this is probably the best single character for the primary subdivision of the genus *Pseudomonas*. The polymer producers are

	No. of flagella	Poly- β -OH-butyrate as cellular reserve	Fluorescent Pigments	Phenazine	Methionine requirement	Denitrification	Growth at $\left\{ \begin{array}{l} 4^{\circ} \\ 41^{\circ} \end{array} \right.$	Extracellular hydrolyses	Gelatin	Poly- β -OH-butyrate	Oxidase reaction	Arginine dihydrolase	Cleavage mechanism for diphenols
Fluorescent group	<i>P. aeruginosa</i>	1	-	+	+	+	+	+	+	+	+	+	ortho
	<i>P. fluorescens</i>	1 >	-	+	v	-	v	+	+	+	-	+	
Pseudomallei group*	<i>P. putida</i>	1 >	-	+	-	-	v	-	-	-	+	+	ortho
	<i>P. multivorans</i>	1 >	+	+	v	-	-	v	+	-	-	+	
Pseudomallei group*	<i>P. pseudomallei</i>	1 >	+	-	-	-	+	+	+	+	+	+	meta
	<i>P. mallei</i>	0	+	+	-	-	v	-	+	+	+	v	
Acidovorans group	<i>P. acidovorans</i>	1 >	+	+	-	-	-	-	-	-	-	+	meta
	<i>P. testosteroni</i>	1 >	+	+	-	-	-	-	-	-	v	-	
Alcaligenes group	<i>P. alcaligenes</i>	1	-	-	-	-	-	+	+	-	-	+	None
	<i>P. pseudoalcaligenes</i>	1	v	-	-	-	-	-	+	v	-	-	
	<i>P. stutzeri</i>	1	-	-	-	-	+	-	v	+	+	+	ortho
	<i>P. maltophilia</i>	1 >	-	-	-	+	-	-	-	+	-	-	
	<i>P. lemoignei</i> †	1	+	-	-	-	-	-	-	-	+	+	None

* Data of Redfearn *et al.* (1986). † Data of Delafeld *et al.* (1965). v = variable.

Table 3. Nutritional characters of diagnostic value for the differentiation of species of aerobic pseudomonads

Utilization as sole carbon and energy source of:	Fluorescent group			Pseudomallei group*			Acidovorans group			Alcaligenes group		
<i>P. aeruginosa</i>	+	+	-	-	-	-	+	+	+	-	-	-
<i>P. fluorescens</i>	+	+	+	-	+	+	+	+	+	-	-	-
<i>P. putida</i>	+	+	-	-	-	-	+	+	+	-	-	-
<i>P. multivorans</i>	+	+	+	+	+	+	+	+	+	+	+	+
<i>P. pseudomallei</i>	+	+	+	+	+	+	+	+	+	+	+	+
<i>P. mallei</i>	+	+	+	+	+	+	+	+	+	+	+	+
<i>P. acidovorans</i>	-	-	-	-	-	-	+	+	+	+	+	+
<i>P. testosteroni</i>	-	-	-	-	-	-	+	+	+	+	+	+
<i>P. alcaligenes</i>	-	-	-	-	-	-	+	+	+	+	+	+
<i>P. pseudoalcaligenes</i>	-	-	-	-	-	-	+	+	+	+	+	+
<i>P. stutzeri</i>	+	+	+	+	+	+	+	+	+	+	+	+
<i>P. maltophilia</i>	+	+	+	+	+	+	+	+	+	+	+	+
<i>P. lemoignei</i> †	+	+	+	+	+	+	+	+	+	+	+	+

* Data of Redfearn *et al.* (1966); † Data of Delafield *et al.* (1965).

+ = positive.

- (+) = usually negative, occasionally positive.

- = always negative.

P. multivorans, *P. pseudomallei*, *P. mallei*, *P. acidovorans*, *P. testosteroni* and *P. lemoignei* sp.nov. (Delafield *et al.* 1965). This character is completely absent from the fluorescent group, *P. stutzeri* and *P. maltophilia*. Only the alcaligenes group is variable, though predominantly positive: two out of six strains of *P. pseudocalcaligenes* fail to produce demonstrable polymer accumulations.

In considering these distributional regularities, it is important to realize that the ability to use poly- β -hydroxybutyrate as a cellular reserve material is, biochemically speaking, a complex character. As Merrick & Doudoroff (1961) showed, the polymer is synthesized from poly- β -hydroxybutyryl-CoA by a specific polymerase, and endogenously hydrolysed to the free hydroxy acid by a system which includes a proteinaceous activator, as well as at least two specific hydrolases. A multi-enzyme system is accordingly involved in the catalysis of this peripheral cycle on the pathway of fatty acid synthesis.

The ability to attack exogenous poly- β -hydroxybutyrate by means of extracellular enzymes is also a taxonomically valuable character. It is possessed by some, but not all, of the species able to synthesize the polymer as a cellular reserve material, and never occurs in species unable to do so. As Delafield *et al.* (1965) showed, the extracellular depolymerase system of *Pseudomonas lemoignei* is quite different from the enzyme system responsible for endogenous hydrolysis of polymer reserves in *Rhodospirillum rubrum* (Merrick & Doudoroff, 1964). Hence the capacity to hydrolyse and grow at the expense of exogenous poly- β -hydroxybutyrate may be an enzymic character independent of the capacity to use the polymer as an intracellular reserve. Species which regularly produce an extracellular depolymerase are *P. pseudomallei* and *P. lemoignei*; the character is variable in *P. testosteroni* and *P. mallei*.

Significance of denitrifying ability

The ability to use nitrate in place of O₂ as a terminal electron acceptor, and hence to grow well anaerobically in a nitrate-containing medium, is another multi-enzyme character that is of considerable taxonomic significance among the aerobic pseudomonads. It is a universal property of *Pseudomonas aeruginosa* (as already shown by Jessen, 1965), *P. pseudomallei*, *P. mallei* and *P. stutzeri*, but is absent from all other species discussed here except *P. fluorescens*. In *P. fluorescens*, some of the subgroups that we recognize as biotypes (B, C, D, F) are uniformly able to denitrify, while others (A, E, G) lack this property. It should be noted that each of these biotypes is distinguishable from the remaining ones by several additional characters, metabolically unrelated to the property of denitrification. Hence denitrification retains some degree of taxonomic significance even within the one species where it is variable, since it serves, in conjunction with other characters, to define biotypes.

The oxidase reaction and its relation to the cytochrome system

The oxidase reaction is positive for every species that we have examined except *Pseudomonas maltophilia*, which was also reported by Thibault (1961) and by Hugh & Leifson (1963) to be oxidase negative. The biochemical basis of the differences between aerobic bacteria in their response to this commonly used diagnostic test is not known. In an attempt to throw some light on this point, we determined dif-

ference spectra (reduced-oxidized) on heavy suspensions of a few representative strains of each species.

The results (Table 4) indicated that the specific cytochrome complement may be an extremely useful taxonomic character for the differentiation of groups and species among the aerobic pseudomonads, although this cannot be definitely established until more strains of each species have been examined. It is particularly notable that the fluorescent, acidovorans and alcaligenes groups all appear internally homogeneous with respect to cytochrome complement. In species that contain a cytochrome of the *c* type, the α -band of the *b*-type cytochrome lies on the shoulder of the *c* peak, and cannot be precisely located; indeed, its very presence in *Pseudomonas stutzeri*, which has an exceptionally high content of *c*-type cytochrome, is somewhat doubtful. However, every species examined shows a β -band at about 527–530 $m\mu$, reportedly characteristic of *b*-type cytochromes (Smith, 1961).

Table 4. Wavelengths ($m\mu$) of the main absorption peaks in the cytochrome difference spectra of selected strains of the different species of aerobic pseudomonads

Species	α -Bands. Cytochrome type			β -Bands. Cytochrome type		Principal Soret band
	<i>a</i>	<i>b</i>	<i>c</i>	<i>b</i>	<i>c</i>	
<i>P. aeruginosa</i>	.	<i>c</i> . 560	552	530	523	428
<i>P. fluorescens</i>	.	<i>c</i> . 560	552	530	523	427
<i>P. putida</i>	.	<i>c</i> . 560	552	530	523	425
<i>P. acidovorans</i>	600	.	553	530	524	425
<i>P. testosteroni</i>	600	.	553	530	523	423
<i>P. alcaligenes</i>	600	<i>c</i> . 560	553	530	523	428
<i>P. pseudoalcaligenes</i>	600	<i>c</i> . 560	551	528	522	423
<i>P. multivorans</i>	628, 597	<i>c</i> . 560	554	527	523	428
<i>P. stutzeri</i>	.	(<i>c</i> . 560)	552	530	523	423
<i>P. maltophilia</i>	628, 597	558	.	530	.	430
<i>P. lemoignei</i>	.	557	552	530	523	422

With regard to the interpretation of the oxidase reaction, it can be seen that *Pseudomonas maltophilia* differs from all other species by virtue of not showing an α -band at 551–554 $m\mu$, or a β -band at 522–524 $m\mu$. These spectral features are both characteristic of cytochromes of the *c* type.

Mr Paul Baumann (unpublished data) has found an analogous situation in the genus *Moraxella*, another group of aerobic bacteria that includes both oxidase-positive and oxidase-negative species: only oxidase-positive strains of *Moraxella* have peaks in their difference spectra referable to a cytochrome of the *c* type. These multiple correlations suggest that a positive oxidase test may indicate the presence of a cytochrome *c* in the terminal respiratory chain, although other possible explanations cannot yet be completely excluded.

The constitutive synthesis of arginine dihydrolase and its relation to the utilization of arginine as a carbon source

The anaerobic conversion of arginine to ornithine, ammonia and CO₂ through action of the so-called 'arginine dihydrolase' system (Hills, 1940) was first described in fluorescent pseudomonads by Slade, Doughty & Slamp (1954) and Sherris, Preston & Shoosmith (1957). This is a reaction of great physiological interest, since

it is accompanied by substrate-level generation of a high-energy phosphate bond (Slade *et al.* 1954), and thus constitutes a mechanism for the anaerobic synthesis of ATP. Indeed, as Sherris *et al.* (1957) have shown by simple but very elegant experiments, it provides the energy necessary for the maintenance of motility under anaerobic conditions by fluorescent pseudomonads, which as a rule rapidly become non-motile when their oxygen supply is depleted, like other strict aerobes. Their data suggested that the arginine dihydrolase system is synthesized constitutively by fluorescent pseudomonads, but is inactive on externally supplied arginine in the presence of O₂, becoming functional only under anaerobic conditions. We have confirmed this surprising fact by more rigorous and detailed unpublished experiments. The method which we recommend for the analysis of arginine dihydrolase activity (see section on Methods) gives positive results only when the enzyme system is constitutively synthesized.

Table 5. *The utilization of arginine by selected strains of Pseudomonas species*

Group	Species	No. of strains examined	No. of positive strains	
			Constitutive arginine dihydrolase	Use of arginine as carbon and energy source
Fluorescent	<i>P. aeruginosa</i>	4	4	4
	<i>P. fluorescens</i>	3	3	3
	<i>P. putida</i>	3	3	3
	<i>P. multivorans</i>	8	0	8
Pseudomallei*	<i>P. pseudomallei</i>	5	5	5
	<i>P. mallei</i>	10	10	10
Acidovorans	<i>P. acidovorans</i>	4	0	0
	<i>P. testosteroni</i>	4	0	0
Alcaligenes	<i>P. alcaligenes</i>	1	1	1
	<i>P. pseudoalcaligenes</i>	6	4†	5‡
	<i>P. stutzeri</i>	13	0	0
	<i>P. maltophilia</i>	2	0	0
	<i>P. lemoignei</i>	1	0	0

* Data of Redfearn *et al.* (1966).

† Strains 66 and 417 negative.

‡ Strain 417 negative.

Using less precise and stringent conditions for the study of the reaction, Sherris *et al.* (1959) and Thornley (1960) attempted to determine the taxonomic value of arginine dihydrolase production as a general character to distinguish aerobic pseudomonads from other Gram-negative eubacterial rods (e.g. coliform bacteria). Their data (obtained principally on fluorescent strains) suggested that it might be useful for this purpose. We have not examined every strain in our collection for the character of constitutive arginine dihydrolase synthesis, but we have tested a number of typical strains representative of each species; and the results shown in Table 5 are based on these samplings. In confirmation of the results of Sherris *et al.* (1959) and Thornley (1960), we find that every fluorescent strain examined is positive. However, this character is far from universal among the other species

examined, being present only in the pseudomallei and alcaligenes groups. As also shown in Table 5, the ability to use arginine as a carbon and energy source for aerobic growth is not necessarily correlated with the possession of a constitutive arginine dihydrolase system, since arginine is a good growth substrate for *Pseudomonas multivorans*, which does not contain a constitutive arginine dihydrolase system. Other aerobic pseudomonads (the acidovorans group, *P. stutzeri*, *P. maltophilia*, *P. lemoignei*) neither contain arginine dihydrolase, nor use arginine as an aerobic source of carbon and energy. In summary, the ability to synthesize arginine dihydrolase constitutively appears to be a useful character for the internal differentiation of species in the genus *Pseudomonas*, and not a generic trait.

Fig. 1. Mechanisms for the oxygenative cleavage of *ortho*-diphenols among pseudomonads.

Certain aromatic ring cleavage mechanisms

A biochemical character which we have examined systematically for the first time is the mechanism of cleavage of two diphenols, catechol and protocatechuate, the last aromatic intermediates in the oxidative metabolism of many aromatic compounds. Two modes of cleavage of these compounds have been described in various pseudomonads (Dagley *et al.* 1960): so-called '*ortho* cleavage' which results in the formation of a *cis-cis*-muconic acid; and so-called '*meta* cleavage' which results in the formation of an α -hydroxy-*cis-cis*-muconic semialdehyde (Fig. 1). Since the metabolic fates of the two classes of products of aromatic ring cleavage are wholly distinct, the cleavage mechanism in effect determines a long series of subsequent biochemical steps, so that two species which differ in cleavage mechanism differ also in many other specific enzymic respects.

The simple aromatic acid most extensively utilized by aerobic pseudomonads is *p*-hydroxybenzoate, which is degraded via protocatechuate. We have therefore tested the great majority of strains which can grow on *p*-hydroxybenzoate for the mode of ring cleavage of protocatechuate by *p*-hydroxybenzoate-grown organisms. Control tests on organisms of the same strains grown on yeast agar were always done, and usually gave negative results, thus confirming many past reports that the enzymes which mediate this cleavage reaction are strictly and specifically inducible. However, occasional strains of the fluorescent group contained the *ortho*-cleavage enzyme for protocatechuate after growth on yeast extract, suggesting that this enzyme is either partially constitutive, or induced by components of yeast extract in the strains in question.

The results of these tests (Table 6) show that the mechanism of cleavage of protocatechuate is a taxonomically significant character, since all the strains of a given species able to perform this reaction use the same cleavage mechanism. Secondly, it is evident that the *meta* cleavage mechanism is relatively rare, being confined exclusively to the two species of the acidovorans group, *Pseudomonas acidovorans* and *P. testosteroni*. The *ortho* cleavage of protocatechuate is characteristic of the entire fluorescent group, *P. pseudomallei*, *P. multivorans* and *P. stutzeri* (although very few strains of the last-mentioned species are able to grow on *p*-hydroxybenzoate).

Table 6. *Mechanisms for ring cleavage of protocatechuate among aerobic pseudomonads, tested after growth on p-hydroxybenzoate*

Species	No. of strains examined	No. of strains showing	
		<i>ortho</i> cleavage	<i>meta</i> cleavage
<i>P. aeruginosa</i>	5	5	0
<i>P. fluorescens</i>	62	62	0
<i>P. putida</i>	34	34	0
Unclassified fluorescent pseudomonads	8	8	0
<i>P. multivorans</i>	8	8	0
<i>P. pseudomallei</i>	6	6	0
<i>P. acidovorans</i>	15	0	15
<i>P. testosteroni</i>	9	0	9
<i>P. stutzeri</i>	3	3	0

Some universally negative nutritional characters of the strains examined

None of the 267 strains examined can grow chemolithotrophically at the expense of H₂. This fact shows that hydrogen chemolithotrophy is neither of random nor of wide occurrence among aerobic pseudomonads, and confers on it a taxonomic significance in the subdivision of this bacterial group that has hitherto been assumed by taxonomists without evidence. We do not consider it a suitable generic character, but at least it can now be used with some confidence as one of the characters of value in the definition of certain *Pseudomonas* species.

None of the 267 strains examined can grow on methanol, methylamine or oxalate. Hence the ability to utilize these compounds can also now be regarded as a significant taxonomic character among aerobic pseudomonads; it is presumably confined to a restricted segment of the genus *Pseudomonas*.

Lastly, it should be noted that none of our strains can grow in an otherwise suitable defined medium that is free from combined nitrogen compounds. Nitrogen fixation is, accordingly, a property absent from the group.

Nutritional versatility of the strains examined

As mentioned in the introduction, the unusually wide range of organic compounds utilizable as carbon and energy sources by *Pseudomonas fluorescens* was established through the comparative nutritional survey of den Dooren de Jong (1926) of a variety of different aerobic bacteria. This particular aspect of his work became well known, as a result of its detailed presentation in Marjory Stephenson's influential book *Bacterial Metabolism* (1939). *P. fluorescens* is, accordingly, often cited as the

epitome of nutritional versatility among bacteria. Since the quantitative aspects of the data on utilization of the 146 organic compounds tested as carbon and energy sources in our study are of some interest in this connexion, and are not easily extractable from the tabulations to be given in later sections, we have summarized them here. In Fig. 2, the range in the number of compounds utilizable by the strains that we assign to each species is expressed in bar form. Where many different strains of a species have been examined, the distribution within the bar is approximately Gaussian, so that the majority of strains cluster around the centre of each bar. In the case of *P. multivorans*, the bar has been prolonged by a dotted line, to

Fig. 2. The numerical range of organic compounds utilizable as sole sources of carbon and energy by certain species of aerobic pseudomonads. The values shown are based on tests with 146 different organic compounds. Data for the pseudomallei group are taken from the paper of Redfearn *et al.* (1966); data for *P. lemoignei* are taken from the paper of Delafield *et al.* (1965) and supplemented by further unpublished information.

indicate a single highly defective strain which falls well outside the normal distribution curve for this species.

In the fluorescent group, we have separated the data for biotypes A and B of *Pseudomonas putida*, since biotype B stands out as the most versatile of fluorescent pseudomonads (75–87 substrates used). The ranges for *P. putida* biotype A (66–82), *P. fluorescens* (59–82) and *P. aeruginosa* (76–82) are slightly lower and quite similar. The very wide span for *P. fluorescens* (all biotypes) is attributable partly to the very large number of strains included, and partly to the fact that the various biotypes differ nutritionally in minor respects from one another. The range for *P. pseudomallei*

(77–87 substrates used) is almost identical with that for *P. putida* biotype B, so that *P. pseudomallei* equals the best record in the fluorescent group. Both are well exceeded by *P. multivorans* (95–108 substrates used, if the single defective strain with a value of 73 is excluded). In the acidovorans group, *P. acidovorans* (68–73) falls only slightly below the fluorescent pseudomonads, while *P. testosteroni* uses markedly fewer compounds (53–63). This group is followed by *P. stutzeri*, whose characteristic nutritional irregularity is shown by a wide numerical span (41–59). At the lower end of the scale are *P. pseudoalcaligenes* (36–45), *P. alcaligenes* (24; one strain), *P. maltophilia* (24–28) and *P. lemoignei* (11; one strain). Fig. 2 confirms the fact, already evident from the nutritional properties of *P. methanica* (Dworkin & Foster, 1956), that the nutritional versatility of different species in the genus *Pseudomonas* differs very widely. It also shows that the numerical extent of the nutritional range is in some cases (e.g. *P. multivorans*) a highly distinctive specific character.

THE FLUORESCENT GROUP

Origins of the strains

The strains belonging to this group will be listed according to their specific assignments.

I. Strains assigned to *Pseudomonas aeruginosa*

45. *P. aeruginosa* strain LA P-6 of Dr S. C. Rittenberg. Isolated from hydrocarbon enrichment. ATCC 17423.
52. *P. aeruginosa* strain B-903 of Dr W. C. Haynes, U.S. Department of Agriculture, Peoria, Ill. ATCC 17429.
54. Same source as 52; strain B-23. ATCC 17431.
55. Same source as 52; strain B-1781. ATCC 17432.
57. Same source as 52; strain B-248. ATCC 17434.
58. Strain of unknown origin obtained from the departmental stock collection, Department of Bacteriology, University of California, Berkeley. ATCC 17435.
131. *P. aeruginosa* strain H-1c of Dr B. Holloway. ATCC 17503.
132. *P. aeruginosa* strain Ph-1, isolated by Dr G. Hegeman from a denitrification enrichment in Berkeley. ATCC 17504.
- 277–296. Independent strains isolated at several different hospitals from a variety of clinical specimens. Collection received from Dr Vera L. Sutter, School of Dentistry, San Francisco Medical Center. ATCC 17643–17662.
416. Strain received from Dr O. Jessen. ATCC 17831.

II. Strains assigned to *Pseudomonas fluorescens* biotype A

12. *P. fluorescens* strain 216 of den Dooren de Jong (1926). Isolated from tap water. ATCC 17397.
126. Strain mesaconic-12 of Dr M. Shilo. Isolated from soil by mesaconate enrichment. ATCC 17498.
182. Strain PJ 73 of Dr O. Jessen. Included in biotype 63 (group IV) by Jessen (1965). ATCC 17550.
184. Strain w4L-2 isolated from water by enrichment with lactate in Berkeley. ATCC 17552.
185. Strain w2L-1, isolated from water by enrichment with lactate in Berkeley. ATCC 17553.

The following 18 strains (186–216) were studied by Jessen (1965) and are all representative of his biotype 63, group V.

186. Strain PJ 79. Isolated and named *P. fluorescens* in the Inst. Infect. Diseases, Tokyo. ATCC 17554.
187. Strain PJ 139. Originally named *P. angulata*. ATCC 17555.
188. Strain PJ 160. Originally named *P. marginata* (NCPPB 316). ATCC 17556.
189. Strain PJ 227. *Pseudomonas* sp., culture 1208 from the Centre International Collection de Culture, Lausanne. ATCC 17557.
192. Strain PJ 239. This strain has been proposed by Dr Muriel E. Rhodes as neotype for *P. fluorescens*. ATCC 13525.
196. Strain PJ 288. Isolated from sputum in Hadeslev, Jutland. ATCC 17563.
197. Strain PJ 290. Isolated from throat swab in Hjørring, North Jutland. ATCC 17564.
198. Strain PJ 302. Isolated from pleural drainage in Slagalse, Mid-Sealand. ATCC 17565.
199. Strain PJ 311. Isolated from stored blood in Holbaek, North Sealand. ATCC 17566.
200. Strain PJ 367. Isolated from sputum (hospital sample) in Copenhagen. ATCC 17567.
201. Strain PJ 368. Isolated from saliva in the Seruminstitut, Copenhagen. ATCC 17568.
202. Strain PJ 372. From heart blood (autopsy), Copenhagen. ATCC 17569.
203. Strain PJ 376. From pleural drainage, Copenhagen. ATCC 17570.
208. Strain PJ 722. From polluted sea water, Øresund, south of Copenhagen. ATCC 17575.
209. Strain PJ 776. Isolated from polluted sea water, Øresund, north of Copenhagen. ATCC 17576.
211. Strain PJ 826. From ground water in south of Sealand. ATCC 17578.
215. Strain PJ 849. From surface water (natural stream) in south of Sealand. ATCC 17582.
216. Strain PJ 885. From ground water, south of Jutland. ATCC 17583.
392. Strain B-2075 of Dr W. C. Haynes, Peoria. Strain 4 of Lysenko, isolated from a caterpillar and named *P. chlororaphis*. ATCC 17813.

III. Strains assigned to *Pseudomonas fluorescens* biotype B

2. Strain P3T-1, isolated from water by L-tryptophan enrichment. ATCC 17387.
93. Strain TT-10 of Dr S. H. Hutner. Isolated from tryptophan enrichment. ATCC 17467.
108. Strain 52-22 of W. R. Sistrom. From naphthalene enrichment. ATCC 17482.

The following strains (400–415) were studied by Jessen (1965), and all are representatives of his biotype 61, group V, except strain 415, which he placed in biotype 62, group V.

400. Strain PJ 185. Named *P. marginalis* (NCPPB 247). Isolated from lettuce. ATCC 17815.
401. Strain PJ 187. Obtained by Dr H. Lautrop from the Edinburgh and East of Scotland School of Agriculture under the name *P. marginalis*. Isolated from dahlia. ATCC 17816.

- 402. Strain PJ 188. Obtained from Dr W. C. Haynes. (NRRL strain B-981.) ATCC 17817.
- 403. Strain PJ 251. Obtained from Dr K. Klinge. (Strain R-2.) ATCC 17818.
- 404. Strain PJ 283. Isolated from pleural fluid. ATCC 17819.
- 405. Strain PJ 362. Isolated from fresh water. ATCC 17820.
- 406. Strain PJ 365. Isolated from fresh water. ATCC 17821.
- 407. Strain PJ 379. Obtained from Dr J. Ørskov, Statens Seruminstitut. ATCC 17822.
- 408. Strain PJ 380. Isolated from fresh water. ATCC 17823.
- 409. Strain PJ 381. Isolated from fresh water. ATCC 17824.
- 410. Strain PJ 384. Isolated from fresh water. ATCC 17825.
- 411. Strain PJ 672. Isolated from sea water. ATCC 17826.
- 412. Strain PJ 833. Isolated from pond water. ATCC 17827.
- 413. Strain PJ 851. Isolated from tap water. ATCC 17828.
- 414. Strain PJ 883. Isolated from well. ATCC 17829.
- 415. Strain PJ 383. Isolated from fresh water. ATCC 17830.

IV. Strains assigned to *Pseudomonas fluorescens* biotype C

- 18. Strain P-7 from the Western Utilization Research and Development Division of the U.S. Department of Agriculture, Albany, California. Psychrophilic organism isolated from hen's egg and named *P. fluorescens*. ATCC 17400.
- 50. Strain P3T-3, isolated in Berkeley from L-tryptophan enrichment of water sample. ATCC 17427.

The following 13 strains were studied by Jessen (1965); his biotype and group classification is indicated in each case.

- 181. Strain PJ 70. Isolated by O. Lysenko (strain 295). Biotype 52, group IV. ATCC 948.
- 191. Strain PJ 236. Received from Dr Muriel E. Rhodes as strain 22.1. Biotype 49, group IV. ATCC 17559.
- 194. Strain PJ 253. Received from Dr K. Klinge as strain R-60. Biotype 49, group IV. ATCC 17561.
- 204. Strain PJ 682. Isolated from polluted sea water, Øresund, Denmark. Biotype 49, group IV. ATCC 17571.
- 205. Strain PJ 686. Isolated from polluted sea water, Øresund, Denmark. Biotype 49, group IV. ATCC 17572.
- 207. Strain PJ 693. Isolated from polluted sea water, Øresund, Denmark. Biotype 49, group IV. ATCC 17574.
- 210. Strain PJ 824. Isolated from ground water, from Ribe, Jutland. Biotype 49, group IV. ATCC 17577.
- 212. Strain PJ 832. Isolated from ground water, Mors, Jutland. Biotype 49, group IV. ATCC 17579.
- 213. Strain PJ 834. From polluted surface water, canal, South Copenhagen. Biotype 49, group IV. ATCC 17580.
- 214. Strain PJ 848. From polluted natural stream, South Copenhagen. Biotype 49, group IV. ATCC 17581.
- 217. Strain PJ 905. From polluted sea water, Greenland (Sukkertoppen). Biotype 49, group IV. ATCC 17584.
- 218. Strain PJ 929. Same origin as 217. Biotype 52, group IV. ATCC 17585.
- 219. Strain PJ 969. From garden soil, Arreskov, Funen. Biotype 49, group IV. ATCC 17586.

V. Strains assigned to *Pseudomonas fluorescens* biotype D

All strains of this group were received from Dr W. C. Haynes, U.S. Department of Agriculture, Peoria, Illinois, as *P. chlororaphis*.

- 30. Strain B-560. ATCC 9446.
- 31. Strain B-561. ATCC 9447.
- 32. Strain B-977. ATCC 17411.
- 35. Strain B-1541. ATCC 17414.
- 388. Strain B-2266. ATCC 17809.
- 389. Strain B-1854. ATCC 17810.
- 390. Strain B-1095. ATCC 17811.
- 391. Strain B-1869. ATCC 17812.
- 392. Strain B-2075. ATCC 17813.
- 393. Strain B-1632. ATCC 17814.
- 394. Strain B-1632. Colony variant ATCC 17814.

VI. Strains assigned to *Pseudomonas fluorescens* biotype E

All strains of this group were received from Dr W. C. Haynes, Northern Utilization Research and Development Division (U.S. Department of Agriculture, Peoria, Illinois) as *P. aureofaciens*.

- 36. Strain B-1482R. Isolated at NRRL (Peoria) from a sample of farm soil in 1953. ATCC 17415.
- 37. Strain B-1543P. Isolated at NRRL from farm soil in 1953. ATCC 17416.
- 38. Strain B-1576. Received from the late Professor A. J. Kluyver as *Pseudomonas polychromogenes*. Isolated from clay that had been suspended in kerosene for 3 weeks. ATCC 13985.
- 39. Strain B-1681. Isolated NRRL from forest soil. ATCC 17417.
- 40. Strain B-1855. Isolated at NRRL from lake water. ATCC 17418.
- 41. Strain B-2265. Isolated by Dr A. Fuchs in 1959 as *P. aureofaciens* var. *non-liquefaciens*. Air contaminant. ATCC 17419.
- 86. Strain B-1855. Isolated from a water sample from Illinois. ATCC 17461.

VII. Strains assigned to *Pseudomonas fluorescens* biotype F

- 83. *P. lemonnierii* strain B-1864 of Dr W. C. Haynes, who received it from Dr P. H. A. Sneath as strain LA. Isolated by Hugo & Turner (1957). ATCC 17458.
- 143. Strain PL, isolated in Berkeley by Miss Diana Loeb from hydrocarbon enrichment. ATCC 17513.

VIII. Strains assigned to *Pseudomonas fluorescens* biotype G

- 1. Strain SCT-1, isolated in Berkeley from a tryptophan enrichment of water sample. ATCC 17386.
- 33. *P. chlororaphis* strain B-1079 received from Dr W. C. Haynes, Peoria. ATCC 17412.
- 34. Ibid., strain B-1098. ATCC 17413.
- 99. Strain RYS-3 of R. Y. Stanier. Isolated from L-tryptophan enrichment. ATCC 17473.
- 124. Strain W4-gly-1, isolated in Berkeley from glycerol enrichment of water sample. ATCC 17496.

149. Fluorescent pseudomonad strain 2440-40 of Dr M. Starr. Isolated from soil by Dr H. Stolp. ATCC 17518.
164. Ibid., strain 2440-15. ATCC 17533.
166. Ibid., strain 2440-28. ATCC 17535.
169. Ibid., strain 2440-183. ATCC 17538.
171. Ibid., strain 2440-105. ATCC 17540.
172. Ibid., strain 2440-110. ATCC 17541.
195. Strain PJ 274 of Dr O. Jessen. Isolated from sputum, Haderslev, Jutland. Included in biotype 52, group IV, by Jessen (1965). ATCC 17562.
206. Strain PJ 692 of Dr O. Jessen. Isolated from polluted sea water, Øresund, Denmark. Included in biotype 52, group IV, by Jessen (1965). ATCC 17573.
267. *P. schuylkilliensis* strain B9 obtained from Dr M. Mandel. ATCC 17634.
269. *P. geniculata* strain B-1606, obtained from Dr M. Mandel. ATCC 17636.
271. *P. geniculata*, strain B-1612 of Dr M. Mandel. ATCC 17638.
272. Ibid., strain B-1603. ATCC 14150.

IX. Strains assigned to *Pseudomonas putida* biotype A

5. *P. putida* strain 5 of den Dooren de Jong (1926). Isolated from soil by enrichment with amylamine. Received from Laboratorium voor Microbiologie, Delft. ATCC 17390.
6. *P. putida* strain 9 of den Dooren de Jong (1926). Isolated from soil by enrichment with propionate. Same source as 5. ATCC 17391.
7. *P. putida* strain of den Dooren de Jong (1926). Isolated from soil by enrichment with phenylaminoacetate. Same source as 5. ATCC 17392.
8. *P. putida* strain 39 of den Dooren de Jong (1926). Isolated from soil by enrichment with heptanoic acid. Same source as 5. ATCC 17393.
26. Strain KA 14-1 isolated by Dr Keichii Hosokawa in Berkeley from soil by kynurenate enrichment. ATCC 17408.
42. Strain LAF-1 of Dr S. C. Rittenberg, University of California, Los Angeles. Nicotine oxidizer. ATCC 17420.
43. Strain PS-1 of Dr S. C. Rittenberg, University of California, Los Angeles. ATCC 17421.
44. Ibid., strain PS-2. ATCC 17422.
49. Strain MB-15 of Dr K. Hosokawa. Isolated from soil by benzoate enrichment. ATCC 17426.
51. Strain M-6 of Dr K. Hosokawa. Isolated from soil by anthranilate enrichment. ATCC 17428.
56. Strain W-1, isolated in Berkeley by Miss Margaret Farley from arginine enrichment from water. ATCC 17433.
76. Strain C 1-A of Dr I. C. Gunsalus, University of Illinois. Isolated from soil by camphor enrichment. ATCC 17452.
77. Strain C 1-B. Same source and origin as 76. ATCC 17453.
81. Strain S-2 of Dr E. E. Snell. Isolated from soil by canavanine enrichment. ATCC 17456.
82. Strain MS of Dr E. E. Snell. Isolated from soil by α -methyl-serine enrichment. ATCC 17457.

87. Strain A.3.1 of R. Y. Stanier. Isolated from soil by alanine enrichment. ATCC 17462.
89. Strain A.3.8 of R. Y. Stanier. Isolated from soil by malate enrichment. ATCC 17464.
90. Strain A.3.12 of R. Y. Stanier. Isolated from soil by lactate enrichment. ATCC 12633.
100. Strain received under the designation *P. ovalis* from Dr Hans Kornberg. ATCC 17474.
111. Strain 53-2 of Dr W. R. Sistrom. Isolated from soil by naphthalene enrichment. ATCC 17485.
115. Strain tyr of Dr M. Shilo. Isolated from soil by tyrosine enrichment. ATCC 17487.
118. Strain meso-1, isolated by Dr M. Shilo from soil by *meso*-tartrate enrichment. ATCC 17490.
119. Strain L-2 of Dr M. Shilo. Isolated from soil by L-tartrate enrichment. ATCC 17491.
122. Strain D-malic-2 of Dr M. Shilo. Isolated from soil by D-malic enrichment. ATCC 17494.
128. Strain MMD of Dr M. Shilo. Isolated from soil by tartrate enrichment. ATCC 17500.
130. Strain N-9 of Dr E. J. Behrman. Isolated from soil by nicotinic acid enrichment. ATCC 17502.
144. Strain B of Dr I. C. Gunsalus, University of Illinois. ATCC 17514.
145. Strain HT, isolated in 1962 by Dr S. Dagley from soil by enrichment on phenylacetate. ATCC 17515.
154. Strain 2440-112 received from Dr M. Starr, University of California, Davis. Isolated by Dr H. Stolp from soil. ATCC 17523.
160. Ibid., strain 2440-133. ATCC 17529.
266. Unidentified strain received from Dr M. J. Coon, Department of Biological Chemistry, University of Michigan, Ann Arbor, Michigan. ATCC 17633.
276. Strain isolated by Dr W. B. Jakoby from soil by tartrate enrichment. ATCC 17642.

X. Strains assigned to *Pseudomonas putida* biotype B.

53. Strain B-3 received from Dr W. C. Haynes. Originally classified as *P. aeruginosa*. ATCC 17430.
96. Strain TR-26 isolated by Dr S. H. Hutner from soil by L-tryptophan enrichment. ATCC 17470.
98. Strain RYS-2, isolated in Berkeley from soil by L-tryptophan enrichment. ATCC 17472.
107. Strain 52-19, isolated in Berkeley by Dr W. R. Sistrom from soil by saligenin enrichment. ATCC 17481.
110. Strain 53-18, isolated by Dr N. Walker at Rothamsted by naphthalene enrichment. ATCC 17484.
153. Strain 2440-103 received from Dr M. P. Starr. ATCC 17522.
157. Strain 2440-4. ATCC 17526.
158. Strain 2440-176. ATCC 17527.
167. Strain 2440-87. ATCC 17536.

XI. Unclassified strains of the fluorescent group

Eleven strains of fluorescent pseudomonads could not be assigned to any of the species or biotypes previously listed.

10. Strain 112 of den Dooren de Jong (1926), isolated from water. ATCC 17395.
17. Strain P-1 from the Western Utilization Research and Development Division of the U.S. Department of Agriculture, Albany, California. Psychrophilic fluorescent pseudomonad isolated from egg. ATCC 17399.
47. Strain F-67 of Dr E. R. Blakley (National Research Council, Canada). Isolated from a cinnamate enrichment culture. ATCC 17425.
59. Strain W-2 isolated by Miss M. Farley in Berkeley from benzoate enrichment. ATCC 17436.
91. Strain A.3.18; originally received from U.S. Department of Agriculture, Peoria, Illinois, as B-8. ATCC 17465.
92. Strain A.3.19, same source as 91; B-14. ATCC 17466.
95. Strain TR-23, isolated by Dr S. H. Hutner from soil by enrichment with tryptophan. ATCC 17469.
101. Strain NIC-24, isolated in Berkeley from soil by enrichment with nicotinate. ATCC 17475.
109. Strain 53-17 of Dr W. R. Sistrom. Isolated by Dr N. Walker at Rothamsted, by enrichment with naphthalene. ATCC 17483.
127. Strain DMM of Dr M. Shilo. Isolated in Berkeley from enrichment with tartrate. ATCC 17499.
322. Strain 62-FC of Dr M. Véron, Institut Pasteur, Paris. Isolated from mineral oil and designated as *P. denitrificans*. ATCC 17688.

General group characters

The production of water-soluble, yellow-green fluorescent pigments is frequently encountered in the aerobic pseudomonads, and has long been used as a major character in the taxonomic subdivision of the genus *Pseudomonas*. It has never been clearly established, however, whether the production of this type of pigment is correlated with other characters that would justify the recognition of the fluorescent pseudomonads as a special subgroup. This is an important problem from the determinative standpoint, since production of fluorescent pigments is strongly influenced by nutritional factors, and can fail completely on media which support excellent growth. Furthermore, it has been shown that occasional strains otherwise typical of *P. aeruginosa* (the one easily identifiable fluorescent species) may be completely unable to produce fluorescent pigment (Véron, 1961; Jessen, 1965).

Of the 175 strains studied by us, 20 did not produce the group-specific fluorescent pigment on slopes of King B medium, or on any other medium used in their examination. These strains might be capable of fluorescent pigment production under some circumstances, but we did not further pursue this question, because our analysis had by then revealed that the members of this generic subgroup share additional properties which permit their identification even when pigment production fails. We shall therefore preface the description of individual species with a discussion of the common group characters.

Fluorescent pseudomonads do not accumulate poly- β -hydroxybutyrate as a

Table 7. *Fluorescent group. The utilization of organic compounds as carbon and energy sources by fluorescent pseudomonads, based on the analysis of 175 strains*

The compounds listed are ones that are: (a) used by a majority of strains, and (b) used by a few or no strains. Accordingly, they are not diagnostically useful for the internal differentiation of species in the fluorescent group.

Compound	No. of positive strains	% of positive strains	Compound	No. of positive strains	% of positive strains
Carbohydrates and sugar derivatives			Non-nitrogenous aromatic and other cyclic compounds		
D-Arabinose	0	0	D-Mandelate	8	5
D-Fucose	0	0	Benzoate	113	65
L-Rhamnose	10	6	<i>o</i> -OH-Benzoate	12	7
D-Glucose	174	99	<i>m</i> -OH-Benzoate	7	4
D-Fructose	159	91	<i>p</i> -OH-Benzoate	167	95
Maltose	1	1	Phenylethanediol	6	3
Cellobiose	0	0	Phthalate	1	1
Salicin	0	0	Isophthalate	0	0
Lactose	0	0	Terephthalate	0	0
Starch	0	0	Naphthalene	4	2
Gluconate	174	99	Phenol	11	6
2-Ketogluconate	169	97	Quinate	167	95
Fatty acids			Testosterone	12	7
Acetate	175	100	Aliphatic amino acids		
Caproate	168	96	L- α -Alanine	174	99
Heptanoate	172	98	D- α -Alanine	172	98
Caprylate	174	99	β -Alanine	174	99
Pelargonate	175	100	L-Threonine	0	0
Caprate	174	99	L-Leucine	166	95
Dicarboxylic acids			L-Isoleucine	160	91
Malonate	160	91	L-Norleucine	0	0
Succinate	175	100	L-Valine	164	94
Maleate	3	2	L-Aspartate	175	100
Fumarate	175	100	L-Glutamate	175	100
Glutarate	170	97	L-Lysine	145	83
Hydroxyacids			DL-Arginine	173	99
L-Malate	175	100	DL-Ornithine	157	90
D-(-)-Tartrate	16	9	DL-Citrulline	132	75
DL- β -OH-Butyrate	173	99	DL- α -Aminobutyrate	4	2
DL-Lactate	175	100	γ -Aminobutyrate	174	99
Glycollate	5	3	δ -Aminovalerate	162	93
DL-Glycerate	169	97	Amino acids and related compounds containing a ring structure		
Poly- β -OH-butyrate	0	0	L-Histidine	168	96
Miscellaneous organic acids			L-Proline	173	99
Citrate	174	99	L-Tyrosine	169	97
α -Ketoglutarate	173	99	D-Tryptophan	1	1
Pyruvate	174	99	Amines		
Aconitate	162	93	Ethanolamine	125	71
Polyalcohols and glycols			Putrescine	172	98
Glycerol	173	99	Spermine	171	98
Ethylene glycol	3	2	Miscellaneous nitrogenous compounds		
2,3-Butyleneglycol	112	64	Betaine	173	99
			Sarcosine	164	94
			Pantothenate	14	8

cellular reserve material, and are unable to attack this polymer by means of extracellular enzymes. They grow well with either nitrate or ammonia as sole nitrogen source. All are oxidase-positive. We have examined only a minority of our strains for constitutive arginine dihydrolase production; but every strain examined gave positive results.

The general nutritional spectrum of these bacteria (Table 7) has many characteristic qualitative features, some of which are valuable in distinguishing them from other aerobic pseudomonads. Two hexoses, glucose and fructose, and two sugar acids, gluconate and 2-ketogluconate, are practically universal substrates. One exceptional strain (41) did grow on either glucose or gluconate; and occasional strains did not grow on either fructose or 2-ketogluconate. No strain of this group grew on D-fucose, D-arabinose, cellobiose, lactose or starch; only one strain grew on maltose, and 10 on L-rhamnose.

Among the fatty acids, acetate, propionate, caproate, heptanoate, caprylate, pelargonate and caprate were used by nearly all strains. Utilization of the C₄ and C₆ fatty acids (butyrate, isobutyrate, valerate, isovalerate) was much less common, and reflects the failure of some biotypes of *Pseudomonas fluorescens* to grow on these compounds. In the dicarboxylic acid series, malonate, succinate and glutarate were used by most strains; but the higher homologues (C₆–C₁₀) were much more rarely attacked; the figures for their utilization largely reflect the fact that they are substrates for *P. aeruginosa*. Other aliphatic organic acids used by most or all strains of the group were: fumarate, L-malate, DL-lactate, DL- β -hydroxybutyrate, DL-glycerate, citrate, α -ketoglutarate, aconitate, pyruvate. Glycollate and maleate were used by only four and three strains, respectively.

A number of simple aromatic compounds are substrates for many fluorescent pseudomonads. The most widely utilized was *p*-hydroxybenzoate, which supported growth of 167 strains. Benzoate and anthranilate were used by much smaller total numbers of strains, primarily because benzoate cannot be used by many strains of *Pseudomonas fluorescens* and anthranilate cannot be used by many strains of *P. putida*. The *o*- and *m*-monohydroxybenzoic acids were used only by occasional strains (12 and 7, respectively). The alicyclic acid, quinate, used by 167 strains, was metabolized through an aromatic pathway, since it is converted metabolically to protocatechuic acid, the first intermediate in the oxidation of *p*-hydroxybenzoate.

Among the amino acids used by most strains as carbon and energy sources were α -alanine, β -alanine, leucine, isoleucine, valine, aspartate, glutamate, arginine, ornithine, histidine, proline, tyrosine, γ -aminobutyrate, δ -aminovalerate. Two amino acids, threonine and norleucine, were not used by any strain, and α -aminobutyrate was used by only one strain. Other very commonly used nitrogenous organic compounds were the polyamines, putrescine and spermine, and the *N*-methyl compounds, betaine and sarcosine.

By selection among the common group characters described above, it is possible to construct a relatively brief list of traits which are diagnostic for fluorescent pseudomonads as a subgroup among the aerobic pseudomonads. These traits, and their frequencies in the 175 strains examined, are listed in Table 8. From the frequencies, one can construct what we term an 'ideal' phenotype, by scoring characters of high frequency as + and characters of low frequency as -. With respect to the sixteen characters which constitute the ideal phenotype, conformity

within the group is excellent: 133 strains show perfect conformity, 38 deviate in one character, and four deviate in two characters (Table 9). These characters were chosen for their value in permitting the differentiation of fluorescent pseudomonads from all the other species and species groups which we have studied. The effectiveness with which they achieve this is also shown in Table 9.

Table 8. *The group characters of greatest differential value in the recognition of the fluorescent pseudomonads, based on the analysis of 175 strains*

Characters	No. of positive strains	Ideal phenotype
1. Production of fluorescent pigment	155	+
2. Poly- β -hydroxybutyrate as cellular reserve material	0	—
3. Utilization of poly- β -hydroxybutyrate as exogenous substrate	0	—
Utilization of		
4. D-Fucose	0	—
5. D-Glucose	174	+
6. Cellobiose	0	—
7. Starch	0	—
8. 2-Ketogluconate	167	+
9. Pelargonate	175	+
10. Benzoate and/or <i>p</i> -hydroxybenzoate	172	+
11. <i>m</i> -hydroxybenzoate	7	—
12. β -Alanine	174	+
13. Norleucine	0	—
14. Arginine	173	+
15. Putrescine and/or spermine	173	+
16. Betaine and/or sarcosine	175	+

Table 9. *Fluorescent group. Number of strains of different species or groups of species of aerobic pseudomonads conforming to the selected sixteen characters (Table 8) which define the 'ideal' fluorescent pseudomonad*

	No. of characters of 'ideal' phenotype												
	16	15	14	13	12	11	10	9	8	7	6	5	4
Fluorescent group	133	38	4	.	.	.	.	.	.	.	.	.	.
Acidovorans group	.	.	.	.	.	.	.	.	1	.	13	8	4
Pseudomallei group	.	.	.	.	.	.	15	13	9	2	1	1	.
<i>P. multivorans</i>	.	.	.	.	.	19	.	.	.	.	.	.	.
<i>P. stutzeri</i>	.	.	.	.	.	.	4	12	1	.	.	.	.
<i>P. maltophilia</i>	.	.	.	.	.	.	.	.	.	23	.	.	.
Alcaligenes group	.	.	.	.	1	6	.	.	.	.	.	.	.
<i>P. lemoignei</i>	.	.	.	.	.	.	.	.	.	1	.	.	.

The three species that most closely approach the ideal phenotype of fluorescent pseudomonads are *Pseudomonas multivorans*, of which all 19 strains deviated in five of the characters chosen; *P. pseudoalcaligenes*, of which all six strains deviated in five characters; and *P. alcaligenes*, of which the single strain examined deviated in only four characters. Accordingly, the characters chosen as diagnostic for the group may be considered highly satisfactory for determinative purposes, except, possibly, in making the distinction from *P. alcaligenes*.

All major biotypes now recognized among the fluorescent pseudomonads are represented in the collection that we have studied, with the one exception of the phytopathogens. It is therefore not certain that this latter subgroup will conform in its entirety to the proposed ideal phenotype of the fluorescent group. Our fragmentary observations on the phytopathogenic pseudomonads are worth mentioning in this connexion. We have completely characterized four named strains, which were received incidentally from Jessen (1965) among representatives of his group V, biotypes 61 and 63. Two of these (our strains 187 and 188), belonging to Jessen's biotype 61, were labelled *Pseudomonas angulata* and *P. marginata*, respectively. They are typical representatives of *P. fluorescens* biotype A. The other two (our strains 400 and 401), belonging to Jessen's biotype 63, were independent isolates labelled *P. marginalis*, and are typical representatives of *P. fluorescens* biotype B. This experience suggests (and the data of Jessen, 1965, are fully concordant) that some phytopathogenic pseudomonads cannot be distinguished on any cultural, physiological or biochemical criteria from *P. fluorescens*; such organisms will obviously fit the ideal phenotype proposed above.

We also examined cursorily a small collection of phytopathogenic fluorescent strains received from Dr M. P. Starr; these included four strains labelled *Pseudomonas syringae*, two labelled *P. morsprunorum* and one labelled *P. tabaci*. These strains constitute a group of gelatin-liquefying fluorescent pseudomonads clearly distinguishable from any biotype of *P. fluorescens* with which we are familiar. Their most striking characteristic is an extremely low growth rate, not only in chemically defined media, but even in yeast extract medium. The organisms from yeast extract cultures were large, swollen and elongated. These observations suggest that the strains in question may have a partial requirement for one or more growth factors not present in yeast extract. However, they can grow (very slowly) in chemically defined media, which has permitted us to make a preliminary scan of their nutritional spectra. They seem to have a far lower degree of nutritional versatility than any other fluorescent pseudomonads. Accordingly, it is quite possible that one additional fluorescent species, consisting of these types of fluorescent plant pathogens, will eventually deserve recognition; and, in view of their apparent nutritional defects, these strains may deviate significantly from the ideal phenotype of the group that we have proposed on the basis of the study of its free-living members. It should be noted that the results of Jessen (1965) also support this conclusion. He placed most of the phytopathogenic fluorescent strains which he examined in a special group (VI), and commented on the unusually slow growth of many of them. Included in Jessen's group VI were strains labelled *P. syringae* and *P. morsprunorum*.

Internal subdivision of the fluorescent group

The only fluorescent pseudomonad now recognizable with reasonable certainty on the basis of several unrelated characters is the type species of the genus *Pseudomonas*, *P. aeruginosa*. The characters now most commonly used for its recognition are: pyocyanine production; growth at 41–43°; and production of slime in a chemically defined medium containing 2-ketogluconate (Haynes, 1951). By far the most extensive comparison of this species with other fluorescent pseudomonads has been made recently by Jessen (1965), who examined a collection of 859 fluorescent strains, of which 354 were identified as *P. aeruginosa*. Jessen has clearly demon-

strated that *P. aeruginosa* is a distinctive and internally uniform biotype. Apart from the characters already mentioned, Jessen found that the strains of this species shared the following characters: denitrifying ability; gelatin liquefaction; negative egg-yolk reaction; failure to grow at 5°; failure to produce acid from sucrose, adonitol, dulcitol, sorbitol or erythritol; acid production from mannitol; monotrichous flagellation, provided that the assessment of this character is made on a statistical basis, as suggested by Lautrop & Jessen (1964).

The classification of other fluorescent pseudomonads remains highly unsatisfactory. Three species (*Pseudomonas lemonnieri*, *P. chlororaphis*, *P. aureofaciens*) have each been defined on a single property, the ability to produce a specific phenazine pigment. Haynes & Rhodes (1962) suggested that *P. chlororaphis* closely resembles the gelatin-liquefying simple fluorescent pseudomonads. Liu (1960), in a serological study on the extracellular antigens of *Pseudomonas* strains, found that *P. aureofaciens*, *P. chlororaphis* and many strains of haemolytic lecithinase-positive *P. fluorescens* showed extensive cross-reactions, which suggested a close relationship among these species. Another important character which clearly differentiates *P. chlororaphis* and *P. aureofaciens* from *P. aeruginosa*, and at the same time justifies associating the first two species with the simple fluorescent pseudomonads, is the fact that both *P. chlororaphis* and *P. aureofaciens* show a guanine + cytosine (GC) content of 63.5 moles % in their DNA. This GC content is similar to that of some simple fluorescent pseudomonads, and markedly lower than the average of 66.5 % found for our strains of *P. aeruginosa* (Mandel, 1961). All these facts do not fit the conclusion of Lysenko (1961), who considered *P. chlororaphis* and *P. aureofaciens* to be closely allied to *P. aeruginosa*.

Among the simple (non-phenazine-producing) fluorescent pseudomonads, the two classical species, *Pseudomonas fluorescens* and *P. putida*, were originally distinguished by Flügge (1886) under the designations *B. fluorescens liquefaciens* and *B. fluorescens non-liquefaciens* on the basis of gelatin liquefaction, a character possessed by the former but not the latter. At least 20 other species (exclusive of the many species of plant pathogens) have since been named; but the descriptions are fragmentary and the alleged differences between named species are tenuous. To judge from recent studies of this group, the primary taxonomic question seems to be whether even the distinction between *P. fluorescens* and *P. putida* can be successfully maintained. Klinge & Gräf (1959) regarded it as valid, and emphasized the egg-yolk reaction as an additional property characteristic of *P. fluorescens* and not shared by *P. putida*. The data of den Dooren de Jong (1926) suggested that some nutritional characters overlooked by later workers might provide additional means for making the distinction. Rhodes (1959, 1961), on the other hand, found it impossible to make clear-cut subdivisions among 184 strains of the *P. fluorescens*-*P. putida* group.

The most thorough taxonomic analysis of the simple fluorescent pseudomonads is that of Jessen (1965). On the basis of twenty-nine diagnostic tests, Jessen found that 354 strains of *Pseudomonas aeruginosa* showed sufficient uniformity to justify their grouping in a single biotype; but application of the same tests to 505 strains of simple fluorescent pseudomonads revealed the existence of no less than 81 distinct biotypes. By clustering biotypes that differed from one another in no more than two characters, Jessen sorted the bulk of these strains into five principal groups designated as II-VI; *P. aeruginosa* constituted group I. These five groups included

Table 10. *Diagnostic characters used in the differentiation of six groups of fluorescent pseudomonads (Jessen, 1965)*

Group	Acid formation in Hugh & Lefson (1953) medium from										Levan from sucrose	Egg-yolk reaction	Growth at		
	Sucrose	Maltose	Trehalose	Raffinose	Erythritol	Adonitol	Sorbitol	Mannitol	Inositol	NO ₃ reduction	Gelatin liquef.		5°	37°	42°
I (<i>P. aeruginosa</i>)	-	-	v	-	-	-	-	+	-	+	+	-	-	+	+
II (192 strains)	v	-	v	-	-	v	-	v	-	+	v	v	+	v	-
III (26 strains)	-	+	v	-	-	-	-	v	v	-	v	-	+	-	-
IV (135 strains)	-	-	+	-	+	+	v	+	+	v	v	+	+	-	-
V (65 strains)	+	-	+	v	+	v	+	+	+	v	+	+	+	v	-
VI (22 strains)	+	-	+	v	+	-	+	+	+	-	v	-	+	-	-

Exceptions to the above patterns: *, 8 positive strains; †, 11 negative strains; ‡, one negative strain; §, three negative strains; ||, one negative strain; ¶, two negative strains.

48 biotypes, but accounted for a much larger fraction of individual strains (440). The remaining 65 strains, scattered through 33 biotypes, constituted intermediate groups. The characters that distinguish the six groups of Jessen are shown in Table 10.

The largest single biotype in Jessen's collection of simple fluorescent pseudomonads, biotype 11 of his group II, corresponds to *Pseudomonas putida*, while his groups IV and V comprised the bulk of the strains that correspond to *P. fluorescens*. His group VI consisted entirely of phytopathogenic strains, although a few phytopathogens were assigned to groups IV and V. Group III consisted of 26 strains, mostly of marine origin, which do not seem readily equatable with any named species in the fluorescent group. Jessen concluded that the extent of intergrading among the simple fluorescent pseudomonads is too great to allow the recognition of species among them.

We have examined a considerably smaller collection of strains than Jessen, but our phenotypic characterization of these strains has been more extensive than his. We have reached much the same general conclusions. Twenty-nine of the 175 fluorescent pseudomonads in our collection could be readily identified as *Pseudomonas aeruginosa* on the basis of the accepted criteria. The screening of carbon and energy sources revealed several hitherto unrecognized nutritional characters which strengthen the differentiation of this species from other fluorescent pseudomonads. In all respects, these 29 strains showed a high degree of uniformity; we therefore fully concur with Jessen that *P. aeruginosa* constitutes a relatively isolated and uniform biotype. The remaining 146 strains were far less uniform, and we agree with Jessen that their subdivision offers a difficult taxonomic problem. Although we have not fully resolved this problem, we believe that the distinction between the two classical species, *P. fluorescens* and *P. putida*, can be maintained on the basis of several different criteria; and that most of the simple fluorescent pseudomonads can be fitted into one or other of these two species. Of the 146 strains which remained after segregation of strains identified as *P. aeruginosa*, 94 could be assigned to *P. fluorescens* (including all other phenazine producers), and 41 to *P. putida*. The remaining 11 strains could not be satisfactorily assigned to either of these species, and will be separately described but not named. In effect, therefore, we propose, for the present, the recognition of only three species in the fluorescent group: *P. aeruginosa*, *P. fluorescens*, *P. putida*. Of these three species, *P. aeruginosa* is internally relatively uniform; but *P. fluorescens* and *P. putida* are not. *P. fluorescens* includes seven biotypes and *P. putida* two biotypes, which are distinguishable from one another by several different characters. Table 11 lists the characters that we believe to be of greatest value in differentiating among the three species of fluorescent pseudomonads. We shall discuss each character in turn.

Flagellar number. In agreement with Lautrop & Jessen (1964) we have found that all strains of *Pseudomonas aeruginosa* are monotrichous, whereas all our remaining strains of fluorescent pseudomonads are multitrichous.

Extracellular enzyme production. The ability to secrete extracellular hydrolytic enzymes (proteases, lecithinases, lipases) is common to *Pseudomonas aeruginosa* and *P. fluorescens*, but almost completely lacking in *P. putida*. In determining these properties we have depended (like most previous workers) on relatively crude and biochemically imprecise methods of assay, namely, tests for the hydrolysis of gelatin and of Tween 80, and for the occurrence of an egg-yolk reaction.

The production of extracellular proteases, as evidenced by the hydrolysis of gelatin, is a universal property of both *Pseudomonas aeruginosa* and *P. fluorescens*, but never occurs in *P. putida*. The egg-yolk reaction is positive only in *P. fluorescens*, though not universal in this species; 77 of the 94 strains examined gave a positive reaction. The mechanism of this reaction is not understood, but it seems to reflect the ability to produce extracellular lecithinases, since Klinge & Gräf (1959) have shown by direct analysis that egg-yolk-positive strains of *P. fluorescens* can liberate inorganic phosphate from lecithin. However, Klinge & Gräf (1959) also found that

Table 11. *Characters of differential value for the internal subdivision of the fluorescent pseudomonads*

	<i>P. aeruginosa</i> (29 strains)	<i>P. fluorescens</i> (94 strains)	<i>P. putida</i> (41 strains)
Flagellar no.	1	> 1	> 1
...	No. of positive strains		
Extracellular enzyme production:			
(a) Proteases (liquefaction of gelatin)	29	94	0
(b) Lecithinases (egg-yolk reaction)	6	77	0
(c) Lipases (hydrolysis of Tween 80)	29	58	3
Temperature relations:			
(a) Growth at 4°	0	93	20
(b) Growth at 41°	29	0	0
Denitrification	29	46	0
Production of pyocyanine	25	0	0
Nutritional properties			
Carbohydrates and sugar derivatives			
D-Xylose	0	51	11
L-Arabinose	0	64	14
D-Mannose	0	91	13
D-Galactose	0	83	7
Sucrose			
Utilization	0	66	3
Levan formation	0	62	0
Trehalose	0	93	0
Saccharate	0	75	40
Mucate	0	84	41
Dicarboxylic acids			
Adipate	29	13	0
Pimelate	25	9	0
Suberate	19	10	0
Azelate	28	11	0
Sebacate	29	13	0
Hydroxyacids			
L-(+)-Tartrate	0	9	25
hydroxymethylglutarate	0	43	0
Other organic acids			
Citraconate	0	36	7
Polyalcohols and glycols			
Erythritol	0	61	1
Sorbitol	0	53	3
meso-Inositol	0	85	0
Adonitol	0	47	1
Alcohol			
Geraniol	29	0	0

Table 11 (cont.)

	<i>P. aeruginosa</i> (29 strains)	<i>P. fluorescens</i> (94 strains)	<i>P. putida</i> (41 strains)
	No. of positive strains		
Non-nitrogenous aromatic and other cyclic compounds			
L-Mandelate	29	3	3
Phenylacetate	0	15	38
Aliphatic amino acids			
Glycine	22	0	34
DL- α -Aminovalerate	0	2	23
Amines			
Benzylamine	0	4	41
Histamine	29	32	40
Butylamine	0	17	38
α -Amylamine	0	27	38
Miscellaneous nitrogenous compounds			
Creatine	0	2	33
Hippurate	0	2	30
Acetamide	29	0	11
Trigonelline	0	34	41
Paraffin hydrocarbon			
n-Hexadecane	21	0	0

P. aeruginosa was a powerful producer of lecithinase, although it typically does not give an egg-yolk reaction. These facts suggest that a positive egg-yolk reaction may denote the ability to produce a specific type of lecithinase, other lecithinases not producing the turbidity characteristic of the egg-yolk reaction.

Lipase production, which we have detected by the hydrolysis of Tween 80, seems to be a less useful character. It is invariably positive in *Pseudomonas aeruginosa*, and was possessed by 58 out of 95 strains of *P. fluorescens*, and 3 out of 41 strains of *P. putida*.

A careful enzymological and immunological study of the extracellular hydrolases produced by fluorescent pseudomonads is much needed. It might well make the taxonomic differentiation between *Pseudomonas aeruginosa* and *P. fluorescens* on the basis of such characters far more precise and reliable than it now is.

Temperature relations. Every strain of *Pseudomonas aeruginosa* grows abundantly at 41°; no strain of *P. fluorescens* or *P. putida* can do so. No strain of *P. aeruginosa* can grow at 4°, whereas every strain of *P. fluorescens* can do so. The ability to grow at 4° was variable in *P. putida*; 20 strains did so.

Denitrification. The capacity to denitrify is universal in *Pseudomonas aeruginosa*, and always absent from *P. putida*. It is a variable character in *P. fluorescens*, found in certain biotypes, but not in others; somewhat less than half of the strains examined were denitrifiers.

Phenazine pigment production. Pyocyanine was produced on slopes of King A medium by 25 of the 29 strains identified as *Pseudomonas aeruginosa*. Strains capable of producing other phenazine pigments were assigned on the basis of overall phenotype to *P. fluorescens*. Specifically, 10 strains shown to produce chlororaphin, 7 strains shown to produce phenazine- α -carboxylic acid, and 1 strain which produced

the blue pigment characteristic of *P. lemonnieri* all fall within the general confines of *P. fluorescens*. In our opinion, the ability to produce these pigments is not a character of sufficient importance to justify the recognition of *P. chlororaphis*, *P. aureofaciens* and *P. lemonnieri* as species distinct from *P. fluorescens*, which these organisms resemble in other respects. None of the strains of *P. putida* was observed to produce phenazine pigments.

Nutritional patterns. Strains of *Pseudomonas fluorescens* characteristically utilize an extensive range of carbohydrates and polyalcohols, whereas strains of *P. aeruginosa* and most strains of *P. putida* do not. For purposes of differentiation, the two most valuable compounds of this class appear to be trehalose and *meso*-inositol. No strain of *P. aeruginosa* or *P. putida* utilized either compound; trehalose was utilized by 93 out of 94 strains of *P. fluorescens*, and inositol by 85 strains. Sucrose utilization is of considerably less value; although sucrose is never used by *P. aeruginosa*, it was used by only 65 out of 94 strains of *P. fluorescens*, and by four strains out of 41 of *P. putida*. *Pseudomonas fluorescens* characteristically attack this sugar by means of levan sucrase (Fuchs, 1959); only five of these Sucrose-positive strains examined by us did not produce levan from sucrose. The rare sucrose-positive strains of *P. putida*, on the other hand, were not levan-formers.

Two very valuable nutritional characters for the recognition of *Pseudomonas aeruginosa* are the ability to grow on the long-chain alcohol geraniol, possessed by all strains; and the ability to grow on one of two aliphatic hydrocarbons, dodecane or hexadecane, possessed by 25 of the 29 strains tested. None of these three compounds can be used by any strain of *P. fluorescens* or *P. putida*.

Pseudomonas putida and *P. aeruginosa* characteristically grow on the complete series of fatty acids from acetate to caprate (C_2 to C_{10}). However, some biotypes of *P. fluorescens* grew very poorly on the *n* and iso C_4 and C_5 fatty acids; in fact, failure to grow on two or more of this group of four fatty acids is practically diagnostic for *P. fluorescens*. This character is, however, not infallible for the recognition of *P. fluorescens*, since three of the biotypes included (biotypes B, D, E) did not share it.

Good growth on the higher homologues of the dicarboxylic acid series (C_6 to C_{10}) is characteristic for *Pseudomonas aeruginosa*. Only two strains of *P. putida* attacked any of these acids, but their growth was very poor. Most strains of *P. fluorescens* likewise did not grow on them, with the exception of some strains belonging to biotype C.

Among the polyalcohols, erythritol, sorbitol, *meso*-inositol and adonitol are never used by *Pseudomonas aeruginosa* and only by exceptional strains of *P. putida*. These compounds were, on the other hand, attacked by at least half of the *P. fluorescens* strains.

The aliphatic alcohols, ethanol, propanol, butanol and isobutanol, are universal substrates for *Pseudomonas aeruginosa* and almost universal for *P. putida*. However, three of the biotypes of *P. fluorescens* (A, E, F) failed completely to grow on ethanol, *n*-propanol and isobutanol. Consequently, the failure to use the alcohols is a sufficient, but not a necessary, criterion for the recognition of *P. fluorescens*.

All strains of *Pseudomonas aeruginosa* grew on L-mandelate, an aromatic substrate only used rarely by *P. fluorescens* (3 strains) and *P. putida* (3 strains). Phenylacetate is an aromatic substrate that is used by most strains of *P. putida*, but never

attacked by *P. aeruginosa*. Most biotypes of *P. fluorescens* are likewise characteristically phenylacetate negative, with the exceptions of biotypes D and E.

Among the amino acids, glycine is never utilized by *Pseudomonas fluorescens*, but can be used by a large majority of the strains of both *P. aeruginosa* and *P. putida*. α -Aminovalerate was used by 23 strains of *P. putida* but by no strain of *P. aeruginosa* or *P. fluorescens*.

Table 12. *Selected characters for the differentiation of the species of fluorescent pseudomonads*

A. Characters for the differentiation between <i>Pseudomonas aeruginosa</i> and <i>P. fluorescens</i>			
	<i>P. aeruginosa</i>	<i>P. fluorescens</i>	
1. Monotrichous flagellation	+	—	
2. Pyocyanine production	+	—	
3. Growth at 4°	—	+	
4. Growth at 41°	+	—	
5. Egg-yolk reaction	—	+	
6. Levan formation from sucrose	—	+	
7. Utilization of trehalose	—	+	
8. Utilization of inositol	—	+	
9. Utilization of geraniol	+	—	
10. Utilization of acetamide	+	—	
B. Characters for the differentiation between <i>P. aeruginosa</i> and <i>P. putida</i>			
	<i>P. aeruginosa</i>	<i>P. putida</i>	
1. Monotrichous flagellation	+	—	
2. Pyocyanine production	+	—	
3. Growth at 41°	+	—	
4. Liquefaction of gelatin	+	—	
5. Denitrification	+	—	
6. Utilization of geraniol	+	—	
7. Utilization of phenylacetate	—	+	
8. Utilization of benzylamine	—	+	
9. Utilization of creatine	—	+	
10. Utilization of hippurate	—	+	
C. Characters for the differentiation between <i>P. fluorescens</i> and <i>P. putida</i>			
	<i>P. fluorescens</i>	<i>P. putida</i>	
1. Gelatin liquefaction	+	—	
2. Egg-yolk reaction	+	—	
3. Utilization of trehalose	+	—	
4. Utilization of inositol	+	—	
5. Utilization of two or more of the following nitrogenous compounds: benzylamine, creatine, hippurate and glycine	—	+	

Pseudomonas putida characteristically grows well on several higher amines (benzylamine, butylamine, α -amylamine) that are not used by *P. aeruginosa*, or by the overwhelming majority of strains of *P. fluorescens*. By far the most useful compound of this class for differential purposes is benzylamine, used by all strains of *P. putida*, and by only four strains of *P. fluorescens*. Creatine and hippurate are two other nitrogenous organic compounds characteristically utilized by *P. putida*, but not by *P. aeruginosa* or *P. fluorescens*. No strain of *P. aeruginosa* attacked either creatine or hippurate; only two strains of *P. fluorescens* attacked each of these

compounds. However, their utilization by *P. putida* is not universal: 33 strains used creatine and 30 hippurate. A nutritional character which has been proposed as specific for *P. aeruginosa* is the ability to grow on acetamide (Bühlmann, Vischer & Bruhin, 1961). All 29 strains tested possessed this property. However, it was also possessed by 11 of the 41 strains of *P. putida*, though by no strain of *P. fluorescens*.

The alkaloid trigonelline (a derivative of nicotinic acid) is a universal substrate for *Pseudomonas putida* and never utilized by *P. aeruginosa*. Two biotypes of *P. fluorescens* (B, G) grow well on it; the remainder are negative, with the exception of four strains of biotype A.

Fig. 3. The differentiation between *Pseudomonas aeruginosa* and *P. fluorescens*, based on a group of ten selected contrasting characters. ▨, *P. aeruginosa*; ▩, *P. fluorescens* (all biotypes); ■, *P. fluorescens* (phenazine-producing biotypes).

From our complete data on the properties of the three fluorescent species, it is possible to select relatively small constellations of characters which permit unambiguous separations between three species (Table 12). A set of ten characters serves to differentiate between *Pseudomonas aeruginosa* and *P. fluorescens*; another set of ten characters differentiates between *P. aeruginosa* and *P. putida*; and a set of five characters differentiates between *P. fluorescens* and *P. putida*. The sharpness with which these selected characters permit interspecific differentiation is shown graphically in Figs. 3–5.

In Fig. 3 all the strains of *Pseudomonas aeruginosa* and of all the biotypes of *P. fluorescens* have been scored on a percentile basis for the extent of their conformity to the ten selected differential characters possessed by *P. fluorescens*. It can be seen that almost 70% of the strains of *P. aeruginosa* possessed none of these characters, while slightly more than 30% possessed one of them. Ninety-three per cent of the strains of *P. fluorescens* possessed either nine or ten of these characters; the remaining 7% possessed either seven or eight. We have calculated separately the degree of conformity of all phenazine-producing strains assigned to

Fig. 4

Fig. 4. The differentiation between *Pseudomonas aeruginosa* and *P. putida*, based on a group of ten selected contrasting characters. ▨, *P. aeruginosa*; ▩, *P. putida* (both biotypes).

Fig. 5

Fig. 5. The differentiation between *Pseudomonas fluorescens* and *P. putida*, based on a group of five selected contrasting characters. ▨, *P. fluorescens* (all biotypes); ▩, *P. putida* (both biotypes).

P. fluorescens (biotypes D, E, F); these data are also shown in Fig. 3 as narrow black bars. Ninety-five per cent of the phenazine-producing strains conformed to all ten characters of *P. fluorescens* and 5% to nine of them. It is accordingly easy to distinguish these phenazine-producing fluorescent pseudomonads from *P. aeruginosa* on the basis of the characters here used.

The extent of conformity of all strains of *Pseudomonas aeruginosa* and *P. putida*

(all biotypes) to the ten selected characters distinctive of *P. putida* is shown in Fig. 4. Again, the separation can be considered highly satisfactory. Figure 5 shows the extent of conformity of all strains of *P. fluorescens* and *P. putida* to the five characters distinctive of *P. putida*. Although the separation in this case is adequate for determinative purposes, it is less clear-cut than in the two cases previously considered. All strains of *P. putida* show perfect conformity with respect to these five characters. Almost 75 % of the strains of *P. fluorescens* show no conformity; but 22 % conform with respect to one character and 3 % with respect to two characters.

Pseudomonas aeruginosa

The properties of *Pseudomonas aeruginosa* which are valuable for its differentiation from other fluorescent pseudomonads have already been discussed. Accordingly, in this section we shall describe, without further comment, the detailed phenotypic traits of the 29 strains that we have examined.

Flagellation: monotrichous in all.

Pyocyanine production: all except strains 57, 280, 281, 283, 284, 285, 286, 416.

Fluorescent pigment: all except strains 57, 280, 281, 283, 284, 285, 286, 416.

Strains 281 and 283 to 295, inclusive, also produced on King B medium slopes a brownish red diffusible pigment which became very dark in time, and masked the presence of other pigments.

Denitrification: vigorous in all.

Gelatin liquefaction: positive in all.

Egg-yolk reaction: negative in all.

Hydrolysis of Tween 80: positive in all, but weak and delayed. In Table 13, we have listed those carbon and energy sources used by 90 % or more of the strains. Table 14 lists substrates that were not universally used, and indicates specifically the negative strains.

Pseudomonas fluorescens

The characters which we propose for the differentiation of *Pseudomonas fluorescens* from *P. putida* and *P. aeruginosa* have already been discussed, and we shall therefore introduce the detailed description of this species by a general survey of its seven biotypes, and of the properties which serve for their recognition. The principal properties which we have found to be of value for this purpose are: phenazine pigment production; denitrification; levan formation from sucrose; utilization of L-arabinose, D-xylose, saccharate, certain fatty acids, hydroxymethylglutarate, sorbitol, adonitol, certain glycols and alcohols, benzoylformate, phenylacetate, butylamine, trigonelline (Table 15). The possibility of subdividing the gelatin-liquefying fluorescent pseudomonads (excluding *P. aeruginosa*) on the basis of levan formation and denitrification has been previously suggested (e.g. by Fuchs, 1959); it is accordingly gratifying to find that a primary subdivision on these characters is correlated with differences in many other unrelated nutritional properties. The ability to produce specific phenazine pigments, heretofore used as the sole justification for the recognition of three gelatin-liquefying species, *P. chlororaphis* (= our biotype D), *P. aureofaciens* (= our biotype E) and *P. lemonnieri* (= our biotype F) is also a significant internal marker in *P. fluorescens*, since it is correlated with a number of nutritional characters. However, these three biotypes

show such high overall phenotypic resemblances to non-phenazine-producing strains of *P. fluorescens* that in our opinion it is unjustified to accord them more than varietal status.

Table 13. *Pseudomonas aeruginosa*. Substrates used by 90% or more of the strains

Carbohydrates and sugar derivatives	Alcohols
D-Ribose	Ethanol
D-Glucose	n-Propanol
D-Fructose*	n-Butanol
Gluconate	Isobutanol
2-Ketogluconate	Geraniol
Fatty acids	Non-nitrogenous aromatic and other cyclic compounds
Acetate	L-Mandelate
Propionate	Benzoylformate*
Butyrate	Benzoate
Isobutyrate	p-Hydroxybenzoate*
Valerate	Quinate*
Isovalerate	Aliphatic amino acids
Caproate	L- α -Alanine
Heptanoate	D- α -Alanine
Caprylate	β -Alanine
Pelargonate	L-Leucine*
Caprate	L-Aspartate
Dicarboxylic acids	L-Glutamate
Malonate	L-Lysine*
Succinate	DL-Arginine
Fumarate	DL-Ornithine
Glutarate	γ -Aminobutyrate
Adipate	δ -Aminovalerate
Azelate*	Amino acids and related compounds containing a ring structure
Sebacate	L-Histidine
Hydroxyacids	L-Proline
L-Malate	L-Tyrosine*
DL- β -Hydroxybutyrate	L-Kynurenine
DL-Lactate	Anthranilate
DL-Glycerate	Amines
Miscellaneous organic acids	Putrescine
Citrate	Spermine
α -Ketoglutarate	Histamine
Pyruvate	Miscellaneous nitrogenous compounds
Aconitate*	Betaine
Laevulinate	Sarcosine*
Itaconate	Acetamide
Mesaconate	Paraffin hydrocarbons
Polyalcohols and glycols	(None)
Mannitol	
Glycerol	
Propyleneglycol*	
2,3-Butyleneglycol*	

* Substrate not utilized by all strains. Negative strains are shown in Table 14.

The properties of specific phenazine pigment production, levan formation from sucrose and denitrification enable us to define six biotypes (A-F) which are positive for at least one of these characters, and which are further differentiable with a

considerable degree of sharpness by nutritional characters. These biotypes encompass 77 of the 94 strains that we assign to *Pseudomonas fluorescens*, leaving a residue of 17 strains which have the general characteristics of the species, but are negative for all the primary characters used to distinguish the other six biotypes. These we have assembled in a final biotype, G, which is far more heterogeneous, as shown by the irregularity of the nutritional data. Biotype G is frankly a provisional group, which lacks the uniformity characteristic of the other six biotypes.

Table 14. *Pseudomonas aeruginosa*. Substrates utilized by a variable number of the strains

Substrate	Total positive strains	Negative strains
D-Fructose	28	290
Pimelate	25	279, 284, 289, 296
Suberate	19	132, 279, 281, 284, 286, 288, 289, 290, 294, 296
Azelate	28	52
D-Malate	8	All except 282, 292, 293
Aconitate	28	416
Propyleneglycol	28	57
2,3-Butyleneglycol	27	45, 416
Benzoylformate	28	279
<i>p</i> -Hydroxybenzoate	28	52
Quinate	26	52, 282, 290
Glycine	22	52, 55, 57, 131, 290, 291, 294
L-Serine	4	All except 281, 282, 293, 416
L-Leucine	28	291
L-Isoleucine	15	132, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290
L-Valine	25	54, 52, 131, 132
L-Lysine	26	52, 54, 55
L-Citrulline	22	52, 54, 55, 58, 131, 132, 416
L-Tyrosine	28	57
L-Phenylalanine	7	All except 45, 54, 55, 57, 132, 279, 293
L-Tryptophan	21	279, 280, 284, 289, 290, 294, 296, 415
Kynurenate	13	54, 58, 132, 277, 278, 279, 280, 281, 283, 287, 288, 289, 290, 292, 295, 296
Ethanolamine	19	277, 279, 280, 281, 285, 286, 287, 291, 292, 293
Sarcosine	26	288, 289, 290
Dodecane	4	All except 45, 58, 131, 132
Hexadecane	21	55, 57, 278, 287, 294, 295, 296

Biotype A. With respect to the general characters of the species, biotype A is typical in almost all respects, being uniformly positive for fluorescent pigment production, gelatin liquefaction, egg-yolk reaction, and growth at 4°. One strain (12) did not hydrolyse Tween 80. Substrates used by 90% or more of the strains are listed in Table 16; and substrates used by only a fraction of the strains in Table 17, which also lists specifically the negative strains.

Biotype B. Not all strains of biotype B produce fluorescent pigment on King B medium; negative for this character were strains 400, 402, 405, 409, 413, 414. All grow at 4°. All liquefy gelatin, but do not hydrolyse Tween 80, and characteristically give a weak or negative egg-yolk reaction; only strains 2, 400, 402, 403, 411, 414 and

Table 15. *Pseudomonas fluorescens*. Characters which serve to differentiate seven biotypes among the 94 strains

	No. of positive strains (percentages in parentheses)						
	Biotype A (24 strains)	Biotype B (19 strains)	Biotype C (15 strains)	Biotype D (10 strains)	Biotype E (7 strains)	Biotype F (2 strains)	Biotype G (17 strains)
Production of							
Chlororaphin	0 (0)	0 (0)	0 (0)	10 (100)	0 (0)	0 (0)	0 (0)
Phenazine- α -carboxylic acid	0 (0)	0 (0)	0 (0)	0 (0)	7 (100)	0 (0)	0 (0)
Insoluble blue phenazine pigment	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	1 (50)	0 (0)
Levan formation from sucrose	24 (100)	19 (100)	0 (0)	10 (100)	7 (100)	2 (100)	0 (0)
Denitrification	0 (0)	19 (100)	15 (100)	10 (100)	0 (0)	2 (100)	0 (0)
Utilization of							
L-Arabinose	23 (96)	18 (95)	0 (0)	0 (0)	7 (100)	2 (100)	14 (82)
D-Xylose	22 (92)	15 (79)	0 (0)	0 (0)	0 (0)	1 (50)	13 (76)
Saccharate	24 (100)	18 (95)	0 (0)	10 (100)	7 (100)	2 (100)	14 (82)
Propionate	22 (92)	19 (100)	13 (87)	9 (90)	7 (100)	0 (0)	14 (82)
Butyrate	0 (0)	15 (79)	0 (0)	9 (90)	4 (57)	2 (100)	6 (35)
Isobutyrate	0 (0)	17 (89)	0 (0)	0 (0)	1 (14)	0 (0)	1 (6)
Valerate	6 (25)	17 (89)	9 (60)	10 (100)	7 (100)	0 (0)	8 (47)
Hydroxymethylglutarate	21 (87)	17 (89)	0 (0)	0 (0)	0 (0)	0 (0)	5 (29)
Sorbitol	22 (92)	19 (100)	2 (13)	0 (0)	0 (0)	2 (100)	8 (47)
Adonitol	22 (92)	2 (11)	15 (100)	0 (0)	0 (0)	0 (0)	8 (47)
Propylene glycol	0 (0)	18 (95)	12 (80)	0 (0)	0 (0)	0 (0)	8 (47)
Ethanol	0 (0)	18 (95)	12 (80)	10 (100)	0 (0)	0 (0)	8 (47)
n-Propanol	0 (0)	18 (95)	12 (80)	10 (100)	0 (0)	0 (0)	8 (47)
Benzoylformate	0 (0)	1 (5)	0 (0)	10 (100)	7 (100)	0 (0)	3 (18)
Phenylacetate	0 (0)	0 (0)	0 (0)	6 (60)	7 (100)	0 (0)	2 (12)
Butylamine	0 (0)	1 (5)	0 (0)	8 (80)	5 (71)	0 (0)	3 (18)
Trigonelline	3 (12)	16 (84)	0 (0)	0 (0)	0 (0)	0 (0)	15 (88)

415 are positive for this latter character. They share with the strains of biotype A the property of levan formation from sucrose, but differ in being, without exception, vigorous denitrifiers. As shown in Table 15, most strains of biotypes A and B share a nutritional property otherwise absent from *Pseudomonas fluorescens*,

Table 16. *Pseudomonas fluorescens* biotype A. Substrates used by 90% or more of the strains

Carbohydrates and sugar derivatives	Polyalcohols and glycols
D-Ribose	Mannitol
D-Xylose*	Sorbitol*
L-Arabinose*	Adonitol*
D-Glucose	Glycerol
D-Mannose	Alcohols
D-Fructose	(None)
Sucrose	Non-nitrogenous aromatic and other
Trehalose	cyclic compounds
Gluconate	<i>p</i> -Hydroxybenzoate*
2-Ketogluconate	Quinate
Saccharate	Aliphatic amino acids
Mucate	L- α -Alanine
Fatty acids	D- α -Alanine
Acetate	β -Alanine
Propionate*	L-Serine*
Caproate	L-Leucine
Heptanoate	L-Isoleucine
Caprylate	L-Valine*
Pelargonate	L-Aspartate
Caprate	L-Glutamate
Dicarboxylic acids	L-Lysine*
Malonate	DL-Arginine
Succinate	DL-Ornithine*
Fumarate	γ -Aminobutyrate
Glutarate	Amino acids and related compounds
Hydroxyacids	containing a ring structure
L-Malate	L-Histidine
DL- β -Hydroxybutyrate	L-Proline*
DL-Lactate	L-Tyrosine
DL-Glycerate*	L-Tryptophan*
Miscellaneous organic acids	Amines
Citrate	Ethanolamine*
α -Ketoglutarate	Putrescine*
Pyruvate	Spermine*
Aconitate	Miscellaneous nitrogenous compounds
Itaconate*	Betaine
Mesaconate*	Sarcosine
	Paraffin hydrocarbons
	(None)

* Substrate not utilized by all strains. Negative strains are shown in Table 17.

namely, ability to grow on hydroxymethylglutarate. The most significant differences in nutritional respects between biotypes A and B concern the utilization of certain fatty acids and alcohols. Most strains of biotype B grow well on butyrate and isobutyrate, propyleneglycol, ethanol and *n*-propanol, whereas none of the strains of biotype A can do so. Another notable feature of biotype B is the high

frequency of utilization of trigonelline, otherwise attacked by only three strains of *P. fluorescens* (all in biotype A). Substrates used by 90 % or more of the strains of biotype B are shown in Table 18; and substrates used by only a fraction of the strains are shown in Table 19, which also lists specifically the negative strains.

Table 17. *Pseudomonas fluorescens* biotype A. Substrates utilized only by a fraction of the strains

Substrate	Total of positive strains	Negative strains
D-Xylose	22	184, 185
L-Arabinose	23	12
Propionate	22	182, 186
Valerate	6	All except 184, 185, 201, 202, 209, 216
Isovalerate	20	12, 126, 211, 392
DL-Glycerate	23	208
Hydroxymethylglutarate	21	184, 185, 198
Laevulinate	6	All except 186, 187, 196, 199, 200, 201
Citraconate	13	126, 184, 185, 201, 202, 203, 209, 215, 216, 392
Itaconate	22	184, 185
Mesaconate	22	184, 185
Erythritol	20	12, 184, 185, 216
Sorbitol	22	184, 185
Inositol	21	184, 185, 216
Adonitol	22	184, 185
2,3-Butyleneglycol	6	All except 187, 189, 198, 199, 203, 215
n-Butanol	4	All except 185, 192, 196, 208
Benzoate	8	All except 126, 188, 189, 196, 199, 201, 209, 216
o-Hydroxybenzoate	1	All except 187
p-Hydroxybenzoate	23	189
L-Serine	23	12
L-Valine	23	392
L-Lysine	22	126, 392
DL-Ornithine	22	184, 185
DL-Citrulline	20	184, 185, 192, 211
δ -Aminovalerate	19	126, 208, 209, 216, 392
L-Proline	23	12
L-Phenylalanine	4	All except 12, 184, 185, 192
L-Tryptophan	22	184, 185
L-Kynurenine	20	184, 185, 208, 216
Kynurenate	12	12, 184, 185, 192, 196, 200, 202, 203, 208, 211, 216, 392
Anthranilate	19	12, 184, 185, 192, 201
Ethanolamine	23	392
Putrescine	22	182, 392
Spermine	23	182
Histamine	13	12, 126, 186, 187, 188, 192, 196, 198, 208, 211, 216
Tryptamine	2	All except 184, 185
Creatine	1	All except 208
Trigonelline	3	All except 184, 185, 215

Biotype C. The strains of biotype C all produce fluorescent pigment and grow at 4°. They all liquefy gelatin and give the egg-yolk reaction. Only a minority can hydrolyse Tween 80, the positive strains being 191, 194, 204, 207, 210, 212. All strains of this biotype are denitrifiers. Unlike the strains of biotypes A and B, they

are unable to grow on sucrose, or to synthesize levan from it. Biotype C also differs from most strains of biotypes A and B by its total inability to grow on L-arabinose, D-xylose or saccharate. It shares with biotype A the inability to grow on butyrate and isobutyrate; but most strains do grow well on propyleneglycol, ethanol and

Table 18. *Pseudomonas fluorescens* biotype B. Substrates utilized by 90% or more of the strains

Carbohydrates and sugar derivatives	Polyalcohols and glycols
D-Ribose	Mannitol
L-Arabinose*	Sorbitol
D-Glucose	Inositol
D-Mannose	Glycerol
D-Galactose	Propyleneglycol*
Sucrose	2,3-Butyleneglycol
Trehalose	Alcohols
Gluconate	Ethanol*
2-Ketogluconate	n-Propanol*
Saccharate*	Isobutanol*
Mucate*	Non-nitrogenous aromatic and other cyclic compounds (None)
Fatty acids	Aliphatic amino acids
Acetate	L- α -Alanine
Propionate	D- α -Alanine
Valerate*	β -Alanine
Isovalerate*	L-Isoleucine
Caproate*	L-Valine
Heptanoate	L-Aspartate
Caprylate	L-Glutamate
Pelargonate	DL-Arginine
Caprate	γ -Aminobutyrate
Dicarboxylic acids	δ -Aminovalerate
Malonate	Amino acids and related compounds containing a ring structure
Succinate	L-Proline
Fumarate	L-Tyrosine
Glutarate*	Amines
Hydroxyacids	Putrescine
L-Malate	Spermine
DL- β -Hydroxybutyrate	Miscellaneous nitrogenous compounds
DL-Lactate	Betaine
DL-Glycerate*	Paraffin hydrocarbons
Hydroxymethylglutarate*	(None)
Miscellaneous organic acids	
Citrate	
α -Ketoglutarate	
Pyruvate	
Aconitate	

* Substrate not used by all strains. Negative strains shown in Table 19.

n-propanol, characters absent from biotype A. Nine of the 15 strains of biotype C share another set of nutritional characters unusual among fluorescent pseudomonads, with the exception of *Pseudomonas aeruginosa*; they grow well and uniformly on the series of dicarboxylic acids from adipate to sebacate. This is one of the characters which den Dooren de Jong believed to distinguish *P. acidovorans* from fluorescent pseudomonads. It is indeed a very nearly universal property in the acidovorans group; but its presence in *P. aeruginosa* and *P. fluorescens* biotype C destroys its value as a differential character for distinguishing the fluorescent

Table 19. *Pseudomonas fluorescens* biotype B. Substrates utilized by only a fraction of the strains

Substrate	Total of positive strains	Negative strains
D-Xylose	15	108, 400, 402, 409
L-Arabinose	18	409
L-Rhamnose	4	All except 401, 403, 414, 415
D-Fructose	13	404, 405, 406, 408, 410, 412
Saccharate	18	409
Mucate	18	410
Butyrate	15	2, 93, 400, 402
Isobutyrate	1	2, 93
Valerate	17	2, 93
Isovalerate	17	2, 93
Caproate	17	2, 93
Glutarate	17	400, 402
D-Malate	6	All except 403, 407, 409, 413, 414, 415
D-(—)-Tartrate	4	All except 401, 403, 411, 414
meso-Tartrate	1	All except 413
Glycollate	1	All except 93
DL-Glycerate	17	400, 402
Hydroxymethylglutarate	17	93, 413
Laevulinate	1	All except 402
Citraconate	6	All except 93, 401, 403, 411, 414, 415
Itaconate	14	2, 407, 412, 413, 415
Mesaconate	13	2, 93, 407, 412, 413, 415
Erythritol	14	2, 93, 405, 407, 413
Adonitol	2	All except 2, 415
Ethyleneglycol	1	All except 93
Propyleneglycol	18	2
Ethanol	18	93
n-Propanol	18	93
n-Butanol	14	2, 93, 400, 402, 411
Isobutanol	17	2, 93
L-Mandelate	1	All except 93
Benzoylformate	1	All except 93
Benzoate	3	All except 93, 407, 413
L-Serine	14	2, 93, 108, 405, 412
L-Leucine	14	400, 402, 405, 406, 408
L-Valine	17	400, 402
L-Lysine	11	2, 108, 400, 402, 404, 405, 407, 412
L-Arginine	17	400, 402
DL-Ornithine	14	2, 400, 402, 407, 413
DL-Citrulline	12	2, 401, 403, 407, 413, 414, 415
L-Histidine	15	405, 406, 409, 412
L-Phenylalanine	7	All except 2, 93, 400, 401, 402, 407, 413
L-Tryptophan	14	400, 401, 403, 407, 413
L-Kynurenine	15	401, 403, 407, 413
Anthranilate	9	All except 2, 93, 406, 408, 410, 411, 412, 414, 415
Ethanolamine	10	2, 400, 403, 404, 405, 406, 408, 409, 410
Histamine	1	All except 413
Tryptamine	2	All except 108, 415
Butylamine	1	All except 413
α-Amylamine	4	All except 2, 93, 108, 413
Sarcosine	14	405, 406, 408, 410, 412
Trigonelline	16	408, 409, 413

group from the acidovorans group. The fluorescent strain described recently by Janota-Bassalik & Wright (1964), which was capable of growing on higher dicarboxylic acids and of denitrifying, is obviously a representative of *P. fluorescens* biotype C.

Substrates used by 90 % or more of the strains of biotype C are shown in Table 20;

Table 20. *Pseudomonas fluorescens* biotype C. Substrates used by 90% or more of the strains

Carbohydrates and sugar derivatives	Non-nitrogenous aromatic and other cyclic compounds
D-Ribose	<i>p</i> -Hydroxybenzoate*
D-Glucose	Quinate*
D-Mannose	Aliphatic amino acids
D-Galactose*	L- α -Alanine
D-Fructose*	D- α -Alanine
Trehalose	β -Alanine
Gluconate	L-Serine
2-Ketogluconate	L-Leucine
Fatty acids	L-Isoleucine
Acetate	L-Valine
Propionate*	L-Aspartate
Caproate*	L-Glutamate
Heptanoate	L-Lysine
Caprylate	DL-Arginine
Pelargonate	DL-Ornithine
Caprate	DL-Citrulline
Dicarboxylic acids	γ -Aminobutyrate
Malonate	δ -Aminovalerate
Succinate	Amino acids and related compounds containing a ring structure
Fumarate	L-Histidine
Glutarate	L-Proline
Hydroxyacids	L-Tyrosine
L-Malate	L-Tryptophan
DL- β -Hydroxybutyrate	L-Kynurenine
DL-Lactate	Kynurenate
DL-Glycerate	Anthranilate
Miscellaneous organic acids	Amines
Citrate	Ethanolamine
α -Ketoglutarate	Putrescine
Pyruvate	Spermine
Polyalcohols and glycols	Miscellaneous nitrogenous compounds
Erythritol	Betaine
Mannitol	Sarcosine
<i>meso</i> -Inositol	Pantothenate
Adonitol	Paraffin hydrocarbons
Glycerol	(None)
Alcohols	
(None)	

* Substrates not used by all strains. Negative strains are shown in Table 21.

and substrates used by only a fraction of the strains in Table 21, which also lists specifically the negative strains.

Biotype D. Biotype D comprises all save three of the strains which we received with the specific designation *Pseudomonas chlororaphis*; the exceptions are strain 392, which is a typical representative of biotype A; and 33 and 34, which we have

assigned to biotype G. In none of these three strains could we observe production of chlororaphin.

All the strains which we assign to biotype D produced chlororaphin on King A medium, and all save one (strain 393) produced fluorescent pigment on King B medium. All grow at 4°. All liquefy gelatin, give the egg-yolk reaction and hydrolyse

Table 21. *Pseudomonas fluorescens* biotype C. Substrates used by only a fraction of the strains

Substrate	Total of positive strains	Negative strains
L-Rhamnose	4	All except 191, 194, 205, 207
D-Galactose	14	204
D-Fructose	14	204
Mucate	10	18, 50, 204, 210, 212
Propionate	13	181, 194
Valerate	9	18, 181, 191, 194, 204, 207
Isovalerate	9	18, 50, 191, 194, 204, 212
Caproate	13	18, 50
Maleate	8	All except 204, 210, 212
Adipate	9	181, 213, 214, 217, 217, 218, 219
Pimelate	9	181, 213, 214, 218, 219
Suberate	9	181, 213, 214, 217, 218, 219
Azelate	9	181, 213, 214, 217, 218, 219
Sebacate	9	181, 213, 214, 217, 218, 219
Aconitate	12	204, 210, 212
Laevulinate	3	All except 181, 191, 194
Citraconate	6	All except 18, 181, 191, 194, 205, 207
Itaconate	6	All except 181, 191, 194, 204, 210, 212
Mesaconate	6	All except 181, 191, 204, 210, 212
Sorbitol	2	All except 181, 218
Propyleneglycol	12	204, 210, 212
2,3-Butyleneglycol	9	181, 191, 194, 204, 210, 212
Ethanol	12	204, 210, 212
n-Propanol	12	204, 210, 212
n-Butanol	10	50, 204, 207, 210, 212
Isobutanol	12	204, 210, 212
Benzoate	6	All except 18, 204, 213, 214, 217, 219
m-Hydroxybenzoate	1	All except 210
p-Hydroxybenzoate	14	212
Quinate	14	50
L-Lysine	14	50
L-Tyrosine	14	212
L-Phenylalanine	3	All except 181, 194, 214
Kynurenate	13	50, 210
Anthranilate	13	18, 212
Benzylamine	1	All except 210
Tryptamine	5	All except 191, 194, 204, 210, 212
α-Amylamine	5	All except 191, 194, 204, 210, 212
Sarcosine	14	212
Creatine	1	All except 210
Hippurate	2	All except 18, 181
Pantothenate	13	18, 50

Tween 80. All produce levan from sucrose and denitrify. The properties of levan formation and denitrification are shared with biotype B, from which biotype D is, however, differentiable by several nutritional characters as well as by chlororaphin

production. No strain of biotype D can use L-arabinose, D-xylose, isobutyrate, hydroxymethylglutarate, sorbitol, propyleneglycol or trigonelline; most strains of biotype D use benzoylformate, phenylacetate and butylamine, which are rarely or never used by biotype B (see Table 15).

Table 22. *Pseudomonas fluorescens* biotype D. Substrates used by 90% or more of the strains

Carbohydrates and sugar derivatives	Alcohols
D-Ribose	(None)
D-Glucose	Non-nitrogenous aromatic and other cyclic compounds
D-Mannose	Benzoylformate
Sucrose	Benzoate
Trehalose	<i>p</i> -Hydroxybenzoate
Gluconate	Quinate
2-Ketogluconate	Aliphatic amino acids
Saccharate	L- α -Alanine
Mucate	D- α -Alanine
Fatty acids	β -Alanine
Acetate	L-Leucine
Propionate*	L-Isoleucine
Butyrate*	L-Valine*
Valerate	L-Aspartate
Isovalerate	L-Glutamate
Caproate*	DL-Arginine
Heptanoate	γ -Aminobutyrate
Pelargonate	δ -Aminovalerate*
Caprylate	Amino acids and related compounds containing a ring structure
Caprate	L-Histidine
Dicarboxylic acids	L-Proline
Malonate	L-Tyrosine
Succinate	L-Tryptophan
Fumarate	L-Kynurenine
Glutarate	Anthranilate
Hydroxyacids	Amines
L-Malate	Putrescine
DL- β -Hydroxybutyrate	Spermine
DL-Lactate	Histamine*
DL-Glycerate*	Miscellaneous nitrogenous compounds
Miscellaneous organic acids	Betaine
Citrate	Paraffin hydrocarbons
α -Ketoglutarate	(None)
Pyruvate	
Itaconate	
Mesaconate	
Polyalcohols and glycols	
Mannitol	
<i>meso</i> -Inositol	
Glycerol	

* Substrates not used by all strains. Negative strains are given in Table 23.

Substrates used by 90% or more of the strains of biotype D are shown in Table 22; and substrates used by only a fraction of the strains in Table 23, which also lists specifically the negative strains.

Biotype E. Biotype E contains the seven strains which were received as *Pseudomonas aureofaciens*. All produced fluorescent pigment on King B medium, and an

orange pigment (presumably phenazine- α -carboxylic acid) on King A medium. All grow at 4°. All hydrolyse gelatin, strain 41 very weakly; and, with the exception of this strain, all give a positive egg-yolk reaction and hydrolyse Tween 80. All produce levan from sucrose, but none can denitrify.

Table 23. *Pseudomonas fluorescens* biotype D. Substrates used by only a fraction of the strains

Substrate	Total of positive strains	Negative strains
D-Galactose	3	30, 32, 35, 388, 391, 393, 394
D-Fructose	7	388, 393, 394
Propionate	9	388
Butyrate	9	391
Caproate	9	394
Adipate	1	All except 389
D-Malate	6	32, 35, 388, 391
L-(+)-Tartrate	6	32, 35, 388, 391
DL-Glycerate	9	391
Aconitate	6	32, 35, 388, 391
Laevulinate	5	32, 388, 389, 390, 391
2,3-Butylene glycol	4	32, 35, 388, 389, 390, 391
Erythritol	4	30, 388, 389, 390, 393, 394
Ethanol	1	All except 394
n-Propanol	1	All except 394
n-Butanol	4	388, 389, 390, 391, 393, 394
o-Hydroxybenzoate	1	All except 389
m-Hydroxybenzoate	2	All except 32, 388
Phenylacetate	6	30, 31, 393, 394
Testosterone	1	All except 391
L-Serine	6	388, 389, 390, 391
L-Valine	9	388
L-Lysine	6	388, 389, 390, 391
DL-Ornithine	5	32, 388, 391, 393, 394
L-Citrulline	5	32, 388, 391, 393, 394
δ -Aminovalerate	9	388
L-Phenylalanine	4	30, 388, 389, 391, 393, 394
Kynurenate	1	All except 391
Ethanolamine	8	390, 391
Histamine	9	391
Butylamine	8	32, 388
α -Amylamine	7	32, 388, 391
Sarcosine	8	388, 390

Strain 41 is atypical of the entire fluorescent group with respect to the utilization of glucose and related compounds. It is the only fluorescent pseudomonad which cannot grow on glucose or gluconate; one of two strains which cannot use glycerol; and one of five which cannot use 2-ketogluconate. Indeed, the only two sugars able to support its growth are the sterically related compounds galactose and L-arabinose. Strain 41 is none the less a vigorous levan former; hence it can cleave sucrose, even though it cannot use this disaccharide as a carbon source, owing to inability to use glucose, the monosaccharide liberated by the action of levan sucrase.

In the general character of its nutritional spectrum biotype E most closely resembles biotype D (= *Pseudomonas chlororaphis*). The principal respects in which

it differs from biotype D are ability to use L-arabinose, and inability to use ethanol and propanol, coupled of course with failure to denitrify and the production of a slightly different kind of phenazine pigment. (The orange pigment of *P. aureofaciens* is the free acid of phenazine- α -carboxamide, oxychlororaphin; chlororaphin

Table 24. *Pseudomonas fluorescens* biotype E. Substrates used by 85% or more of the strains

Carbohydrates and sugar derivatives	Alcohols
D-Ribose*	(None)
L-Arabinose	Non-nitrogenous aromatic and other
D-Glucose*	cyclic compounds
D-Mannose*	Benzoylformate
D-Galactose	Benzoate*
D-Fructose*	<i>p</i> -Hydroxybenzoate
Sucrose*	Phenylacetate
Trehalose*	Quinate
Gluconate*	Aliphatic amino acids
2-Ketogluconate*	L- α -Alanine
Saccharate	β -Alanine
Mucate	L-Serine
Fatty acids	L-Leucine
Acetate	L-Isoleucine
Propionate	L-Valine
Valerate	L-Aspartate
Isovalerate	L-Glutamate
Caproate	DL-Arginine
Heptanoate	DL-Ornithine
Caprylate	γ -Aminobutyrate
Pelargonate	Amino acids and related compounds con-
Caprate	taining a ring structure
Dicarboxylic acids	L-Histidine
Malonate	L-Proline
Succinate	L-Tyrosine
Fumarate	L-Phenylalanine
Glutarate	L-Tryptophan
Hydroxyacids	L-Kynurenine
L-Malate	Anthranilate
DL- β -Hydroxybutyrate	Amines
DL-Lactate	Ethanolamine
DL-Glycerate	Putrescine
Miscellaneous organic acids	Spermine
Citrate	Histamine*
α -Ketoglutarate	α -Amylamine
Pyruvate	Miscellaneous nitrogenous compounds
Aconitate	Betaine
Itaconate	Sarcosine
Mesaconate	Paraffin hydrocarbons
Polyalcohols and glycols	(None)
Mannitol	
<i>meso</i> -Inositol	
Glycerol*	

* Substrates not used by all strains. Negative strains are given in Table 25.

is a reduced dimer of phenazine- α -carboxamide; Sneath, 1960). The two biotypes D and E are accordingly easily differentiable from one another. Biotype E shares with biotype A the ability to form levan from sucrose, and the inability to denitrify; but they differ from one another in many nutritional characters.

Characters which are absent from biotype E but overwhelmingly positive in biotype A include: utilization of xylose, hydroxymethylglutarate, sorbitol and adonitol; conversely, all strains of biotype E can use benzoylformate and phenylacetate, while none of biotype A can do so. Substrates used by six out of seven or by all strains of biotype D are shown in Table 24; substrates used by only a fraction of the strains in Table 25, which also lists the negative strains.

Table 25. *Pseudomonas fluorescens* biotype E. Substrates utilized by only a fraction of the strains

Substrate	Total of positive strains	Negative strains
D-Ribose	6	41
D-Glucose	6	41
D-Mannose	6	41
D-Fructose	6	41
Sucrose	6	41
Trehalose	6	41
Gluconate	6	41
2-Ketogluconate	6	41
Butyrate	4	37, 39, 40
Isobutyrate	1	All except 41
Laevulinate	5	36, 37
Glycerol	6	41
2,3-Butyleneglycol	5	38, 41
Benzoate	6	37
DL-Citrulline	2	All except 36, 86
DL- α -Aminovalerate	1	All except 36
δ -Aminovalerate	4	37, 38, 41
Kynurenate	1	All except 36
Histamine	6	41
Butylamine	5	36, 37

Biotype F. This biotype, which corresponds to the named species *Pseudomonas lemonnieri*, is unfortunately represented by only two strains, so that the weight which should be placed on the characters that appear to differentiate it from other biotypes of *P. fluorescens* is uncertain. Strain 83 is an authentic strain studied by Hugo & Turner (1957), whereas strain 143 was isolated in Berkeley. The recognition of *P. lemonnieri* by means of its most distinctive character, the production of a blue insoluble phenazine pigment (Starr *et al.* 1960), is not easy in our experience. In the first place, King A medium, which is otherwise a very reliable medium for demonstrating the production of phenazine pigments, fails completely with strains of this biotype; and it is necessary to use another special medium (see Methods). Even on this special medium, the results were erratic; strain 83 produced only blue sectors, indicating that the population was genetically heterogeneous with respect to the character of specific phenazine production; and strain 143 gave negative results, even though it had been tentatively identified as *P. lemonnieri* by virtue of its characteristic colour upon primary isolation several years ago. However, in other respects these two independent isolates are very similar, and share a number of properties which seem to set them apart from the biotypes so far described. Both produce fluorescent pigment on King B medium. Both liquefy gelatin and give an egg-yolk reaction; one strain (83) hydrolyses Tween 80. Both grow at 4°. Both are

levan formers and denitrifiers. Hugo & Turner (1957) reported monotrichous flagellation for the strains of *P. lemonnieri* studied by them. However, we found both the strains studied by us to be multitrichous; this has been confirmed by Dr H. Lautrop (personal communication).

Table 26. *Pseudomonas fluorescens* biotype F. Substrates utilized by the two strains

Carbohydrates and sugar derivatives	Polyalcohols and glycols (<i>cont.</i>)
D-Ribose	meso-Inositol
L-Arabinose	Glycerol
D-Glucose	2,3-Butyleneglycol
D-Mannose	Alcohols
D-Galactose	n-Butanol
D-Fructose	Non-nitrogenous aromatic and other
Sucrose	cyclic compounds
Trehalose	Benzoate
Gluconate	o-Hydroxybenzoate
Saccharate	Quinate
Mucate	Aliphatic amino acids
Fatty acids	L- α -Alanine
Acetate	D- α -Alanine
Butyrate	β -Alanine
Caproate	L-Serine
Heptanoate	L-Leucine
Caprylate	L-Isoleucine
Pelargonate	L-Valine
Caprate	L-Aspartate
Dicarboxylic acids	L-Glutamate
Malonate	L-Lysine
Succinate	DL-Arginine
Fumarate	γ -Aminobutyrate
Glutarate	δ -Aminovalerate
Hydroxyacids	Amino acids and related compounds con-
D-Malate	taining a ring structure
L-Malate	L-Histidine
L-(+)-Tartrate	L-Proline
DL- β -Hydroxybutyrate	L-Tyrosine
DL-Lactate	L-Phenylalanine
DL-Glycerate	Amines
Miscellaneous organic acids	Ethanolamine
Citrate	Putrescine
α -Ketoglutarate	Spermine
Pyruvate	Miscellaneous nitrogenous compounds
Aconitate	Betaine
Citraconate	Sarcosine
Polyalcohols and glycols	Paraffin hydrocarbons
Mannitol	(None)
Sorbitol	

On the properties of denitrification and levan formation, the strains resemble biotypes B and D, from which however they are distinguishable by failure to grow on propionate, valerate, isovalerate, ethanol and *n*-propanol. Furthermore, biotype F can be distinguished from biotype B by failure to grow on hydroxymethylglutarate, propyleneglycol and trigonelline; and from biotype D by failure to grow on benzoylformate, phenylacetate and butylamine, and by growth on L-arabinose and sorbitol.

The compounds which can be utilized as sources of carbon and energy are shown in Table 26. The strains differ in only three nutritional respects: utilization of D-xylose and 2-ketogluconate (positive for strain 83) and of ornithine (positive for strain 143).

Table 27. *Pseudomonas fluorescens* biotype G. Substrates utilized by 90 % or more of the strains

Carbohydrates and sugar derivatives	Alcohols
D-Ribose*	(None)
D-Glucose	Non-nitrogenous aromatic and other
D-Mannose*	cyclic compounds
D-Fructose	p-Hydroxybenzoate
Trehalose	Quinate
Gluconate	Aliphatic amino acids
2-Ketogluconate	L- α -Alanine
Fatty acids	β -Alanine
Acetate	L-Serine*
Caproate	L-Leucine*
Heptanoate	L-Isoleucine
Caprylate	L-Valine
Pelargonate	L-Aspartate
Caprate	L-Glutamate
Dicarboxylic acids	DL-Arginine
Malonate	DL-Ornithine*
Succinate	γ -Aminobutyrate
Fumarate	Amino acids and related compounds con-
Glutarate*	taining a ring structure
Hydroxyacids	L-Histidine*
L-Malate	L-Proline
DL- β -Hydroxybutyrate	L-Tyrosine
DL-Lactate	Amines
DL-Glycerate	Putrescine*
Miscellaneous organic acids	Spermine*
Citrate	Miscellaneous nitrogenous compounds
α -Ketoglutarate*	Betaine*
Pyruvate	Sarcosine
Aconitate	Trigonelline*
Polyalcohols and glycols	Paraffin hydrocarbons
Mannitol*	(None)
Glycerol	

* Substrates not used by all strains. Negative strains are given in Table 28.

Biotype G. This is much the least satisfactory biotype of *Pseudomonas fluorescens* and consists essentially of strains with the general character of the species which remained unaccounted for after the recognition of biotypes A–F. All except strains 164 and 172 produce fluorescent pigment. All are gelatin liquefiers; most give an egg-yolk reaction (negative: strains 169, 172, 206, 267) and hydrolyse Tween 80 (negative: strains 149, 206, 267, 272). One strain (267) is unable to grow at 4°, an otherwise invariable property of *P. fluorescens*. None can denitrify. None can form levan from sucrose, though strains 99, 164, 172 and 269 can grow at the expense of this sugar, thus apparently constituting exceptions to the rule that *P. fluorescens* attacks sucrose by means of a levan sucrase. Substrates used by 90 % or more of the strains are shown in Table 27; and substrates used by only a fraction of the

strains, in Table 28. The characteristic nutritional irregularity of biotype G, shown in the data of Table 15, has already been mentioned. However, this irregularity is at least in part a reflexion of the fact that biotype G assembles representatives of what might well have emerged (had more strains been examined) as additional biotypes of *P. fluorescens*. Some indication of this can be obtained from the perusal of Table 28, which shows certain repeated clusterings of strains. This is particularly true for strains 1, 99 and 124, which differ from the rest of biotype G in many nutritional characters, both positive and negative.

Table 28. *Pseudomonas fluorescens* biotype G. Substrates utilized by only a fraction of the strains

Substrate	Total of positive strains	Negative strains
D-Ribose	16	124
D-Xylose	13	1, 99, 124, 271
L-Arabinose	14	1, 99, 124
L-Rhamnose	2	All except 267, 272
D-Mannose	15	124, 164
D-Galactose	14	1, 99, 124
Sucrose	4	All except 99, 164, 172, 269
Saccharate	14	1, 99, 124
Mucate	13	1, 99, 124, 272
Propionate	14	267, 269, 271
Butyrate	3	All except 1, 99, 124
Isobutyrate	1	All except 1
Valerate	5	All except 1, 99, 124, 206, 269
Isovalerate	8	33, 34, 149, 164, 166, 172, 267, 271, 272
Glutarate	16	164
Adipate	3	All except 1, 99, 124
Suberate	1	All except 1
Azelate	2	All except 1, 99
Sebacate	4	All except 1, 99, 124, 272
D-Malate	4	All except 195, 267, 269, 271
L-Serine	15	99, 124
L-Leucine	15	267, 272
L-Lysine	13	33, 34, 267, 271
DL-Ornithine	15	169, 271
DL-Citrulline	10	169, 171, 172, 206, 269, 271, 272
DL- α -Aminobutyrate	1	All except 166
DL- α -Aminovalerate	1	All except 1
δ -Aminovalerate	14	33, 34, 267
L-Histidine	16	272
L-Phenylalanine	10	1, 124, 166, 171, 195, 271, 272
L-Tryptophan	11	164, 169, 171, 172, 206, 269
L-Kynurenine	11	164, 169, 171, 172, 206, 269
Kynurenate	2	All except 99, 124
Anthranilate	11	164, 169, 171, 172, 206, 269
Ethanolamine	13	164, 169, 172, 269
Benzylamine	3	All except 1, 124, 171
Putrescine	16	169
Spermine	16	169
Histamine	3	All except 1, 99, 124
Tryptamine	1	All except 272
Butylamine	3	All except 1, 99, 124
α -Amylamine	4	All except 1, 99, 124, 149
Betaine	15	124, 267

Table 28 (cont.)

Substrate	Total of positive strains	Negative strains
Nicotinate	9	1, 99, 164, 169, 171, 172, 206, 269
Trigonelline	15	1, 99
D-(-)-Tartrate	1	All except 169
L-(+)-Tartrate	1	All except 267
meso-Tartrate	11	1, 99, 124, 169, 206, 272
Hydroxymethylglutarate	5	All except 33, 34, 149, 166, 195
α -Ketoglutarate	16	267
Laevulinate	1	All except 99
Citraconate	9	1, 33, 34, 99, 124, 169, 172, 206
Itaconate	9	99, 164, 169, 171, 172, 206, 269, 272
Mesaconate	7	All except 1, 33, 34, 149, 166, 195, 271
Erythritol	8	1, 99, 164, 169, 171, 172, 206, 269, 271
Mannitol	16	124
Sorbitol	8	1, 99, 124, 164, 169, 171, 172, 206, 269
meso-Inositol	11	164, 169, 171, 172, 206, 269
Adonitol	8	1, 99, 124, 164, 169, 171, 172, 206, 269
Propyleneglycol	8	All except 1, 99, 124, 149, 166, 195, 267, 272
2,3-Butyleneglycol	8	All except 1, 33, 34, 99, 149, 166, 195, 272
Ethanol	8	All except 1, 99, 124, 149, 166, 195, 267, 272
n-Propanol	10	33, 34, 171, 172, 206, 269, 271
n-Butanol	12	33, 34, 206, 267, 271
Isobutanol	11	33, 34, 172, 206, 269, 271
L-Mandelate	2	All except 99, 124
Benzoylformate	3	All except 1, 99, 124
Benzoate	5	All except 1, 99, 124, 164, 171
Phenylacetate	2	All except 99, 124
Testosterone	1	All except 124

Pseudomonas putida

Of the 175 strains belonging to the fluorescent group that we studied, 52 are unable to hydrolyse gelatin or to give an egg-yolk reaction, do not denitrify, do not grow at 41°, and do not produce phenazine pigment. By virtue of this constellation of characters, they are separable both from *Pseudomonas aeruginosa* and from the seven biotypes of *P. fluorescens*. The failure to hydrolyse gelatin is, of course, the one character which has classically defined *P. putida* (= *P. fluorescens non-liquefaciens* Flüggé), and we were therefore initially inclined to regard all these strains as being representative of *P. putida*. In nutritional respects, however, they are not an internally uniform group. A total of 32 strains constitute a fairly homogeneous major subgroup, and we have decided arbitrarily to consider them the central biotype of *P. putida*; we shall term this subgroup biotype A. Another nine strains, internally homogeneous, resemble biotype A in most respects, but are more versatile nutritionally, and will be described as *P. putida* biotype B. The remaining 11 strains present unsolved taxonomic problems, and we shall simply describe them as 'unclassified strains of the fluorescent group'.

Biotype A. All 32 strains produce fluorescent pigment on King B medium. Only two (42, 144) can hydrolyse Tween 80. The ability to grow at 4° is relatively rare,

being possessed only by strains 6, 26, 49, 56, 90, 100, 111, 115, 119, 122, 145 (eleven in all). Substrates used by 90 % or more of the strains are shown in Table 29; substrates used by a fraction of the strains are listed in Table 30, with negative strains included.

Table 29. *Pseudomonas putida* biotype A. Substrates used by 90% or more of the strains

Carbohydrates and sugar derivatives	Non-nitrogenous aromatic and other cyclic compounds
D-Glucose	Benzoate
D-Fructose*	<i>p</i> -Hydroxybenzoate
Gluconate	Phenylacetate*
2-Ketogluconate	Quinate*
Saccharate	Aliphatic amino acids
Mucate	L- α -Alanine*
Fatty acids	D- α -Alanine*
Acetate	β -Alanine
Propionate	L-Leucine
Butyrate*	L-Isoleucine
Valerate	L-Valine*
Isovalerate	L-Aspartate
Caproate	L-Glutamate
Heptanoate*	L-Lysine
Caprylate	DL-Arginine
Pelargonate	DL-Ornithine
Caprate	DL-Citrulline
Dicarboxylic acids	γ -Aminobutyrate
Succinate	δ -Aminovalerate
Fumarate	Amino acids and related compounds containing a ring structure
Glutarate*	L-Histidine
Hydroxyacids	L-Proline
L-Malate	L-Tyrosine
DL- β -Hydroxybutyrate	L-Phenylalanine*
DL-Lactate	Amines
Miscellaneous organic acids	Benzylamine
Citrate	Putrescine
α -Ketoglutarate*	Spermine*
Pyruvate	Histamine
Aconitate	Butylamine*
Polyalcohols and glycols	α -Amylamine*
Glycerol*	Miscellaneous nitrogenous compounds
Alcohols	Betaine
Ethanol*	Sarcosine
<i>n</i> -Propanol*	Trigonelline
<i>n</i> -Butanol*	Paraffin hydrocarbons
Isobutanol	(None)

* Substrates not used by all strains. Negative strains are listed in Table 30.

Biotype B. The nine strains of biotype B, with the exception of 98, produce fluorescent pigment. One strain (153) hydrolyses Tween 80. This group of strains can be most readily distinguished from biotype A by their ability to grow on L-tryptophan and L-kynurenine (used by no strain of biotype A) and on anthranilate (used by only two strains of biotype A, 51 and 276). In some respects, biotype B is more fluorescens-like than biotype A: all strains can grow at 4°, and the utilization of sugars is more extensive than in biotype A. The most useful sugar for the dif-

ferentiation between biotypes A and B is galactose, used by no strain of biotype A, and by seven out of nine strains of biotype B. Biotype B has a significantly wider general nutritional spectrum than biotype A, as shown by the much higher frequencies with which such substrates as testosterone, phenol, salicylate, naphthalene and tryptamine are attacked. Substrates used by 90 % or more of the strains are shown in Table 31. Table 32 gives the list of substrates attacked by a fraction of the strains, with negative strains indicated. The only two substrates utilized by a large fraction of the strains of biotype A and by few or none of the strains of biotype B are α -aminovalerate (used only by strain 110 of biotype B) and nicotinate (Table 33). The nutritional characters of tryptophan, kynurenine, galactose and nicotinate utilization appear to be the only absolute criteria for differentiating the two biotypes.

Unclassified strains

Of the 11 strains in this category, all save one (109) produce fluorescent pigment. Eight of them (10, 17, 47, 59, 91, 92, 127, 322) could be regarded as strains of *Pseudomonas putida*, which they resemble in failing to denitrify, to hydrolyse gelatin or to give an egg-yolk reaction, and in rarely hydrolysing Tween 80 (only strain 322 positive). A possible exception is strain 10, which was described by den Dooren de Jong (1926) as liquefying gelatin. In our hands, however, strain 10 has given a faint reaction under the patch by the test of Skerman (1959) after several days of incubation, probably attributable to lysis and release of intracellular proteases. What distinguishes these eight strains from biotypes A and B of *P. putida* is essentially their much more limited nutritional versatility, as shown by the data in Tables 34 and 35. The many typical nutritional characters of *P. putida* which they lack would have conferred an undesirable heterogeneity on this species had they been included.

The remaining three unclassified strains (95, 101, 109) present a different problem. They are the only strains which appear to be intermediate between *Pseudomonas fluorescens* and *P. putida* with respect to the key nutritional characters that otherwise serve well to distinguish these species. They are putida-like in the sense that none can denitrify, hydrolyse gelatin or give an egg-yolk reaction; and only one (strain 101) can hydrolyse Tween 80. All grow at 4°. Two (95, 101) can use trehalose, a character universally positive in *P. fluorescens* and universally negative in *P. putida*; and they share with several biotypes of *P. fluorescens* the inability to grow on butyrate, isobutyrate, ethanol or *n*-propanol. Strain 109 cannot grow on trehalose, but is fluorescens-like in its ability to use D-xylose, L-arabinose, D-galactose, hydroxymethylglutarate, sorbitol or adonitol; and in its failure to use butyrate. The nutritional characters of these three exceptional strains are shown in Tables 36 and 37.

Neotype strains in the fluorescent group

According to the catalogue of the American Type Culture Collection (1964) no less than three different neotype strains have been proposed for *Pseudomonas aeruginosa*: the cultures carrying ATCC numbers 10145, 14209 and 14216. We were unfortunately not aware of this fact until recently, and none of these strains was included in our study. However, in view of the phenotypic uniformity and

Table 30. *Pseudomonas putida* biotype A. Substrates used by a fraction of the strains

Substrate	Total of positive strains	Negative strains
D-Ribose	17	8, 26, 49, 51, 56, 77, 87, 100, 111, 115, 118, 119, 130, 144, 266
D-Xylose	5	All except 5, 8, 87, 100, 119
L-Arabinose	6	All except 6, 100, 118, 119, 128, 130
D-Mannose	6	All except 5, 87, 119, 145, 266, 276
D-Fructose	30	44, 100
Sucrose	3	All except 145, 154, 160
Butyrate	31	100
Isobutyrate	28	8, 100, 111, 115
Heptanoate	31	118
Malonate	21	5, 42, 43, 44, 49, 51, 56, 87
Glutarate	31	89
D-Malate	18	5, 8, 42, 43, 44, 56, 87, 100, 118, 119, 128, 130, 154, 160
D-(-)-Tartrate	5	All except 6, 82, 100, 119, 128
D-(+)-Tartrate	23	5, 6, 8, 118, 122, 130, 145, 154, 160
meso-Tartrate	16	5, 6, 8, 42, 43, 44, 49, 51, 56, 87, 100, 119, 130, 145, 154, 160
Glycollate	3	All except 51, 82, 145
α -Ketoglutarate	31	89
Laevulinate	8	All except 7, 49, 81, 118, 130, 266, 276
Citraconate	2	All except 118, 130
Itaconate	6	All except 8, 100, 118, 119, 130, 276
Mesaconate	5	All except 8, 100, 118, 119, 130
Erythritol	1	All except 118
Mannitol	3	All except 8, 100, 144
Sorbitol	1	All except 118
Adonitol	1	All except 8
Glycerol	31	122
Ethyleneglycol	1	All except 145
Propyleneglycol	26	5, 7, 8, 115, 118, 130
2,3-Butyleneglycol	23	6, 7, 8, 81, 89, 100, 118, 119, 130
Ethanol	31	8
n-Propanol	31	8
n-Butanol	30	8, 100
D-Mandelate	7	All except 6, 7, 49, 81, 89, 90, 111
L-Mandelate	2	All except 89, 90
Benzoylformate	15	All except 6, 7, 26, 42, 43, 44, 49, 81, 89, 90, 111, 115, 122, 145, 276
o-Hydroxybenzoate	3	All except 51, 111, 115
m-Hydroxybenzoate	3	All except 43, 44, 56
p-Hydroxybenzoate	30	76, 77
Phenylacetate	29	8, 100 and 111
Phenylethanediol	5	All except 81, 89, 90, 111, 160
Naphthalene	1	All except 111
Phenol	5	All except 51, 76, 77, 89, 144
Quinate	30	76, 77
Testosterone	1	All except 130
Glycine	26	8, 81, 100, 118, 119, 130
L- α -Alanine	31	160
D- α -Alanine	31	8
L-Serine	20	5, 6, 7, 51, 76, 77, 82, 89, 119, 122, 128, 144
L-Valine	31	5
DL- α -Aminobutyrate	2	All except 76, 266
DL- α -Aminovalerate	22	8, 26, 56, 81, 100, 115, 118, 119, 130, 276

Table 30 (cont.)

Substrate	Total of positive strains	Negative strains
L-Phenylalanine	31	145
Kynurenate	4	All except 8, 26, 100, 115
Anthranilate	2	All except 51, 276
Ethanolamine	17	6, 7, 26, 49, 51, 76, 77, 81, 82, 87, 89, 90, 118, 130, 266
Spermine	31	81
Tryptamine	7	All except 6, 7, 89, 115, 118, 122, 130
Butylamine	29	8, 49, 100
α -Amylamine	29	8, 49, 100
Creatine	28	82, 128, 144, 266
Hippurate	23	81, 89, 100, 118, 122, 128, 130, 144, 266
Pantothenate	1	All except 145
Acetamide	10	All except 43, 44, 51, 56, 76, 77, 87, 145, 154, 160
Nicotinate	21	5, 6, 7, 49, 82, 115, 122, 128, 144, 266

distinctiveness of *P. aeruginosa*, it seems probable that almost any strain would be a satisfactory neotype.

A neotype strain has been proposed for *Pseudomonas fluorescens* (NCTC 10038 = ATCC 13525) by Rhodes (1959). Since she did not accept as valid the distinction between *P. fluorescens* and *P. putida*, it was chosen by her to be representative of this whole assemblage. We have examined the strain in question (our designation: 192), which proves to be a typical representative of *P. fluorescens* biotype A. We therefore concur (on different grounds) with the proposal of Rhodes. Acceptance of this neotype will have the consequence of fixing the designation *P. fluorescens* on our biotype A, if subsequent work shows that some or all of the biotypes of *P. fluorescens* merit species rank.

Apparently no neotype strain has been proposed for *Pseudomonas putida*. Accordingly, we propose our strain 90 (former designation A.3.12 = ATCC 12633), which is a typical member of *P. putida* biotype A. This particular strain has been used over the past 20 years in many different biochemical and physiological studies, and there is consequently a large body of information about its metabolism and physiology.

Nomenclatural problems in the fluorescent group

The account of the properties of *Pseudomonas fluorescens* has presented the evidence which leads us to classify the three phenazine producers *P. chlororaphis*, *P. aureofaciens* and *P. lemonnieri* as biotypes of *P. fluorescens*. These three names could perhaps be preserved as varietal designations for our biotypes D, E and F.

A much more difficult nomenclatural problem is presented by the many named gelatin-liquefying fluorescent species which do not produce phenazine pigments. Most of these are plant pathogens; and the question of the possible synonymy of these species with *Pseudomonas fluorescens* must await a detailed comparative study of the phytopathogenic fluorescent pseudomonads by the methods which we have used to characterize *P. fluorescens*. However, Haynes (1957) also recognized in the latest (7th) edition of *Bergey's Manual of Determinative Bacteriology* a total of nine

Table 31. *Pseudomonas putida* biotype B. Substrates used by 90% or more of the strains

Carbohydrates and sugar derivatives	Non-nitrogenous aromatic and other
L-Arabinose*	cyclic compounds
D-Glucose	Benzoate
D-Fructose	<i>p</i> -Hydroxybenzoate
Gluconate	Phenylacetate
2-Ketogluconate	Quinate
Saccharate*	Testosterone*
Mucate	Aliphatic amino acids
Fatty acids	Glycine
Acetate	L- α -Alanine
Propionate	D- α -Alanine
Butyrate	β -Alanine
Valerate	L-Serine*
Isovalerate	L-Leucine
Caproate	L-Isoleucine
Heptanoate	L-Valine
Caprylate	L-Aspartate
Pelargonate	L-Glutamate
Caprate	L-Lysine*
Dicarboxylic acids	DL-Arginine
Malonate	DL-Ornithine
Succinate	γ -Aminobutyrate
Fumarate	δ -Aminovalerate
Glutarate	Amino acids and related compounds con-
Hydroxyacids	taining a ring structure
L-Malate	L-Histidine*
DL- β -Hydroxybutyrate	L-Proline
DL-Lactate	L-Tyrosine
DL-Glycerate	Phenylalanine*
Miscellaneous organic acids	L-Tryptophan
Citrate	L-Kynurenine
α -Ketoglutarate	Anthranilate
Pyruvate	Amines
Aconitate	Benzylamine
Polyalcohols and glycols	Putrescine
Glycerol	Spermine
Alcohols	Histamine*
Ethanol*	Tryptamine
<i>n</i> -Propanol*	Butylamine
<i>n</i> -Butanol	α -Amylamine
Isobutanol*	Miscellaneous nitrogenous compounds
	Betaine
	Sarcosine
	Trigonelline
	Paraffin hydrocarbons
	(None)

* Substrates not used by all strains. Negative strains are listed in Table 32.

non-phytopathogenic, gelatin-liquefying species in addition to *P. fluorescens*. They are: *P. reptilivora*, *P. caviae*, *P. boreopolis*, *P. effusa*, *P. fairmontensis*, *P. myxogenes*, *P. schuylkilliensis*, *P. geniculata*, *P. septica*. We have examined one strain with the label *P. schuylkilliensis* (strain 267) and three with the label *P. geniculata* (strains 269, 271, 272). All were assigned to *P. fluorescens* biotype G. Since there is nothing in the published descriptions of any of these nine species which would permit a

distinction from *P. fluorescens*, we believe that they should all be reduced to synonymy with it, unless good evidence to the contrary can be obtained through a new examination of the type strains, if any of these is still in existence.

An entirely comparable nomenclatural problem exists for the fluorescent species which do not liquefy gelatin. There is a number of named plant pathogens in this

Table 32. *Pseudomonas putida* biotype B. Substrates used by a fraction of the strains

Substrate	Total of positive strains	Negative strains
D-Ribose	4	53, 96, 98, 107, 110
D-Xylose	6	96, 153, 157
L-Arabinose	8	98
D-Mannose	7	110, 153
D-Galactose	7	98, 153
Sucrose	1	All except 158
Saccharate	8	98
Isobutyrate	7	98, 107
Adipate	1	All except 107
Sebacate	1	All except 107
D-Malate	2	All except 96, 107
D-(-)-Tartrate	2	All except 98, 158
L-(+)-Tartrate	2	All except 96, 107
Laevulinate	3	53, 96, 98, 110, 153, 157
Citraconate	5	96, 98, 110, 157
Mannitol	5	98, 110, 157, 158
Sorbitol	2	All except 53, 107
2,3-Butyleneglycol	3	53, 107, 110, 153, 157, 158
Ethanol	8	107
n-Propanol	8	107
Isobutanol	8	107
D-Mandelate	1	All except 53
L-Mandelate	1	All except 53
Benzoylformate	1	All except 53
o-Hydroxybenzoate	6	96, 98, 158
Phenylethanediol	1	All except 53
Naphthalene	2	All except 107, 110
Phenol	6	53, 158, 167
Testosterone	8	110
L-Serine	8	98
L-Lysine	8	107
DL-Citrulline	5	96, 153, 157, 158
DL- α -Aminovalerate	1	All except 110
L-Histidine	8	53
L-Phenylalanine	8	107
D-Tryptophan	1	All except 110
Kynurenate	6	96, 98 and 110
Ethanolamine	3	All except 98, 158, 167
Histamine	8	98
Creatine	5	98, 107, 153, 157
Hippurate	7	98, 110
Acetamide	1	All except 110

category, the status of which *vis-à-vis* *Pseudomonas putida* cannot be decided at present. Haynes (1957) also listed several non-phytopathogenic species, in addition to *P. putida*, namely: *P. striata*, *P. ovalis*, *P. incognita*, *P. rugosa*, *P. convexa*,

Table 33. *Pseudomonas putida* biotypes A and B. Selected characters diagnostically important for the differentiation of biotypes A and B

Character	Biotype A		Biotype B	
	No. of positive strains	% of positive strains	No. of positive strains	% of positive strains
Growth at 4°	11	34	9	100
Utilization of				
D-Xylose	5	16	6	67
L-Arabinose	6	19	8	89
D-Mannose	6	19	7	78
D-Galactose	0	0	7	78
Mannitol	3	9	5	56
o-Hydroxybenzoate	3	9	6	67
Phenol	5	16	6	67
Testosterone	1	3	8	89
α -Aminovalerate	22	69	1	11
L-Tryptophan	0	0	9	100
L-Kynurenine	0	0	9	100
Anthranilate	2	6	9	100
Tryptamine	7	22	9	100
Nicotinate	21	66	0	0

Table 34. *Unclassified fluorescent pseudomonads. Substrates utilized by seven or all eight strains (10, 17, 47, 59, 91, 92, 127, 322)*

Carbohydrates and sugar derivatives	Non-nitrogenous aromatic and other
D-Glucose	cyclic compounds
Gluconate	Quinate*
Fatty acids	Aliphatic amino acids
Acetate	L- α -Alanine
Propionate	D-Alanine*
Caproate*	β -Alanine*
Caprylate*	L-Serine*
Pelargonate	L-Leucine*
Caprate*	L-Isoleucine*
Dicarboxylic acids	L-Valine*
Succinate	L-Aspartate
Fumarate	L-Glutamate
Glutarate*	DL-Arginine
Hydroxyacids	γ -Aminobutyrate*
L-Malate	δ -Aminovalerate*
DL-Lactate	Amino acids and related compounds con-
DL-Glycerate	taining a ring structure
Miscellaneous organic acids	L-Histidine*
Citrate*	L-Proline*
α -Ketoglutarate	Amines
Pyruvate*	Putrescine
Polyalcohols and glycols	Spermine*
Glycerol	Miscellaneous nitrogenous compounds
Alcohols	Betaine
(None)	Sarcosine
	Paraffin hydrocarbons
	(None)

* Substrates utilized by seven out of eight strains. Negative strains are indicated in Table 35.

Table 35. *Unclassified fluorescent pseudomonads. Substrates utilized by a fraction of the eight strains*

Substrate	Total of positive strains	Negative strains
D-Ribose	3	10, 47, 59, 127, 322
D-Xylose	1	All except 92
L-Arabinose	2	All except 10, 92
D-Mannose	1	All except 92
D-Galactose	2	All except 10, 92
D-Fructose	6	10, 91
Maltose	1	All except 322
2-Ketogluconate	4	17, 47, 127, 322
Saccharate	5	17, 47, 91
Mucate	5	17, 47, 91
Butyrate	5	10, 59, 127
Isobutyrate	4	10, 59, 91, 322
Valerate	5	10, 59, 92
Isovalerate	3	10, 59, 92, 127, 322
Caproate	7	10
Heptanoate	6	10, 92
Caprylate	7	10
Caprate	7	322
Malonate	4	17, 91, 92, 322
Glutarate	7	127
D-Malate	4	10, 59, 92, 127
D- α -Alanine	7	10
β -Alanine	7	10
L-Serine	7	10
L-Leucine	7	322
L-Isoleucine	7	322
L-Valine	7	10
L-Lysine	5	10, 127, 322
DL-Ornithine	6	10, 59
DL-Citrulline	6	10, 127
γ -Aminobutyrate	7	322
δ -Aminovalerate	7	322
L-Histidine	7	322
L-Proline	7	92
L-Tyrosine	6	10, 92
L-Phenylalanine	5	10, 127, 322
Ethanolamine	5	10, 59, 92
Benzylamine	1	All except 17
Spermine	7	92
Histamine	2	All except 59, 92
Butylamine	4	10, 59, 92, 127
α -Amylamine	3	10, 17, 92, 127, 322
Creatine	1	All except 92
Hippurate	1	All except 92
Nicotinate	1	All except 92
Trigonelline	3	10, 17, 47, 91, 322
D-(-)-Tartrate	3	17, 47, 91, 127, 322
L-(+)-Tartrate	1	All except 127
meso-Tartrate	4	10, 47, 59, 322
DL- β -Hydroxybutyrate	6	92, 127
Hydroxymethylglutarate	1	All except 127
Citrate	7	322
Pyruvate	7	10
Aconitate	3	10, 17, 47, 91, 322
Laevulinate	1	All except 322

Table 35 (*cont.*)

Substrate	Total of positive strains	Negative strains
Citraconate	1	All except 322
Itaconate	2	All except 92, 322
Mesaconate	2	All except 92, 322
Mannitol	1	All except 92
<i>meso</i> -Inositol	1	All except 92
Propyleneglycol	6	10 and 322
2,3-Butyleneglycol	5	10, 92, 322
Ethanol	5	47, 92, 127
<i>n</i> -Propanol	5	47, 92, 127
<i>n</i> -Butanol	3	10, 47, 59, 92, 127, 322
Isobutanol	6	10, 322
Benzoylformate	1	All except 59
Benzoate	2	All except 92, 322
<i>p</i> -Hydroxybenzoate	6	92, 322
Phenylacetate	3	10, 59, 92, 127, 322
Quinate	7	322
Glycine	2	All except 127, 322

Table 36. *Unclassified fluorescent pseudomonads. Substrates utilized by strains 95, 101 and 109*

Carbohydrates and sugar derivatives	Alcohols
D-Ribose	(None)
D-Glucose	Non-nitrogenous aromatic and other
D-Mannose	cyclic compounds
D-Fructose	(None)
Gluconate	Aliphatic amino acids
2-Ketogluconate	L- α -Alanine
Mucate	β -Alanine
Fatty acids	L-Leucine
Acetate	L-Isoleucine
Propionate	L-Aspartate
Heptanoate	L-Glutamate
Caprylate	DL-Arginine
Pelargonate	DL-Citrulline
Caprate	γ -Aminobutyrate
Dicarboxylic acids	δ -Aminovalerate
Malonate	Amino acids and related compounds con-
Succinate	taining a ring structure
Fumarate	L-Histidine
Glutarate	L-Proline
Hydroxyacids	Amines
L-Malate	Ethanolamine
DL- β -Hydroxybutyrate	Putrescine
DL-Lactate	Spermine
DL-Glycerate	Miscellaneous nitrogenous compounds
Miscellaneous organic acids	Betaine
Citrate	Sarcosine
α -Ketoglutarate	Trigonelline
Pyruvate	Paraffin hydrocarbons
Aconitate	(None)
Polyalcohols and glycols	
Mannitol	
Glycerol	

Table 37. *Unclassified fluorescent pseudomonads. Substrates used by one or two of the strains 95, 101 and 109*

Substrate	Strain			Substrate	Strain		
	95	101	109		95	101	109
D-Xylose	—	—	+	n-Propanol	—	—	+
L-Arabinose	—	—	+	n-Butanol	—	±	—
D-Galactose	—	—	+	Benzoate	+	—	—
Sucrose	+	+	—	o-Hydroxybenzoate	—	—	+
Trehalose	+	+	—	m-Hydroxybenzoate	+	—	—
Saccharate	—	+	+	p-Hydroxybenzoate	+	+	—
Caproate	+	—	+	Naphthalene	—	—	+
D-Malate	—	—	+	Quinate	+	+	—
D-(—)-Tartrate	—	—	+	Testosterone	+	+	—
L-(+)-Tartrate	—	+	—	Glycine	±	+	—
meso-Tartrate	—	—	+	D- α -Alanine	+	+	—
Glycollate	—	—	+	L-Serine	—	+	—
Hydroxymethylglutarate	—	—	+	L-Valine	+	+	—
Laevulinate	—	±	—	L-Lysine	+	±	—
Citraconate	—	—	+	DL-Ornithine	+	+	—
Itaconate	±	—	+	L-Tyrosine	+	—	—
Mesaconate	—	—	+	L-Phenylalanine	+	±	—
Sorbitol	—	—	+	L-Tryptophan	+	—	—
meso-Inositol	+	±	—	L-Kynurenine	+	+	—
Adonitol	—	—	+	Anthranilate	±	±	—
Ethyleneglycol	—	—	+	Histamine	—	+	+
Propyleneglycol	—	—	+	Butylamine	+	—	—
2,3-Butyleneglycol	—	—	±	α -Amylamine	+	—	—
Ethanol	—	—	+	Nicotinate	—	+	+

P. eisenbergii. There is nothing in the published descriptions of these species which would permit their differentiation from *P. putida*, so again we recommend reduction to synonymy, unless type strains still exist which can be shown to differ significantly from *P. putida* as we have described it.

Many workers (e.g. Lautrop & Jessen, 1964) apply the name *Pseudomonas ovalis* to strains which we consider to be typical of *P. putida*. This name, first used by Chester (1901), is derived from the characteristic shape of the organisms, which are small, plump and oval. In our experience, this shape occurs, either exclusively or predominantly, in many strains of *P. putida*. However, other typical strains of *P. putida* are longer and are indistinguishable on a morphological basis from *P. fluorescens*. Jessen (1965) has made similar comments about the variability of shape in his biotype 11, group II, which encompasses strains that we would assign to *P. putida*. It therefore seems evident that the shape of the organisms cannot be considered a reliable character for the definition of a species among the fluorescent pseudomonads which do not liquefy gelatin; and there is nothing else in Chester's description of *P. ovalis* which justifies a separation from *P. putida*. Since *P. putida* is clearly the older specific name, first applied in 1889 by Trevisan, and derived from Flügge's trinomial *Bacillus fluorescens putidus*, we cannot see any justification for the continued use of the name *P. ovalis*.

THE ACIDOVORANS GROUP

Twenty-six strains were assigned to this group, within which we recognize two species: *Pseudomonas acidovorans* den Dooren de Jong (15 strains), and *P. testosteroni* Marcus & Talalay (nine strains). Two strains could not be given a definite specific assignment, but appear on their general characters to be members of this group.

Origins of the strains(a) Strains assigned to *Pseudomonas acidovorans* den Dooren de Jong (1926)

14. Strain 7 of den Dooren de Jong (1926). Isolated from soil by aerobic enrichment with acetamide. Received from Laboratorium voor Microbiologie, Delft. ATCC 15668.
24. *P. desmolytica* strain 4B of Dr M. Yano (no published description). Isolated from soil by aerobic enrichment with *p*-hydroxybenzoate. Received from Institute of Applied Microbiology, Tokyo. ATCC 17406.
29. *P. indoloxidans* Gray (1928). Isolated from soil through aerobic enrichment with indole. Authentic strain; ATCC 9355.
61. Strain 60.79 of Dr P. Thibault, Institut Pasteur. Strain 452-3 of Dr R. Hugh. Isolated from a pharyngeal swab. ATCC 17438.
62. Strain 60.78 of Dr P. Thibault. Strain 553-3 of Dr R. Hugh. Isolated from pharyngeal swab. ATCC 17439.
80. Strain A of Dr U. Bachrach (1957). Received from Professor S. Dagley, Leeds University. ATCC 17455.
102. Strain B₂ aba of Professor H. Kornberg, Leicester University; originally isolated at Oxford by Dr June Lascelles. ATCC 17476.
103. Strain nic SH₂ of Dr S. H. Hutner, Haskins Laboratories, N.Y. Isolated from soil by aerobic enrichment with nicotinate about 1947. ATCC 17477.
105. Strain rys-4. Isolated from soil by aerobic enrichment with tryptophan by R. Y. Stanier at Berkeley in 1950. ATCC 17479.
106. Strain tr-7 of Dr S. H. Hutner, Haskins Laboratories, N.Y. Isolated from soil by aerobic enrichment with tryptophan about 1947. ATCC 17480.
114. Strain 5 of Dr M. Shilo. Isolated from soil by aerobic enrichment with maleate at Berkeley in 1956. ATCC 17486.
125. Strain 11 of Dr M. Shilo. Isolated from soil by aerobic enrichment with mesaconate at Berkeley in 1956. ATCC 17497.
129. Strain 59 of Dr M. Shilo. Isolated from soil by aerobic enrichment with maleate at Berkeley in 1956. ATCC 17501.
146. *P. desmolytica* Gray & Thornton (1928). Authentic strain; ATCC 15005.
148. Strain c-1-o of Professor I. C. Gunsalus, University of Illinois. Origin uncertain. ATCC 17517.

(b) Strains assigned to *Pseudomonas testosteroni* Marcus & Talalay (1956)

15. Strain 21 of den Dooren de Jong (1926). Isolated from soil by aerobic enrichment with fumarate. Received from Laboratorium voor Microbiologie, Delft; ATCC 15667.

16. Strain 31 of den Dooren de Jong (1926). Isolated from soil by aerobic enrichment with bromosuccinate. Same source as strain 15. ATCC 15668.
25. *P. desmolytica* strain 1123 from Institute of Applied Microbiology, Tokyo. Isolated from soil by aerobic enrichment with anthranilate. ATCC 17407.
27. Strain KA-11 of Dr K. Hosokawa. Isolated from soil by aerobic enrichment with kynurenate in 1963 at Berkeley. ATCC 17409.
28. Strain KA-14-2. Same history as strain 27. ATCC 17410.
78. *P. testosteroni*, type strain of Marcus & Talalay (1956). Isolated at Berkeley from soil by Dr Paul Talalay, using aerobic enrichment with testosterone, about 1953. Authentic strain received via Professor I. C. Gunsalus in 1964. ATCC 11996.
79. NCTC 8893. Received from Professor S. Dagley, Leeds University, in 1964. ATCC 17454.
138. Strain FD-30 of Dr F. Delafield. Isolated from soil by aerobic enrichment with poly- β -hydroxybutyrate at Berkeley in 1961. ATCC 17510.
139. Strain FD-32. Same history as 138. ATCC 17511.

(c) Unclassified strains

60. Strain A-35 of Dr P. Thibault, Institut Pasteur. Isolated from human pleural fluid. ATCC 17437.
298. Strain 6084 of Dr P. Thibault, Institut Pasteur. ATCC 17664.

General group characters

The acidovorans group consists of non-pigmented, nutritionally versatile aerobic pseudomonads which share a distinctive nutritional spectrum and certain unique metabolic properties. All strains are multitrichous. When grown at 30° on yeast agar slopes, the strains of *Pseudomonas testosteroni* show a predominance of the monotrichous type of flagellation. The flagellar index, as defined by Lautrop & Jessen (1964), is low; but it increases considerably when the organisms are grown in liquid medium at 10–12°. According to these authors, therefore, the flagellation should be considered as multitrichous. In two strains (78, 79) it was also possible to show the formation of lateral flagella of shorter wavelength when the organisms were grown at low temperatures. Strain 78 has a higher percentage of this type of flagella.

All strains of the group can accumulate poly- β -hydroxybutyrate as a cellular reserve material. Four strains of *Pseudomonas testosteroni* (15, 16, 138, 139) attack poly- β -hydroxybutyrate by means of extracellular enzymes and use it as an exogenous carbon source; this property is absent from the rest of the group. Thus the ability to produce extracellular enzymes for the hydrolysis of poly- β -hydroxybutyrate is not a universal property in this species, as it is in *P. pseudomallei* (Redfearn *et al.* 1966) and in certain hydrogenomonads (unpublished observations). It should be noted that this property was selected in the isolation of strains 138 and 139, since they were enriched with poly- β -hydroxybutyrate as carbon and energy source.

None of the members of this group hydrolyses gelatin or gives an egg-yolk reaction; all can hydrolyse Tween 80. Growth factors are not required, and ammonia or nitrate can be used as nitrogen source. No strain can denitrify. The temperature range is relatively narrow; growth does not occur at either 4° or 41°, except for

the aberrant strain 298, which grows at 41°. Several strains grow at 37°, although poorly. The oxidase reaction is positive.

Table 38 lists the organic compounds that can be used as carbon and energy sources by 90% or more of the strains of *Pseudomonas acidovorans* and *P. testosteroni*; this list accordingly represents the common nutritional core characteristic

Table 38. *Acidovorans* group. Substrates utilized by 90% or more of the strains

Carbohydrates and sugar derivatives	Polyalcohols and glycols
Gluconate	(None)
Saccharate	Alcohols
Mucate	<i>n</i> -Butanol
Fatty acids	Non-nitrogenous aromatic and other
Acetate	cyclic compounds
Propionate	<i>m</i> -Hydroxybenzoate
Butyrate	<i>p</i> -Hydroxybenzoate
Valerate	L-Tyrosine
Isovalerate	L-Phenylalanine
Caproate	L-Kynurenine*
Dicarboxylic acids	Kynurenate*
Succinate	Aliphatic amino acids
Fumarate	Glycine*
Glutarate	L- α -Alanine*
Adipate	D- α -Alanine*
Pimelate	L-Leucine
Suberate	L-Isoleucine
Azelate	L-Aspartate
Sebacate	L-Glutamate
Hydroxyacids	L-Norleucine
D-Malate	L-Glutamate
L-Malate	Amino acids and related compounds con-
DL- β -Hydroxybutyrate	taining a ring structure
DL-Lactate	L-Histidine
Glycollate	L-Proline
DL-Glycerate	Amines
Hydroxymethylglutarate*	(None)
Miscellaneous organic acids	Miscellaneous nitrogenous compounds
Citrate	Hippurate*
Citraconate	Paraffin hydrocarbons
Itaconate	(None)
α -Ketoglutarate	
Pyruvate	
Aconitate	
Laevulinate	

* Not utilized by all strains. Negative strains are given in Tables 39, 40.

for the group. Tables 39 and 40 supplement the nutritional picture for *P. acidovorans* and *P. testosteroni*, respectively; they list all compounds used by a fraction of the strains, together with specific indications of the negative strains.

Although *Pseudomonas acidovorans* and *P. testosteroni* are nutritionally versatile, they cannot attack glucose or most other carbohydrates. As a consequence, they can be easily confused with the much less nutritionally versatile alcaligenes group when examined by customary methods. The pattern of fatty acid utilization is highly characteristic. With some exceptions for isobutyrate, the fatty acids from

acetate to caproate are excellent substrates. One strain of *P. acidovorans* grows on heptanoate, and caprate supports the growth of eight strains of the same species; caprylate and pelargonate are completely non-utilizable by both species. The dicarboxylic acids, from succinate to sebacate, support uniformly excellent growth.

Table 39. *Pseudomonas acidovorans*. Compounds used by a fraction of the strains

Substrate	Total of positive strains	Negative strains
Mannitol	14	114
meso-Inositol	7	14, 29, 61, 80, 105, 114, 125, 129
Glycerol	5	All except 24, 29, 114, 125, 146
Heptanoate	1	All except 103
Caprate	8	14, 102, 103, 105, 106, 114, 146
Isobutanol	4	All except 24, 29, 103, 148
2,3-Butyleneglycol	7	29, 61, 62, 80, 114, 125, 129, 146
D-Tryptophan	14	62
L-Kynurenine	14	62
Kynurenate	14	62
Phenol	1	All except 114
Glycine	14	114
D- α -Alanine	14	61
β -Alanine	12	105, 106, 114
DL- α -Aminobutyrate	14	61
γ -Aminobutyrate	13	24, 114
DL- α -Aminovalerate	13	61, 103

Table 40. *Pseudomonas testosteroni*. Compounds used by a fraction of the strains

Substrate	Total of positive strains	Negative strains
Isobutyrate	6	25, 28, 79
D-(—)-Tartrate	2	All except 138, 139
meso-Tartrate	4	25, 27, 28, 78, 79
Poly- β -hydroxybutyrate	4	All except 15, 16, 138, 139
Mesaconate	4	All except 16, 25, 27, 138
Hydroxymethylglutarate	8	25
n-Propanol	7	15, 16
Isopropanol	1	All except 25
Geraniol	2	79, 138
Benzoylformate	8	15
Benzoate	7	15, 25
Phthalate	2	All except 138, 139
Terephthalate	2	All except 138, 139
Nicotinate	6	25, 27, 78
Trigonelline	6	25, 27, 78
L- α -Alanine	7	15, 16
L-Valine	7	All except 25
DL- α -Aminovalerate	2	All except 25, 27
Hippurate	8	15

Most strains grow on five organic acids (glycollate, citraconate, itaconate, mesaconate, hydroxymethylglutarate) which are not widely utilized by other aerobic pseudomonads, and never all attacked by one strain of other species. The pattern

with respect to aromatic acids is likewise unusual. Two aromatic acids, *m*- and *p*-hydroxybenzoate, are universal substrates. However, no strain can grow on anthranilate or salicylate, and benzoate is used only by some strains of *P. testosteroni*. The pattern of amino acid utilization is also distinctive. Norleucine, which is wholly unutilizable by the fluorescent pseudomonads, *P. multivorans* and *P. pseudomallei* (all nutritionally versatile organisms) supports the growth of all acidovorans strains. Fourteen out of 15 strains of *P. acidovorans* grow on DL- α -aminobutyrate,

Table 41. *Acidovorans* group. The group characters of greatest differential value in the recognition of the acidovorans group, based on the analysis of 26 strains

Characters	Number of positive strains	Ideal phenotype
1. Poly- β -hydroxybutyrate as cellular reserve material	26	+
Utilization of		
2. L-arabinose	0	—
3. D-Glucose	0	—
4. D-Galactose	0	—
5. 2-Ketogluconate	0	—
6. Saccharate	25	+
7. Pelargonate	0	—
8. Adipate, pimelate, suberate, azelate, sebacate	26	+
9. Glycollate	24	+
10. Laevulinate	26	+
11. Itaconate	26	+
12. <i>m</i> -Hydroxybenzoate	25	+
13. Norleucine	25	+
14. Putrescine	0	—

Table 42. *Acidovorans* group. Number of strains of different species or groups of species of aerobic pseudomonads conforming to the selected fourteen characters (Table 41) which define the ideal phenotype of the acidovorans group

	No. of characters of 'ideal' phenotype														
	14	13	12	11	10	9	8	7	6	5	4	3	2	1	0
Acidovorans group	24	.	1	1	.	.	.	.	.	.	.	.	.	.	.
Fluorescent group	.	.	.	.	.	.	.	.	1	24	31	44	48	27	.
Pseudomallei group	.	.	.	.	.	.	.	.	.	1	23	10	7	.	.
<i>P. multivorans</i>	.	.	.	.	.	.	.	.	13	6	.	.	.	.	.
<i>P. stutzeri</i>	.	.	.	.	.	.	.	1	8	8	.	.	.	.	.
<i>P. maltophilia</i>	.	.	.	.	.	.	.	.	.	23	.	.	.	.	.
Alcaligenes group	.	.	.	.	.	.	.	.	4	2	1	.	.	.	.
<i>P. lemoignei</i>	.	.	.	.	.	.	.	1	.	.	.	.	.	.	.

a property shared with only three other strains in our collection. In contrast, no strain can grow on serine, lysine, arginine or ornithine, and only one strain on valine. These are all amino acids commonly used by other nutritionally versatile *Pseudomonas* species. Lastly, it should be noted that the acidovorans group is completely unable to attack higher amines and *N*-methyl compounds which are generally used by other nutritionally versatile species. Although the nutritional spectrum of the acidovorans group shows some degree of overlap with the nutritional spectrum of

the fluorescent pseudomonads, *P. multivorans* and *P. pseudomallei*, in many particulars it is a complementary one.

We selected a total of fourteen characters to constitute an ideal phenotype for the acidovorans group (Table 41). Every strain of *Pseudomonas acidovorans* and *P. testosteroni* conforms perfectly to this ideal phenotype, while the two unclassified strains that we have assigned to the group deviate from it in two and three characters, respectively. This is shown in Table 42, together with the degree of conformity of the strains of all other species to the ideal phenotype of the acidovorans group. The degree of separation is very satisfactory.

Table 43. *The differences between Pseudomonas acidovorans (15 strains) and P. testosteroni (9 strains)*

GC (moles %) of DNA, average value	<i>P. acidovorans</i> 66.8	<i>P. testosteroni</i> 61.8
Growth on	(Numbers of positive strains)	
D-Fructose	15	0
Mannitol	14	0
Malonate	15	0
Maleate	15	0
L-(+)-Tartrate	15	0
Quinate	15	0
Ethanol	15	0
Phenylacetate	15	0
D-Tryptophan	14	0
L-Tryptophan	15	0
Benzoylformate	0	8
Benzoate	0	7
β -Alanine	12	0
DL- α -Aminobutyrate	14	0
γ -Aminobutyrate	13	0
δ -Aminovalerate	15	0
Acetamide	15	0
Testosterone	0	9

Differentiation of two species: Pseudomonas acidovorans and P. testosteroni

As a result of our early screening of the acidovorans group, we were inclined to consider that it contained only one nutritionally somewhat irregular species. However, the analyses of Mandel (1966) have shown that strains of this group can be sharply subdivided on the guanine+cytosine (GC) content of their DNA, some showing a value of about 67 moles % GC, and others of about 62. A closer scrutiny of the nutritional data then revealed that many of the 'irregularities' were systematically correlated with differences in GC content. The correlated differences in GC content and in nutritional properties (Table 43) evidently justify the recognition of two species. For reasons that will be discussed shortly, we assign the strains of high GC content to *P. acidovorans* and those of low GC content to *P. testosteroni*.

Nomenclatural problems and designations of type strains

The recognition of two species in the acidovorans group posed a nice nomenclatural problem. One of the original strains to which den Dooren de Jong (1926) gave the name *Pseudomonas acidovorans* (strain 14) is in the high GC group, and the other

two (strains 15, 16) in the low GC group. Since there is no objective way of determining which most clearly typifies *P. acidovorans*, we have arbitrarily decided to retain this name for strain 14; accordingly, the high GC group becomes *P. acidovorans* den Dooren de Jong, with strain 14 as the type. Our choice automatically reduces to synonymy the later specific names for strain 29 (*P. indoloxidans* Gray, 1928) and for strain 146 (*P. desmolytica* Gray & Thornton, 1928). One additional nomenclatural point requires discussion. Hugh (*ATCC Catalogue*, 7th edition, 1964) has named strain 61 (ATCC 12816) *Comamonas terrigena*, a new combination for *Vibrio terrigenus* Günther (1894). Günther's strain is long since lost, and his description of it had the inadequacy typical of that era. In our opinion, it could never be certainly recognized upon re-isolation from nature, and the specific name might well have been consigned to oblivion. However, Hugh (1962) chose to revive it as an earlier synonym for *Vibrio percolans* Mudd and Warren (1923), a strain of which (ATCC 8461) still exists, and has been designated by Hugh as the neotype. We have briefly examined this strain, and find that it cannot grow on a chemically defined medium. Whatever it may be, it is clearly not identical with our strain 61 (ATCC 12816), which Hugh also considered to be '*Comamonas terrigena*'. The epithet '*terrigena*' therefore cannot be used as an earlier synonym for '*acidovorans*'.

In the low GC group, the oldest isolates are strains 15 and 16, which den Dooren de Jong (1926) assigned to *Pseudomonas acidovorans*, a name we propose to reserve for the high GC group. Strain 25, received under the label *P. desmolytica*, does not resemble Gray & Thornton's type strain of this species (our strain 146), which falls in the high GC group, and is therefore synonymous with *P. acidovorans*. The only other name previously applied to a low GC strain, *P. testosteroni* Marcus & Talalay (1956), becomes the correct designation for this group; and the strain of Marcus & Talalay (our strain 78) is the holotype. *P. testosteroni* was named on the basis of a single striking character, its ability to grow on testosterone and related steroids. Fortunately, this nutritional character proves to be taxonomically significant, since it is possessed by all low GC strains and by no high GC strains (Table 43).

The danger of naming a species on the basis of one striking character possessed by a single isolate is well illustrated by *Pseudomonas indoloxidans* of Gray (1928), now a synonym for *P. acidovorans*. Gray's strain could oxidize indole to the water-insoluble blue compound, indigotin, which accumulated as crystals in and around the bacterial growth on an indole-containing medium. The test for indigotin formation must be conducted on a medium furnished with a utilizable carbon and energy source, since indole itself cannot so serve. Tested under these conditions, Gray's original strain (our strain 29) still forms indigotin abundantly, *but is the only strain in the acidovorans group to do so*. In all other respects, strain 29 is typical of *P. acidovorans*. We have screened all other strains in our collection for the ability to form indigotin from indole and have found only three (strains 109, 110, 187) which possess this property. All three are fluorescent pseudomonads. It is accordingly evident that this striking biochemical property has no taxonomic significance.

Ecological considerations and enrichment

The frequency with which strains of the acidovorans group have been isolated from soil by aerobic enrichment with a variety of organic compounds (see strain histories) shows that this group is widespread in soil. The nutritional data suggest

enrichment methods which should be relatively or absolutely specific for the bacteria of this group. Our own experience shows that they often predominate in aerobic enrichments with higher dicarboxylic acids and kynurenate incubated at 30°. Enrichments at the same temperature with either maleate or D-tryptophan should be highly specific for *Pseudomonas acidovorans*, since these compounds are very rarely used by strains belonging to other species of aerobic pseudomonads.

Table 44. *Properties of strains 60 and 298. Only positive nutritional characters included*

Character	Strain no.		Character	Strain no.	
	60	298		60	298
GC in DNA (moles %)	59	64	Citraconate	—	+
Growth on			Itaconate	+	+
Gluconate	+	+	Mesaconate	—	+
Acetate	+	+	α -Ketoglutarate	+	+
Propionate	+	+	Pyruvate	+	+
Butyrate	+	+	Aconitate	+	+
Valerate	+	+	Laevulinate	+	+
Isovalerate	+	+	<i>m</i> -Hydroxybenzoate	+	—
Caproate	+	+	<i>p</i> -Hydroxybenzoate	+	+
Succinate	+	+	L-Tyrosine	+	—
Fumarate	+	+	L-Phenylalanine	+	—
Glutarate	+	+	D- α -Alanine	+	—
Adipate	+	+	L-Leucine	+	+
Pimelate	+	+	L-Isoleucine	+	+
Suberate	+	+	Norleucine	+	—
Azelate	+	+	L-Aspartate	+	+
Sebacate	+	+	L-Glutamate	+	+
D-Malate	+	+	L-Proline	+	+
L-Malate	+	+			
DL- β -Hydroxybutyrate	+	+			
DL-Lactate	+	+			

Properties of unclassified strains

The two remaining strains that we assign somewhat tentatively to the acidovorans group are 60 and 298. As shown in Table 44 their DNA has a GC content of 59 (strain 60) and of 64 (strain 298) %. Both strains accumulate poly- β -hydroxybutyrate as a cellular reserve material. These two strains are unable to grow on more than 20 compounds which are used by *Pseudomonas testosteroni*, so that they give the appearance of being physiological mutilates which have lost a large number of enzyme systems basic to the acidovorans group, but still retain a few of its distinctive properties. The most serious objection to regarding them as defective strains of *P. testosteroni* is that the GC content of their DNA is not identical with that of typical *P. testosteroni* strains.

THE ALCALIGENES GROUP

Seven strains of our collection have been assigned to this group, which we divide into two species: *Pseudomonas alcaligenes* (one strain), and *P. pseudoalcaligenes* (six strains).

Origins of the strains

- 63. Received from Dr P. Thibault, Institut Pasteur, Paris, as strain x-59. Isolated 1956 from sinus discharge. ATCC 17440.
- 65. Received from Dr P. Thibault, Institut Pasteur, Paris, as strain R-160. Isolated 1960 as contaminant of culture medium. ATCC 17442.
- 66. Received from Dr P. Thibault, Institut Pasteur, Paris, as strain 6076. ATCC 17443.
- 142. Strain 1577 of Dr R. Hugh. Isolated from swimming pool water. Suggested as neotype (Hugh & Ikari, 1964). ATCC 14909.
- 297. Received from Dr P. Thibault as strain 55.111. Isolated from rabbit blood culture. ATCC 17663.
- 299. Received from Dr P. Thibault, NCTC 8769. ATCC 17665.
- 417. *P. alcaligenes* ATCC 12815.

General group characters

Monias (1928) briefly described a group of non-pigmented pseudomonads, unable to liquefy gelatin or produce acid from sugars, to which he gave the name *Pseudomonas alcaligenes*. Similar strains were encountered by Thibault (1961) in a systematic study of bacteria which presented the physiological characters of the genus *Alcaligenes*. Thibault showed that the category of motile Gram-negative rods which are oxidase-positive and do not hydrolyse gelatin or produce acid from sugars includes both peritrichous and polar forms. In effect, they are distinguishable in terms of conventional diagnostic criteria only by the mode of insertion of the flagella. The 16 polarly flagellated strains studied by Thibault could, in turn, be subdivided on the basis of the number of flagella: some were monotrichous, others multitrichous. Through the kindness of Dr P. Thibault, we have been able to examine many of his polarly flagellate strains.

Ikari & Hugh (1963) examined by conventional taxonomic tests 12 strains which they considered to be representative of *Pseudomonas alcaligenes* Monias; they proposed a neotype strain for the species (Hugh & Ikari, 1964). Even in terms of an extended series of tests, this organism was defined almost entirely by negative physiological criteria: the only positive physiological criteria for its recognition given by Hugh & Ikari (1964) were the presence of catalase, ability to attack citrate, to reduce nitrate to nitrite and to give a positive oxidase reaction. These authors also showed that the flagellation was monotrichous.

The characteristics of the acidovorans group as revealed by conventional diagnostic tests are very similar to those of the alcaligenes group, and the two cannot in fact be distinguished by such cultural and physiological properties as attack on gelatin, oxidase reaction, and reactions in the carbohydrate medium of Hugh & Leifson (1953). This was revealed when we examined the collection of strains received from Dr P. Thibault under the tentative designation of *Pseudomonas alcaligenes*. Two were typical strains of *P. acidovorans*, and two were atypical strains of the acidovorans group. There remained a residue of strains which did not possess the characters of the acidovorans group. We have compared these strains with the neotype strain of *P. alcaligenes* proposed by Hugh & Ikari (strain 142), and with one other strain which had been studied by Hugh & Ikari (strain 417).

Table 45. *Alcaligenes* group. Substrates utilized by all strains

Carbohydrates and sugar derivatives (None)	Alcohols (None)
Fatty acids	Non-nitrogenous aromatic and other cyclic compounds (None)
Acetate	Aliphatic amino acids
Propionate	L- α -Alanine
Butyrate	β -Alanine
Caprylate	L-Leucine
Pelargonate	L-Glutamate
Caprate	γ -Aminobutyrate
Dicarboxylic acids	δ -Aminovalerate
Succinate	Amino acids and related compounds con- taining a ring structure
Fumarate	(None)
Hydroxyacids	Amines
L-Malate	Spermine
DL-Lactate	Miscellaneous nitrogenous compounds (None)
Miscellaneous organic acids	Paraffin hydrocarbons (None)
α -Ketoglutarate	
Pyruvate	
Polyalcohols and glycols (None)	

Table 46. *Alcaligenes* group. Substrates utilized by a fraction of the strains

Substrate	Total of positive strains	Negative strains
D-Fructose	5	142
Valerate	5	142, 417
Caproate	3	66, 142, 297, 417
Glutarate	6	142
D-Malate	6	142
DL- β -Hydroxybutyrate	6	142
DL-Glycerate	5	66, 142
Citrate	3	66, 297, 299, 417
Itaconate	6	142
Mesaconate	6	142
Glycerol	2	All except 297, 299
Propyleneglycol	5	63, 142
Ethanol	6	142
n-Propanol	6	142
n-Butanol	6	142
Glycine	3	63, 66, 142, 417
D- α -Alanine	5	142, 417
L-Serine	6	142
DL-Arginine	6	417
DL-Ornithine	1	All except 66
L-Histidine	5	66, 417
L-Proline	4	63, 66, 417
L-Tyrosine	6	142
L-Phenylalanine	6	142
Ethanolamine	6	142
Putrescine	6	142
Betaine	6	142
Sarcosine	3	63, 142, 297, 299

All these strains show monotrichous flagellation. Four of them (63, 5, 297, 299) accumulate poly- β -hydroxybutyrate as a cellular reserve material, but none of them can attack the polymer by means of extracellular enzymes. None is a denitrifier. Growth factors are not required. Both ammonia and nitrate can serve as sole nitrogen sources. All are oxidase-positive. None can grow at 4°, but all grow at 41°. although the growth of strains 3, 65 and 417 at this temperature is relatively poor, Using Skerman's method, we found a feeble attack on gelatin by strains 142, 299 and 417; Hugh & Ikari (1964) reported strain 142 to be gelatinase negative.

Table 45 gives the list of the substrates attacked by all the strains of the alcaligenes group, and Table 46 lists those substrates used by a fraction of the strains. Only 19 out of the total of 146 compounds tested are universal substrates. Beyond this, the proposed neotype strain of Hugh & Ikari (strain 142) diverges in nutritional respects from the remaining six strains, which constitute an internally homogeneous group. Strain 142 fails to grow on fructose, glutarate, D-malate, DL- β -hydroxybutyrate, itaconate, mesaconate, ethanol, *n*-propanol, *n*-butanol, L-serine, L-tyrosine, L-phenylalanine, ethanolamine, putrescine and betaine, a total of fifteen compounds used by all other strains. Citrate, used by strain 142 as reported by Hugh & Ikari (1964), is used by two of the six other strains. All strains except 417 can grow on DL-arginine, though growth of some is feeble.

Table 47. *Alcaligenes* group. The group characters of greatest differential value in the recognition of the alcaligenes group, based on the analysis of seven strains

Characters	No. of positive strains	Ideal phenotype
1. Monotrichous flagellation	7	+
2. Growth at 41°	7	+
3. Denitrification	0	—
Utilization of		
4. Glucose	0	—
5. Gluconate	0	—
6. Caprylate	7	+
7. <i>m</i> -Hydroxybenzoate	0	—
8. <i>p</i> -Hydroxybenzoate	0	—
9. β -Alanine	7	—
10. L-Serine	6	+
11. L-Aspartate	0	—
12. Spermine	7	+

The nutritional spectrum of the group recalls in many respects that of the fluorescent pseudomonads; but it is much narrower, notably in the areas of sugars and sugar-acids, and of aromatic compounds. The ability of some of the members of the group to accumulate poly- β -hydroxybutyrate, and the inability to use sugars (with the exception of fructose) are characters of the alcaligenes group also found in the acidovorans group. However, an ideal phenotype which separates the alcaligenes group from all other aerobic pseudomonads can be easily constructed (Table 47). Table 48 shows the degree of conformity of all species and groups to this ideal phenotype.

Differentiation of two species: Pseudomonas alcaligenes and P. pseudoalcaligenes, sp. nov.

In the light of the nutritional data, it seems evident that there are two distinct species in the alcaligenes group: *Pseudomonas alcaligenes*, uniquely represented in our collection by strain 142, the proposed neotype strain of Hugh & Ikari (1964); and a second species, represented by the other six strains, which is nutritionally more versatile. The data reported by Mandel (1966) for the GC content of the DNA support this conclusion: strain 142 has 66 % GC, and all other strains, 63. We propose to recognize strains 3, 65, 66, 297, 299 and 417 as a new species, *P. pseudoalcaligenes*, with strain 63 (Thibault's strain x-59) as the type strain.

Table 48. *Alcaligenes* group. Number of strains of different species and groups of species of aerobic pseudomonads conforming to the selected twelve characters (Table 47) which define the ideal phenotype of the alcaligenes group

	No. of characters of 'ideal' phenotype										
	12	11	10	9	8	7	6	5	4	3	2
Alcaligenes group	6	1	.	.	.	.	.	.	.	.	.
Fluorescent group	.	.	.	.	.	6	92	64	10	3	.
Acidovorans group	.	.	.	.	.	.	.	1	.	13	12
Pseudomallei group	.	.	.	.	.	2	15	11	9	4	.
<i>P. multivorans</i>	.	.	.	.	.	.	8	8	3	.	.
<i>P. stutzeri</i>	.	.	.	.	.	2	2	9	3	1	.
<i>P. maltophilia</i>	.	.	.	.	.	.	4	19	.	.	.
<i>P. lemoignei</i>	.	.	.	.	.	1	.	.	.	.	.

The ecology of the alcaligenes group is at present wholly obscure. Strains have been isolated only by direct plating on complex media, either from clinical specimens or from water samples.

PSEUDOMONAS MULTIVORANS SP. NOV.

We propose this name for a homogeneous and distinctive group of aerobic pseudomonads, particularly remarkable for the range and number of organic compounds that they can use as carbon and energy sources. This group was first described, but not named, by Morris & Roberts (1959). They studied 16 strains from soil and two strains from river water, isolated in Trinidad by direct plating on glucose yeast-extract peptone agar. These pseudomonads accumulate large amounts of poly- β -hydroxybutyrate, a property explored by Forsyth *et al.* (1958) and Hayward, Forsyth & Roberts (1959), who were the first to emphasize the absence of this property from fluorescent pseudomonads and its probable value in the taxonomy of the genus *Pseudomonas*. Another feature of many strains in this group is conspicuous and unusual pigmentation: according to Morris & Roberts, they form 'characteristic opaque yellow colonies which in some strains change after 2-3 days to an intense reddish-purple colour'.

Under the label of *Chromobacterium ianthinum*, one of the Trinidad strains (strain 426 of our collection) was examined by Sneath (1956, 1960) in the course of his taxonomic analysis of the genus *Chromobacterium*. He concluded that it did not belong to this genus, and did not fit any of the classical descriptions of *C. ianthinum* (Sneath, 1960). The brief characterization of the group by Morris & Roberts and the

fuller description of one strain by Sneath seem to be the only published accounts of the properties of this species. Our description is based on the study of 19 strains. Two were isolated in Berkeley; the remaining 17 included some of the original Trinidad isolates, together with other strains later isolated in Great Britain.

Origins of the strains

85. Received from Dr W. C. Haynes, Northern Regional Research Laboratory Peoria, Illinois, under the designation B 2320. Isolated by Miss M. B. Morris (at the time, curator of the Hankey Culture Collection, Trinidad) from the Maraval River, Trinidad. ATCC 17460.
104. Isolated at Berkeley from soil by Dr G. Hegeman, using D-mandelate enrichment at 30°. ATCC 17478.
249. Strain S-4-AN, isolated at Berkeley from soil using anthranilate enrichment at 41°. ATCC 17616.
382. Isolated by Dr A. C. Hayward, Commonwealth Mycological Institute, Kew, Surrey, England, in 1958 from forest soil, Trinidad (isolate 8). Other designations: Sneath D 335, and NCIB 9085. ATCC 17759.
383. Isolated by Dr A. C. Hayward in 1958 from forest soil, Trinidad (isolate 61). Other designations: NCIB 9086, and Sneath D 336. ATCC 17760.
384. Isolated by Dr A. C. Hayward in 1958 from rotting tree trunk, Trinidad (isolate 7c). Other designations: NCIB 9087 and Sneath D 337. ATCC 17761.
385. Received from Dr A. C. Hayward under the designation IMI B 2056. Isolated by Dr R. G. Mitchell from urine in urinary tract infection at Bristol Royal Infirmary, 1964. ATCC 17763.
386. Received from Dr A. C. Hayward, isolated by Dr R. G. Mitchell from baby's thermometer in Bristol Royal Infirmary. ATCC 17763.
387. Received from Dr A. C. Hayward under the designation IMI B 2572, NCIB 8507. Studied by Sneath (1960) under the designation D 338. ATCC 17764.
396. Received from Dr A. C. Hayward as strain B 2394. Isolated from urine in urinary tract infection of child, at Bristol Royal Infirmary, 1964. ATCC 17765.
- 397, 398, and 399. Received from Dr A. C. Hayward under the designations B 2443, B 2253 and B 2384. These three strains have the same origin as 396. ATCC 17766, 17767 and 17768, respectively.
420. Received from Mr W. Hodgkiss, Torry Research Station, Aberdeen, Scotland (NCIB 9088; HCC 79). From soil, Trinidad. ATCC 17769.
421. Received from Mr W. Hodgkiss (NCIB 9089 and HCC B 175). From soil, Trinidad. ATCC 17770.
422. Received from Mr W. Hodgkiss (NCIB 9091; HCC B 338). From soil, Trinidad. ATCC 17771.
423. Received from Mr W. Hodgkiss (NCIB 9092; HCC B 339). From soil, Trinidad. ATCC 17772.
424. Received from Mr W. Hodgkiss (NCIB 9093; HCC B 340). ATCC 17773.
425. Received from Mr W. Hodgkiss (Sneath TI = Sneath B 269; NCIB 9135). Isolated from forest soil in Trinidad. ATCC 17774.

General properties

All strains are multitrichous. All accumulate poly- β -hydroxybutyrate as a cellular reserve material, but are unable to attack exogenous poly- β -hydroxybutyrate. Sixteen of the 19 strains hydrolyse gelatin (negative strains are 249, 385, 398); and, with the exception of one strain (398), they give an egg-yolk reaction. All are vigorous lipase producers, as judged by the hydrolysis of Tween 80. None can denitrify. They do not require growth factors, and can use both ammonia and nitrate as nitrogen sources. They are oxidase positive. Half of the strains grow well at 41°, and all can grow at 37°. None can grow at 4°.

Morris & Roberts (1959) divided their strains into two subgroups on the basis of pigmentation: some were yellow on glucose yeast-extract peptone agar, others various shades of brown, red, violet and purple. From a strain of the latter type, they isolated a purple pigment, and identified it unequivocally as a phenazine derivative. In our experience, pigment production is a very variable character in any given strain. Purple-pigmented strains show a bewildering variety of colours when grown on solid chemically defined media with a series of different carbon sources, the pigments being sometimes exclusively associated with the organisms, sometimes also diffusible. In accord with Morris & Roberts, we find that some strains never show red or purple pigmentation, producing only yellow pigments which are partly cellular and partly diffusible. Two strains in our collection (249, 385) are completely non-pigmented. Strains 85, 383, 387, 420, 421, 422 and 425 produce the purple diffusible pigment, and strains 104, 382, 386, 396, 397, 398, 399, 423 and 424 give a yellow diffusible pigment. These two pigments are readily produced in the King A medium. Differences in pigmentation are not correlated with any of the other phenotypic characters that we have studied, and therefore seem to be devoid of taxonomic significance.

Compounds which are used by 90 % or more of the strains of *Pseudomonas multivorans* are shown in Table 49. Details for substrates utilized by a fraction of the strains are given in Table 50.

Five strains (249, 396, 397, 398, 399) lack a series of nutritional characters otherwise practically constant for the group; they cannot grow on glycollate, histamine, testosterone, acetamide or (with the exception of 249) *o*-hydroxybenzoate. Strain 385 appears to be the most defective in the whole collection; it is unable to grow at the expense of valerate, isovalerate, caproate, caprylate, caprate, D-malate, hydroxymethylglutarate, citraconate, benzoate, DL-citrulline, kynurenate, anthranilate, ethanolamine, benzylamine, spermine, butylamine, α -amylamine, sarcosine or hippurate, which are practically universal substrates for the rest of the strains.

Apart from this, the group shows a striking nutritional uniformity. In a qualitative sense, the nutritional spectrum overlaps that of the fluorescent pseudomonads, with which *Pseudomonas multivorans* also shares the *ortho* type of ring cleavage in the metabolism of benzoate and *p*-hydroxybenzoate. The only compounds utilized by a substantial number of fluorescent pseudomonads that are not utilized by *P. multivorans* are D-arabinose, erythritol, D-(–)-tartrate, itaconate, mesaconate, geraniol, propyleneglycol, glycine, L-leucine, creatine and pantothenate; but none of these compounds is a universal substrate in the fluorescent group. There is, however, a remarkable *quantitative* difference between the nutritional

Table 49. *Pseudomonas multivorans*. Substrates used by 90%
or more of the strains

Carbohydrates and sugar derivatives	Polyalcohols and glycols
D-Ribose	Mannitol
D-Arabinose	Sorbitol
L-Arabinose	meso-Inositol
D-Fucose	Adonitol
D-Glucose	Glycerol
D-Mannose	2,3-Butyleneglycol
D-Galactose	Alcohols
D-Fructose	(None)
Sucrose	Non-nitrogenous aromatic and other
Trehalose*	cyclic compounds
Cellobiose	Benzoate*
Salicin	m-Hydroxybenzoate
Gluconate	p-Hydroxybenzoate
2-Ketogluconate	Phenylacetate
Saccharate	Quinate
Mucate	Aliphatic amino acids
Fatty acids	L- α -Alanine
Acetate	β -Alanine
Butyrate	L-Threonine
Isobutyrate*	L-Aspartate
Valerate*	L-Glutamate
Isovalerate*	L-Lysine
Caproate*	DL-Arginine
Heptanoate	DL-Citrulline*
Caprylate*	γ -Aminobutyrate
Pelargonate	δ -Aminovalerate
Caprate*	Amino acids and related compounds con-
Dicarboxylic acids	taining a ring structure
Malonate	L-Histidine
Succinate	L-Proline
Fumarate	L-Tyrosine
Glutarate	L-Phenylalanine
Adipate	L-Tryptophan
Pimelate	L-Kynurenine*
Suberate	Kynurenate*
Azelate	Anthranilate*
Sebacate	Amines
Hydroxyacids	Benzylamine*
D-Malate*	Putrescine
L-Malate	Spermine*
meso-Tartrate	Tryptamine
DL- β -Hydroxybutyrate	Butylamine*
DL-Lactate	α -Amylamine*
DL-Glycerate	Miscellaneous nitrogenous compounds
Hydroxymethylglutarate*	Betaine
Miscellaneous organic acids	Sarcosine*
Citrate	Paraffin hydrocarbons
α -Ketoglutarate	(None)
Pyruvate	
Aconitate*	
Laevulinate	
Citraconate*	

* Not used by all strains. Negative strains are listed in Table 50.

Table 50. *Pseudomonas multivorans*. Substrates utilized by a fraction of the strains

Substrate	Total of positive strains	Negative strains
D-Xylose	9	85, 383, 384, 385, 387, 420, 421, 422, 423, 425
L-Rhamnose	15	421, 423, 424, 425
Trehalose	18	398
Maltose	3	All except 384, 423, 425
Propionate	16	249, 385, 387
Isobutyrate	18	249
Valerate	18	385
Isovalerate	18	385
Caproate	18	385
Caprylate	18	385
Caprate	18	385
D-Malate	17	383, 385
L-(+)-Tartrate	12	85, 383, 384, 385, 387, 420, 425
Glycollate	13	249, 385, 396, 397, 398, 399
Hydroxymethylglutarate	17	249, 385
Aconitate	18	425
Citraconate	17	249, 385
Ethanol	10	249, 383, 384, 385, 420, 421, 422, 423, 425
n-Propanol	8	249, 383, 384, 385, 387, 420, 421, 422, 423, 424, 425
n-Butanol	12	249, 420, 421, 422, 423, 424, 425
Isobutanol	7	All except 85, 104, 386, 396, 397, 398, 399
L-Mandelate	4	All except 104, 382, 386, 424
Benzoylformate	15	249, 385, 387, 421
Benzoate	17	385, 387
o-Hydroxybenzoate	14	385, 396, 397, 398, 399
Phthalate	1	All except 249
Phenol	9	249, 382, 384, 385, 386, 396, 397, 398, 399, 423
Testosterone	13	249, 385, 396, 397, 398, 399
L-Serine	16	423, 424, 425
L-Isoleucine	7	All except 85, 104, 382, 420, 422, 423, 424
L-Valine	8	85, 249, 382, 385, 387, 396, 397, 398, 399, 424, 425
DL-Ornithine	18	399
DL-Citrulline	17	249, 385
D-Tryptophan	2	All except 383, 384
L-Kynurenine	18	422
Kynurenate	18	385
Anthranilate	18	385
Ethanolamine	16	383, 384, 385
Benzylamine	17	385, 387
Spermine	18	385
Histamine	11	249, 385, 387, 396, 397, 398, 399, 421
Butylamine	18	385
α -Amylamine	18	385
Sarcosine	17	385, 422
Hippurate	16	385, 387, 421
Acetamide	13	249, 385, 396, 397, 398, 399
Nicotinate	11	104, 382, 383, 384, 385, 386, 421, 424
Trigonelline	10	104, 382, 383, 384, 385, 386, 421, 424, 425
n-Dodecane	7	All except 85, 383, 387, 420, 422, 423, 425
n-Hexadecane	7	All except 249, 383, 387, 420, 421, 422, 423

spectra of the two groups. No *single* strain of fluorescent pseudomonad that we have examined can grow on the total number of substrates that are used by the least nutritionally versatile strains of *P. multivorans* (excluding the defective strain 385). Particularly characteristic of *P. multivorans* is the very wide range of carbohydrates utilized, coupled with utilization of all the higher dicarboxylic acids and nearly all simple aromatic acids: this is a nutritional constellation which does not occur in the fluorescent group. In addition, all strains of *P. multivorans* can use a few substrates which are completely non-utilizable by the fluorescent pseudomonads that we have examined: D-fucose, cellobiose, salicin and L-threonine.

Table 51. *Pseudomonas multivorans*. The characters of greatest differential value in the recognition of *P. multivorans*, based on the analysis of 19 strains

Characters	No. of positive strains	Ideal phenotype
1. Poly- β -hydroxybutyrate as cellular reserve material	19	+
Utilization of:		
2. D-Arabinose	19	+
3. D-Fucose	19	+
4. Cellobiose	19	+
5. Saccharate and mucate	19	+
6. 2,3-Butyleneglycol	19	+
7. Sebacate	19	+
8. meso-Tartrate	19	+
9. Citraconate	17	+
10. o-Hydroxybenzoate	14	+
11. m-Hydroxybenzoate	19	+
12. L-Threonine	19	+
13. DL-Ornithine	18	+
14. Tryptamine	19	+

Table 52. *Pseudomonas multivorans*. Number of strains of different species or groups of species of aerobic pseudomonads conforming to the selected 14 characters (Table 51) which define the ideal phenotype of *Pseudomonas multivorans*

	No. of characters of 'ideal' phenotype														
	14	13	12	11	10	9	8	7	6	5	4	3	2	1	0
<i>P. multivorans</i>	13	4	2	.	.	.	.	.	.	.	.	.	.	.	.
Fluorescent group	.	.	.	.	.	.	.	.	4	12	43	74	39	3	.
Acidovorans group	.	.	.	.	.	.	.	7	12	5	.	2	.	.	.
Pseudomallei group	.	.	.	.	.	.	.	1	25	15	.	.	.	.	.
<i>P. stutzeri</i>	.	.	.	.	.	.	.	.	.	.	5	8	2	2	.
<i>P. maltophilia</i>	.	.	.	.	.	.	.	.	.	.	.	.	.	23	.
Alcaligenes group	.	.	.	.	.	.	.	.	.	.	.	.	.	5	2
<i>P. lemoignei</i>	.	.	.	.	.	.	.	.	.	.	.	.	.	1	.

Phenotypic comparison between Pseudomonas multivorans and other species of aerobic pseudomonads

In Table 51 we give a list of selected characters used for the definition of the ideal phenotype of *P. multivorans*; 13 of the strains in our collection fit this ideal phenotype in all fourteen characters, four strains coincide in thirteen of the characters, and

two strains in twelve (Table 52). All other groups and species of aerobic pseudomonads examined by us deviate from the *P. multivorans* phenotype in at least seven of the fourteen selected characters.

Proposal of type strain

As type strain for the species *Pseudomonas multivorans* we propose strain 382, one of the Trinidad isolates. This strain produces yellow diffusible pigment, liquefies gelatin, gives the egg-yolk reaction and grows at 41°. It can grow on all the substrates attacked by some *P. multivorans* strains (see Tables 49 and 50) with the exception of maltose, isobutanol, phthalate, phenol, L-valine, nicotinate, trigonelline, *n*-dodecane and *n*-hexadecane.

Ecological considerations and enrichment

It is a remarkable fact that *Pseudomonas multivorans*, by far the most nutritionally versatile aerobic pseudomonad so far examined, should have been first isolated so recently. One possible explanation might be that it is, like *P. pseudomallei* (Redfearn *et al.* 1966), a geographically restricted species, abundant only in tropical regions; this would fit with its high temperature maximum. Unfortunately, the conditions of isolation of the strains from Trinidad and Great Britain throw no light on its ecology. However, the two strains isolated in Berkeley were both obtained by enrichments, albeit unusual ones. Strain 104 emerged from a deliberate search by Dr G. Hegeman for a strain that lacked mandelic racemase, and hence could grow on L-, but not on D-mandelate. An aerobic liquid enrichment containing L-mandelate as carbon and energy source was inoculated with pond water and incubated at 30°. Colonies were isolated by plating on L-mandelate agar medium. This plate was replicated on a plate of D-mandelate agar, and a colony which failed to grow on the replica plate was isolated.

Strain 249 was isolated by one of us from an aerobic liquid enrichment with anthranilate as carbon and energy source, inoculated with soil and incubated at 41°, in the hope of discovering a specific enrichment method for *Pseudomonas aeruginosa*. This hope was unfulfilled, since from several soil samples so enriched the organism which we subsequently recognized as *P. multivorans* was isolated. The same enrichment, conducted at 30°, typically yields fluorescent pseudomonads other than *P. aeruginosa* as the predominant microflora. These facts suggest that, at 30°, *P. multivorans* is likely to be outgrown in aerobic enrichments by fluorescent pseudomonads, other nutritionally versatile aerobic pseudomonads (e.g. *P. acidovorans*, *P. testosteroni*) or members of the *Moraxella* group, even though the medium contains a carbon and energy source that is readily utilizable by *P. multivorans*. Consequently, unless some form of post-selection is applied (as in the isolation of strain 104), *P. multivorans* will probably be overlooked on plates, being a minority member of the enriched population. In enrichments at 41°, on the other hand, competition from other nutritionally versatile aerobic pseudomonads (with the exceptions of *P. aeruginosa*, *P. pseudomallei* and possibly *P. stutzeri*) is automatically eliminated. Further work on this ecological problem is obviously needed.

PSEUDOMONAS STUTZERI

A good general description of this species was published some years ago by van Niel & Allen (1952), who also retraced its chequered taxonomic history. We shall present data, based on the study of 17 strains, which supplement their description.

Origins of the strains

Strains 220 to 229 were received from Dr H. Lautrop, Statens Seruminstitut, Copenhagen. They were isolated from different clinical specimens submitted for bacteriological diagnosis.

220. Strain AB 180. ATCC 17587.

221. Strain AB 201. ATCC 17588.

222. Strain AB 205. ATCC 17589.

223. Strain AB 207. ATCC 17590.

224. Strain AB 212. ATCC 17591.

225. Strain AB 217. ATCC 17592.

226. Strain AB 316. ATCC 17593.

227. Strain AB 317. ATCC 17594.

228. Strain AB 320. ATCC 17595.

229. Strain AB 322. ATCC 17596.

275. Strain 41.4 of van Niel & Allen (1952). Received from Dr P. V. Liu, University of Louisville, Louisville, Ky. ATCC 17641.

Strains 316 to 321 were received from Dr M. Véron, Institut Pasteur, Paris; like the strains of Dr Lautrop, they were isolated from clinical specimens.

316. Strain 62C. ATCC 17682.

318. Strain 62DA. ATCC 17684.

319. Strain 62GN. ATCC 17685.

320. Strain 62HA. ATCC 17686.

321. Strain 62HB. ATCC 17687.

419. Isolated in Berkeley from soil by L-(+)-tartrate enrichment. ATCC 17832.

General properties

All strains that we have examined are monotrichous. Poly- β -hydroxybutyrate is not accumulated as a cellular reserve material; nor can it be attacked by extra-cellular enzymes when furnished as an exogenous substrate. Gelatin is not hydrolysed, and the egg-yolk reaction is negative. Most strains can hydrolyse Tween 80. Growth factors are not required, and ammonia or nitrate can serve as the nitrogen source. The temperature maximum is high; well over half the strains in our collection can grow at 41°. The oxidase test is positive.

Two outstanding properties have traditionally served to identify this species: its denitrifying ability, and its unusual colony structure. The colonies of freshly isolated strains are light brown in colour and of the rough form, wrinkled, dry and coherent. However, as van Niel & Allen (1952) noted, smooth colony variants frequently predominate after laboratory cultivation for some time. Most of the strains received by us from other laboratories proved to have a smooth colony form.

Furthermore, many of these strains grew anaerobically with nitrate only after a lag of several days; however, once vigorous growth and denitrification had occurred, subcultures into the same medium grew and denitrified promptly. Interestingly enough, several of the strains which had possessed a smooth colony form upon their receipt from other laboratories regained the rough colony form characteristic of fresh isolates from soil after one or two passages through denitrification media. These observations suggest that aerobic cultivation on complex media free of

Table 53. *Pseudomonas stutzeri*. Substrates used by 90% or more of the strains

Carbohydrates and sugar derivatives	Polyalcohols and glycols
D-Glucose	Glycerol*
Sucrose	Propyleneglycol*
Maltose*	Alcohols
Starch*	Ethanol
Fatty acids	<i>n</i> -Propanol*
Acetate	<i>n</i> -Butanol*
Butyrate*	Non-nitrogenous aromatic and other
Caproate*	cyclic compounds
Heptanoate*	(None)
Caprylate	Aliphatic amino acids
Pelargonate*	Glycine*
Caprate	L- α -Alanine
Dicarboxylic acids	D- α -Alanine
Succinate	L-Glutamate
Fumarate	Amino acids and related compounds con-
Glutarate*	taining a ring structure
Sebacate*	L-Proline*
Hydroxyacids	Amines
L-Malate	(None)
DL- β -Hydroxybutyrate	Miscellaneous nitrogenous compounds
DL-Lactate	(None)
Glycollate	Paraffin hydrocarbons
Miscellaneous organic acids	(None)
Citrate*	
α -Ketoglutarate*	
Pyruvate	
Citraconate*	
Itaconate*	
Mesaconate*	

* Not used by all strains. Negative strains are given in Table 54.

nitrate counterselects both the denitrifying ability and the rough colony form characteristic of *Pseudomonas stutzeri*, a point which should be kept in mind by those who are attempting to identify the species. In our experience, every strain can regain the capacity for active denitrification as a result of prolonged growth under conditions selective for this ability. The 'training' process can often be accelerated by initial cultivation in a nitrate-containing medium under semi-anaerobic conditions followed by transfer to a fresh tube of the same medium incubated under strictly anaerobic conditions. Some but not all strains will also regain the typical colony form as a result of this treatment.

As already mentioned in the section on the oxidase reaction and its relation to the cytochrome system, suspensions of *Pseudomonas stutzeri* show very strong

Table 54. *Pseudomonas stutzeri*. Substrates used by a fraction of the strains

Substrate	Total of positive strains	Negative strains
D-Mannose	4	All except 226, 227, 275, 318
D-Fructose	11	222, 225, 275, 316, 319, 419
Maltose	16	316
Starch	15	229 (strain 316 hydrolyses starch, but does not grow on it)
Gluconate	13	228, 319, 320, 419
Saccharate	7	All except 222, 223, 228, 229, 318, 320, 321
Mucate	7	All except 222, 223, 228, 229, 318, 320, 321
Propionate	7	All except 221, 222, 226, 227, 229, 316, 318
Butyrate	16	275
Valerate	11	220, 229, 275, 318, 321, 419
Isovalerate	14	275, 320, 321
Caproate	15	275, 320
Heptanoate	16	223
Pelargonate	16	223
Malonate	13	228, 229, 320, 419
Glutarate	16	228
Adipate	1	All except 223
Suberate	1	All except 223
Azelate	13	228, 229, 275, 419
Sebacate	15	275, 419
D-Malate	14	223, 227, 419
L-(+)-Tartrate	1	All except 419
DL-Glycerate	5	All except 223, 275, 319, 320, 419
Hydroxymethylglutarate	3	All except 222, 226, 227
Citrate	16	318
α -Ketoglutarate	16	228
Aconitate	11	222, 226, 227
Citraconate	15	223, 419
Itaconate	16	223
Mesaconate	16	223
Mannitol	9	226, 227, 316, 318, 319, 320, 321, 419
Glycerol	16	275
Ethyleneglycol	1	All except 419
Propyleneglycol	16	221
2,3-Butyleneglycol	13	226, 227, 229, 275
n-Propanol	16	221
n-Butanol	16	273
Isobutanol	12	226, 227, 229, 275, 320
Benzoate	2	All except 316, 319
Testosterone	1	All except 226, 227, 318, 419
L-Phenylalanine	3	All except 316, 319
Glycine	16	318
L-Isoleucine	13	275, 319, 321, 419
L-Valine	13	228, 229, 275, 321
L-Aspartate	13	221, 229, 275, 419
γ -Aminobutyrate	6	All except 223, 226, 227, 229, 319, 419
δ -Aminovalerate	4	All except 225, 226, 319, 419
L-Proline	16	419
L-Tyrosine	14	223, 229, 318
Kynurenate	2	All except 223, 275, 319
Ethanolamine	6	All except 220, 224, 225, 316, 319, 320
p-Hydroxybenzoate	4	All except 319
Putrescine	13	229, 316, 321, 419
Spermine	2	All except 319, 320
Betaine	1	All except 223

absorption bands of a cytochrome of the *c* type. This probably explains the characteristic colour of growth on solid media.

The substrates used by all or most strains of *Pseudomonas stutzeri* are shown in Table 53. Detailed data for substrates used by a fraction of the strains are given in Table 54. A total of 72 compounds can be used as carbon and energy sources by some strains of *P. stutzeri*; but only seventeen compounds (24% of the total) are substrates for all strains. These figures clearly indicate the marked nutritional heterogeneity of this species as compared, for instance, with a relatively homogeneous species like *P. multivorans*, for which the number of universally utilized substrates represents about 60% of the total number of compounds used by one or more of the strains of the species. The nutritional spectrum of *P. stutzeri* is very irregular; furthermore, the results of successive screenings have not always been completely reproducible. In these as in other respects, *P. stutzeri* seems to be a species prone to undergo considerable variation.

The nutritional data presented in Table 53 show, however, that *Pseudomonas stutzeri* does possess some characteristic features, although almost every nutritional character useful for the identification of the species is lacking in an occasional

Table 55. *Pseudomonas stutzeri*. The group characters of greatest differential value in the recognition of *P. stutzeri*, based on the analysis of 17 strains

Characters	No. of positive strains	Ideal phenotype
1. Monotrichous flagellation	17	+
2. Denitrification	17	+
Utilization of:		
3. L-Arabinose	0	—
4. D-Galactose	0	—
5. Cellobiose	0	—
6. Maltose	16	+
7. Starch	16	+
8. 2-Ketogluconate	0	—
9. Propyleneglycol	16	+
10. Glycollate	17	+
11. <i>p</i> -Hydroxybenzoate	3	—
12. β -Alanine	0	—
13. L-Histidine	0	—

exceptional strain. One of the diagnostically most valuable properties is the ability to grow on maltose and starch, the latter substrate being attacked by extracellular amylases. Other unusual nutritional features are the good growth on glycollate, itaconate, mesaconate and propyleneglycol. With respect to amino acids, useful features are the good growth on glycine, and the failure to grow on histidine. Higher amines and *N*-methyl compounds are rarely and sporadically used. Aromatic compounds are likewise rarely utilized; the typical irregularity of this species is shown by the fact that two strains can grow on benzoate, and three others on *p*-hydroxybenzoate.

Thirteen characters have been selected to define the ideal phenotype of *Pseudomonas stutzeri* (Table 55). As shown in Table 56, twelve strains fit the ideal phenotype in all characters, four strains deviate in one character, and 1 strain in two characters.

These thirteen characters permit a clear differentiation of *P. stutzeri* from other groups and species of aerobic pseudomonads.

In their discussion of this species, van Niel & Allen (1952) noted that denitrifying enrichments furnished with tartrate as a carbon and energy source, first used by van Iterson (1902), are particularly useful for the isolation of *Pseudomonas stutzeri*. We were therefore surprised to discover that none of the 16 strains first examined, including 275, the strain of van Niel & Allen (1952), reportedly isolated by denitrifying enrichment with tartrate, could grow on any isomer of tartrate. We therefore set up a denitrifying enrichment with L-tartrate, using a soil inoculum, which did in fact yield a strain of *P. stutzeri* (strain 419) that could grow on L-tartrate, though not on the D- and meso-isomers.

Table 56. *Pseudomonas stutzeri*. Number of strains of different species or groups of aerobic pseudomonads conforming to the selected 13 characters (Table 55) which define the ideal phenotype

	No. of characters of 'ideal' phenotype												
	13	12	11	10	9	8	7	6	5	4	3	2	1
<i>P. stutzeri</i>	12	4	1	.	.	.	.	.	.	.	.	.	.
Fluorescent group	.	.	.	.	.	1	1	27	10	48	35	16	37
Acidovorans group	.	.	.	.	.	.	11	3	12	.	.	.	.
Pseudomallei group	.	.	.	.	.	.	.	.	.	26	6	4	5
<i>P. multivorans</i>	.	.	.	.	.	.	.	.	.	.	.	3	10
<i>P. maltophilia</i>	.	.	.	.	.	.	.	23	.	.	.	.	.
Alcaligenes group	.	.	.	.	.	1	3	2	1	.	.	.	.
<i>P. lemoignei</i>	.	.	.	.	.	1	.	.	.	.	.	.	.

As Palleroni & Doudoroff (1965) pointed out, some investigators have confused *Pseudomonas stutzeri* with a hydrogenomonad, *P. saccharophila*. This confusion is probably caused by similarities in carbohydrate utilization, specifically the ability to utilize starch. These two species are, however, readily distinguishable by other characters, notably denitrifying ability (not possessed by *P. saccharophila*) and H_2 chemolithotrophy (not possessed by *P. stutzeri*). *P. stutzeri* also resembles *P. pseudomallei* in some respects (colony structure, denitrifying ability, high temperature maximum). However, it can be readily distinguished from *P. pseudomallei* by its lesser nutritional versatility and particularly by its failure to accumulate poly- β -hydroxybutyrate, or to decompose this polymer by means of extracellular enzymes.

The strain of van Niel & Allen (strain 275) is atypical of the species (or has become atypical) in several respects. It is therefore not a desirable neotype strain, and we propose instead strain 221.

DNA base composition and internal subdivision of Pseudomonas stutzeri

The GC content of the DNA of our strains of *Pseudomonas stutzeri* has been analysed by Mandel (1966). The analyses revealed the existence of two clearly differentiated subgroups within the species. Strains 220, 223, 224, 225, 228 and 320 have GC contents that average 62 moles %, while strains 221, 222, 226, 227, 229, 275, 316, 318, 319, 321 and 419 show values which cluster around 65 %. These differences in GC content suggested that two species might be combined in *P. stutzeri*. Unfortunately, the difference in GC content is not correlated with a large number of pheno-

typic differences, as in the acidovorans group. The one nutritional character which shows a high degree of correlation with GC content is the ability to grow on propionate, possessed only by strains with 65 moles % GC; but it is absent from strains 275, 319, 321 and 419, all of which do have 65 moles % GC. The ability to grow at high temperatures also appeared to show some correlation with GC content, and we investigated this aspect of the problem further. In attempting to determine the maximum temperature of growth, we found that none of the strains of *P. stutzeri* grow at 45°, but that five strains (221, 222, 229, 316, 318) grew at 44°. These five strains have 65 moles % GC in their DNA. The temperature that seems to define most clearly the two groups of strains is 43°; all high GC strains, with the exception of 275 and 419, can grow at this temperature; none of the strains of low GC content can do so. Strain 419 can grow at 42°; but so does strain 228, which has 62 moles % GC. As can be seen from the nutritional data presented in Tables 53 and 54, strain 419 is highly atypical in nutritional respects. It cannot grow on fructose, gluconate, valerate, malonate, azelate, sebacate, D-malate, citraconate, aconitate, isoleucine, aspartate, proline or putrescine, which are all attacked by the majority of other strains. On the other hand, it is the only strain capable of utilizing ethyleneglycol and L-(+)-tartrate for growth.

Possibly other correlations between phenotypic characters and DNA composition in the strains assigned to *Pseudomonas stutzeri* may be revealed by further investigation. However, it would certainly be premature to split *P. stutzeri* into two species on the basis of the minor and unreliable phenotypic differences just discussed; and we are likewise reluctant to do so purely on the basis of differences in the GC content of the DNAs.

PSEUDOMONAS MALTOPHILIA

This species was named by Hugh & Ryschenkow (1960), who subsequently (1961) published a detailed description of it, based on conventional taxonomic criteria. We have examined 23 strains, consisting of the type strain (strain 67) and a large collection of strains which had been isolated in France or French dependencies, and identified as *Pseudomonas maltophilia* by Thibault (1961).

Origins of the strains

All these strains were received from Dr P. Thibault, Institut Pasteur, Paris, under the designation *Pseudomonas maltophilia*.

67. Strain 6077. Isolated by Hugh & Ryschenkow (1961) from the oropharyngeal region of patient with mouth cancer and described as the type strain. Strain 810-2 of Hugh. ATCC 13637.
68. Strain R-72. Isolated as contaminant of a streptomycin solution. ATCC 17444.
69. Strain X-53. Isolated from pleural pus. ATCC 17445.
70. Strain CB. Obtained from cerebrospinal fluid. ATCC 17446.
71. Strain R-173. Isolated from contaminated tissue culture. ATCC 17447.
72. Strain Hamze. From blood culture. ATCC 17448.
73. Strain X-55. Obtained from a case of arthritis. ATCC 17449.
74. Strain X-80. Isolated from human lung. ATCC 17450.
75. Strain R-123. Isolated from blood culture. ATCC 17451.
300. Strain X-56. Isolated from contaminated tissue culture. ATCC 17666.
301. Strain X-57. Isolated from contaminated tissue culture. ATCC 17667.

- 302. Strain x-61. Isolated from sinusitis pus. ATCC 17668.
- 303. Strain R-41. Origin unknown. Received from French Equatorial Africa. ATCC 17669.
- 304. Strain R-147. Obtained from contaminated tissue culture. ATCC 17670.
- 305. Strain x-51. Isolated from blood culture. ATCC 17671.
- 306. Strain x-54. Isolated from sputum. ATCC 17672.
- 307. Strain x-60. Isolated from blood culture. ATCC 17673.
- 308. Strain x-62. Obtained from cerebrospinal fluid. ATCC 17674.
- 309. Strain x-77. Isolated from tissue culture. ATCC 17675.
- 310. Strain x-83. Isolated from sinusitis pus. ATCC 17676.
- 311. Strain x-57. Isolated from otitis pus. ATCC 17677.
- 312. Strain R-122. Isolated from blood culture. ATCC 17678.
- 313. Strain R-146. Isolated from tissue culture. ATCC 17679.

General properties

Pseudomonas maltophilia is multitrichous. It does not accumulate poly- β -hydroxybutyrate as a cellular reserve material, and cannot attack exogenous polymer. Nutritionally, *P. maltophilia* has proved to be a remarkably uniform biotype, which possesses several very distinctive characters not revealed by previous conventional taxonomic studies (Hugh & Ryschenkow, 1960, 1961). Every strain requires methionine, and only methionine, as growth factor; this requirement has been independently established by Iizuka & Komagata (1964) for five strains of Japanese origin. It is, furthermore, the only species of aerobic pseudomonad examined that cannot use nitrate as a principal nitrogen source. It is also the only oxidase-negative aerobic pseudomonad that we have encountered. All strains hydrolyse gelatin and are strongly lipolytic, but do not give an egg-yolk reaction. They do not denitrify.

Some of the strains produce a yellow intracellular pigment which does not diffuse into the medium. This pigment cannot be extracted from the organisms by treatment with organic solvents; it is, however, water soluble. Although we have not further investigated this point, it seems likely that the yellow colour is due to flavins.

The range of utilizable carbon and energy sources is restricted; only 23 of the 145 compounds tested support the growth of all or most strains (Table 57). Substrates used by a fraction of the strains are listed in Table 58. Notable features of the nutritional spectrum are the ability to use the disaccharides cellobiose, maltose and lactose. In fact, *Pseudomonas maltophilia* is the only species of our collection characterized by the ability to use lactose. The strains grow on lactose rather slowly, and the growth yield does not seem to be very high when lactose is used at the same concentration as the other carbohydrates in the nutritional screening. This poor growth presumably reflects the fact that galactose cannot be attacked by *P. maltophilia*. Other nutritional properties of interest include the very limited range of fatty acids utilized (acetate, propionate, valerate); the failure to grow on glutarate, β -hydroxybutyrate or aspartate, otherwise well-nigh universal substrates for aerobic pseudomonads; and the failure to grow on polyalcohols, aromatic compounds or amines. Strain 72 is aberrant: it can grow at the expense of D-xylose, isobutyrate, ethanol, *n*-propanol and L-aspartate, but not on malonate. The type strain of the

species, which was proposed by Hugh & Leifson (1963), is 67 (ATCC 13637). It can grow on: D-glucose, D-mannose, D-fructose, sucrose, trehalose, maltose, cellobiose, salicin, lactose, acetate, propionate, valerate, malonate, succinate, fumarate, L-malate, DL-lactate, citrate, α -ketoglutarate, pyruvate, aconitate, L- α -alanine, D- α -alanine, L-glutamate, L-histidine, L-proline.

Table 57. *Pseudomonas maltophilia*. Substrates used by 90% or more of the strains

Carbohydrates and sugar derivatives	Polyalcohols and glycols
D-Glucose	(None)
D-Mannose	Alcohols
Sucrose	(None)
Trehalose	Non-nitrogenous aromatic and other cyclic
Maltose	compounds
Cellobiose	(None)
Salicin	Aliphatic amino acids
Lactose	L- α -Alanine
Fatty acids	D- α -Alanine
Acetate	L-Glutamate
Propionate	Amino acids and related compounds con-
Valerate	taining a ring structure
Dicarboxylic acids	L-Histidine
Malonate*	L-Proline
Succinate	Amines
Fumarate	(None)
Hydroxyacids	Miscellaneous nitrogenous compounds
L-Malate	(None)
DL-Lactate	Paraffin hydrocarbons
Miscellaneous organic acids	(None)
Citrate	
α -Ketoglutarate	
Pyruvate	

* Not used by all strains. Negative strains are listed in Table 58.

Table 58. *Pseudomonas maltophilia*. Substrates utilized by a fraction of the strains

Substrate	Total of positive strains	Negative strains
D-Xylose	1	All except 303
D-Fructose	15	69, 74, 300, 301, 303, 307, 310, 312
Isobutyrate	2	All except 300, 302
Malonate	22	72
Aconitate	12	68, 69, 71, 300, 301, 304, 307, 309, 310, 312, 313
Ethanol	1	All except 72
n-Propanol	2	All except 72, 310
L-Aspartate	1	All except 72

Phenotypic comparison between Pseudomonas maltophilia and other species of aerobic pseudomonads. The ideal phenotype of *P. maltophilia*, defined on the basis of a total of ten selected characters, is shown in Table 59. This selected group of characters permits a very satisfactory separation from other species of aerobic

pseudomonads (Table 60). In fact, *P. maltophilia* appears to be by far the most isolated species, in terms of overall phenotype, among the aerobic pseudomonads that we have examined.

Ecological considerations

The ecology of *Pseudomonas maltophilia* is obscure. The restricted nutritional spectrum and requirement for methionine should be serious handicaps in competition with more nutritionally versatile aerobic pseudomonads. Yet the records of its repeated isolation from clinical specimens (Hugh & Ryschenkow, 1961; Dr P. Thibault, personal communication) as well as from water and raw milk (Hugh & Ryschenkow, 1961) suggest that it is widespread in nature. Iizuka & Komagata (1964) isolated it from enrichments with kerosene and crude oil inoculated with soil from petroleum zones in Japan, and reported that fresh isolates utilized hydrocarbons, but rapidly lost this ability when grown on nutrient agar slopes. None of the strains in our collection grew on dodecane or hexadecane.

Table 59. *Pseudomonas maltophilia*. The group characters of greatest differential value in the recognition of *P. maltophilia* based on the analysis of 23 strains

Characters	No. of positive strains	Ideal phenotype
1. Methionine requirement	23	+
2. Oxidase reaction	0	—
3. Utilization of nitrate as nitrogen source	0	—
Utilization of:		
4. Lactose	22	+
5. Cellobiose	23	+
6. Aspartate	1	—
7. β -Hydroxybutyrate	0	—
8. Glutarate	0	—
9. Glycerol	0	—
10. Maltose	23	+

Table 60. *Pseudomonas maltophilia*. Number of strains of different species or groups of species of aerobic pseudomonads conforming to the selected 10 characters (Table 59) which define the ideal phenotype of *P. maltophilia*

	No. of characters of 'ideal' phenotype										
	10	9	8	7	6	5	4	3	2	1	0
<i>P. maltophilia</i>	21	2	.	.	.	.	.	.	.	.	.
Fluorescent group	.	.	.	.	.	.	.	.	1	8	166
Acidovorans group	.	.	.	.	.	.	.	.	.	21	5
Pseudomallei group	.	.	.	.	.	.	.	7	32	2	.
<i>P. multivorans</i>	.	.	.	.	.	.	.	.	3	16	.
<i>P. stutzeri</i>	.	.	.	.	.	.	.	1	3	13	.
Alcaligenes group	.	.	.	.	.	.	1	.	4	2	.
<i>P. lemoignei</i>	.	.	.	.	.	.	.	1	.	.	.

DISCUSSION

Nutritional spectra: their taxonomic utility, physiological interpretation, and limitations

The long-ignored discovery of den Dooren de Jong (1926) that the nutritional spectrum of aerobic pseudomonads includes phenotypic traits of deep taxonomic significance has been overwhelmingly substantiated by our work. Every species and group of species can use a characteristic range of organic compounds as carbon and energy sources. The fact that a strain can use a given organic compound as sole source of carbon and energy demonstrates its possession of a special complement of enzymes, necessary to convert this primary substrate into one or more members of the general intermediary metabolic pool of the cell. When the biochemistry of the pathway is known, the nutritional datum immediately provides information about the enzymic constitution of the strain in question. Present biochemical knowledge tells us, for example, that a fluorescent pseudomonad able to grow on L-tryptophan can synthesize the twelve specific enzymes which mediate the conversion of this primary substrate within the cell to L-alanine, succinate and acetyl-CoA (Behrman, 1962). The ability of a fluorescent pseudomonad to grow on L-tartrate, on the other hand, is dependent upon the synthesis of only two specific proteins: a stereospecific permease, and L-tartrate dehydrase which converts the primary substrate immediately to oxalacetate (Shilo, 1957). In other words, nutritional characters, which numerical taxonomists would treat as characters of equivalent weight, are in fact far from equivalent to one another when we examine them in terms of their underlying enzymic mechanisms. In enzymic terms (and hence also in terms of structural genes) growth on L-tryptophan is a 12-unit character, growth on L-tartrate only a 2-unit character. To make really refined phenotypic comparisons among aerobic pseudomonads by the methods of numerical taxonomy (Sokal & Sneath, 1963) we require much more biochemical information about these bacteria than we now possess; and it is largely for this reason that we have resisted the temptation to feed our data to a computer. A purely nutritional analysis can also completely conceal deep biochemical differences. Nearly all strains of the acidovorans and fluorescent groups grow well on *p*-hydroxybenzoate, and with closely similar growth yields. *Nutritionally* speaking, the two groups are phenotypically identical in this particular respect. However, the metabolic pathways which they use for the dissimilation of this substrate diverge after the first step-reaction, and knowledge of this fact, obtainable only by biochemical analysis, leads to the recognition that the members of each group can synthesize at least four specific enzymes not synthesized by the other.

The inability of a strain to use a given compound as sole source of carbon and energy for growth is subject to several possible interpretations. The strain in question may simply be sensitive to the toxic effect of the compound at the concentration used, and may in fact possess the complete array of enzymes required for its utilization. The failure of many strains of *Pseudomonas fluorescens* to grow on butyrate and isobutyrate at a concentration of 0.1 % (w/v), while growing well on the same concentration of higher and lower homologues of the fatty acid series, might well prove to be interpretable in this way.

Failure to grow on a compound may occur despite the presence in the cell of the

complete array of enzymes necessary for its metabolism, if penetration is mediated by a specific permease not synthesized by the strain in question. The failure of some nutritionally versatile species (e.g. *Pseudomonas testosteroni*, *P. acidovorans*) to grow on glucose might be attributable to impermeability.

Lastly, failure to grow on a compound may be caused by the inability to synthesize one or more of the enzymes necessary for its metabolism. If the dissimilation of a compound necessitates synthesis of a considerable number of specific enzymes, the significance that should be attached to absence of growth can be determined only by a biochemical analysis. For example, in the case of L-tryptophan, lack of growth could be explicable as the result of a failure to synthesize only the first enzyme of the specific sequence, tryptophan pyrrolase. This is probably true of *Pseudomonas aeruginosa*, since strains of this species typically grow well on L-kynurenine, the second intermediate of the tryptophan pathway, but not on L-tryptophan. Many of these strains give rise to tryptophan-positive mutants when patches are replicated on tryptophan agar. From the purely determinative standpoint, this is a nuisance, since it means that tryptophan utilization is an irregular character in *P. aeruginosa*, difficult to score in plate tests. However, for a profounder understanding of the specific phenotype it is a deeply significant fact, since it shows that the ability to use tryptophan as a carbon and energy source is immanent in the genome of the species, and requires only the appearance and selection of a mutation in a gene governing synthesis of an early enzyme to achieve full phenotypic expression. If, on the other hand, it could be shown that a species unable to grow on L-tryptophan was unable to synthesize any of the specific enzymes of the tryptophan pathway, this would suggest that the entire set of genes governing the tryptophan pathway might be absent from the genome. With respect to tryptophan metabolism, such an organism would differ radically from *P. aeruginosa*, even though both might be scored nutritionally as 'tryptophan negative'.

Accordingly, the determination of nutritional spectra, useful as these may be for practical taxonomic purposes, represents only a first approximation (and a very crude one) towards characterizing the phenotypes of the aerobic pseudomonads. In effect, our present nutritional findings point the way to a much more refined analysis of the different species at the biochemical level. At least such analyses can now be conducted, as they could not be in the past, on strains which have a meaningful taxonomic identity.

*Prospective developments: towards a biochemical taxonomy
of the aerobic pseudomonads*

As already pointed out, different species may carry out the attack on a common primary substrate in biochemically different ways. This important fact, illustrated particularly clearly by the fluorescent and acidovorans groups, deserves further discussion. The most extensive evidence for the existence of metabolic divergences between the fluorescent and acidovorans groups has been accumulated through many different biochemical studies on the oxidation of aromatic substrates. This body of work is summarized in the form of two comparative metabolic maps (Figs. 6, 7). These figures show how two commonly attacked aromatic compounds, tryptophan and *p*-hydroxybenzoic acid, are metabolized by the two groups. The main features of the metabolic map for the fluorescent group have been known for

many years. However, the metabolic map for the acidovorans group could not be previously constructed, because the biochemical data on which it is based had been obtained piecemeal with various strains, for the most part designated by those who

Fig. 6. Convergent pathways for metabolism of L-tryptophan and *p*-hydroxybenzoic acid in the fluorescent group of pseudomonads. The metabolic relationship of benzoate to these two pathways is also shown.

worked with them as '*Pseudomonas* sp.', or 'an unidentified *Pseudomonas* isolated from soil'. It is the present taxonomic analysis of many of these strains which has enabled us to give coherence to the biochemical information. For example, the existence of two divergent pathways for tryptophan metabolism among aerobic pseudomonads was definitely established through comparative studies as long ago

as 1951 by Stanier, Hayaishi & Tsuchida. These authors noted that fluorescent pseudomonads all used the pathway leading through anthranilate to β -ketoadipate. Of the strains then shown to use the pathway leading through kynurenate, three which survived have now been identified by us as *P. acidovorans* (strains 103, 105,

Fig. 7. Divergent pathways for metabolism of L-tryptophan and *p*-hydroxybenzoic acid in the *acidovorans* group of pseudomonads.

106). Stanier *et al.* (1951) also noted that the ability to oxidize D-tryptophan at a high rate is possessed only by strains that use the quinoline pathway. As we have now shown, the nutritional character of growth on D-tryptophan is a distinctive specific trait of *P. acidovorans*. Consequently, although we have not yet examined the mode of tryptophan metabolism by other strains of this species, it is reasonable to assume that they all use the quinoline pathway. The second species of the acidovorans group, *P. testosteroni*, cannot grow on either isomer of tryptophan; but all strains grow well on kynurenate, and somewhat less well on L-kynurenine. These facts suggest that *P. testosteroni* probably also synthesizes the specific enzymes of the quinoline pathway, but cannot synthesize the two enzymes (tryptophan pyrrolase, formylkynurenine formamidase) used by all aerobic pseudomonads to convert tryptophan to kynurenine.

Under the designation 'Ps. B₂ aba', strain 102 of *P. acidovorans* has been the subject of numerous biochemical investigations by H. L. Kornberg and his associates. Francis, Hughes, Kornberg & Phizackerley (1963) have found that the oxidation of L-malate by this strain is catalysed by a soluble NAD-dependent dehydrogenase, whereas in a strain of *P. putida* the same reaction is catalysed by a particle-bound oxidase. This observation suggests that some of the enzymic differences between the fluorescent and acidovorans groups may be very profound indeed, and extend deep into the central intermediary metabolism of the cell.

This work could not have been done without the aid of many people. The following individuals helped us at various times in the preparation of media, the isolation of cultures, the performance of experiments or the analysis of data: Mrs Kristin Sunada, Mrs Sharron Gentry, Mrs Rebecca Contopoulou, Miss Riyo Kunisawa, Miss Margaret Farley, Mr Michael Adelberg and Mr Henri Cohen. We are deeply grateful to them all. We should also like to acknowledge the kindness of Dr Keiichi Hosokawa, who let us use unpublished biochemical tests which he had developed and first applied; and of the late Professor David Hackett, who taught us how to determine cytochrome difference spectra.

Many scientists generously contributed strains from their own collections. Not all these donations can be acknowledged here. However, particularly valuable material was received from Drs H. Lautrop and O. Jessen (Statens Seruminstitut, Copenhagen, Denmark); Drs P. Thibault and M. Véron (Service de Microbie Technique, Institut Pasteur, Paris, France); Dr J. W. M. la Rivière (Laboratorium voor Microbiologie, Technische Hoogeschool, Delft, Holland); Mr W. Hodgkiss (Department of Scientific and Industrial Research, Torry Research Station, Aberdeen, Scotland); Dr A. C. Hayward (Commonwealth Mycological Institute, Kew, Surrey, England); Dr W. C. Haynes (Northern Regional Research Laboratory of the U.S. Department of Agriculture, Peoria, Illinois); Dr Vera Sutter (School of Dentistry, University of California, San Francisco); Dr M. P. Starr (Department of Bacteriology, University of California, Davis, California). Dr Jessen also put at our disposal his then unpublished thesis on the fluorescent pseudomonads, and Dr Lautrop provided precious advice, encouragement and warning throughout the course of the work.

This study was supported by Grant AI-1808 from the National Institutes of Health, U.S. Public Health Service, and by Grant GB-4112 from the National Science Foundation.

REFERENCES

- American Type Culture Collection, Catalogue of Cultures* (1964). 7th edition. Rockville, Maryland.
- AUDUREAU, A. (1940). Étude du genre *Moraxella*. *Ann. Inst. Pasteur*, **64**, 126.
- BACHRACH, U. (1957). The aerobic breakdown of uric acid by certain pseudomonads. *J. gen. Microbiol.* **17**, 1.
- BEHRMAN, E. J. (1962). Tryptophan metabolism in *Pseudomonas*. *Nature, Lond.* **196**, 150.
- BÜHLMANN, X., VISCHER, W. A. & BRUHIN, H. (1961). Die Identifizierung nicht Pyocyanin-bildender Stämme von *Pseudomonas aeruginosa*. *Zent. Bakt. ParasitKde* (1. Abt. Orig.), **183**, 368.
- BURRI, R. & STUTZER, A. (1895). Über Nitrat zerstörende Bakterien und den durch dieselben bedingten Stickstoffverlust. *Zentbl. Bakt. ParasitKde* (2. Abt.), **1**, 257, 350, 392, 422.
- CHESTER, F. D. (1901). *A Manual of Determinative Bacteriology*. New York: MacMillan.
- COHEN-BAZIRE, G., SISTROM, W. R. & STANIER, R. Y. (1957). Kinetic studies of pigment synthesis by non-sulphur purple bacteria. *J. cell. comp. Physiol.* **49**, 25.
- DAGLEY, S., EVANS, W. C. & RIBBONS, D. W. (1960). New pathways in the oxidative metabolism of aromatic compounds by micro-organisms. *Nature, Lond.* **188**, 560.
- DAVIS, G. H. G. & PARK, R. W. A. (1962). A taxonomic study of certain bacteria currently classified as *Vibrio* species. *J. gen. Microbiol.* **27**, 101.
- DELAFIELD, F. P., DOUDOROFF, M., PALLERONI, N. J., LUSTY, C. J. & CONTOPOULOU, R. (1965). Decomposition of poly- β -hydroxybutyrate by pseudomonads. *J. Bact.* **90**, 1455.
- DEMAIN, A. L. (1965). Contamination of commercial L-leucine preparations with methionine and cystine. *J. Bact.* **89**, 1162.
- DEN DOOREN DE JONG, L. E. (1926). *Bijdrage tot de kennis van het mineralisatieproces*. Rotterdam: Nijgh and Van Ditmar.
- DEN DOOREN DE JONG, L. E. (1927). Über Protaminophage-Bakterien. *Zentbl. Bakt. ParasitKde* (2. Abt.), **71**, 193.
- DERBY, T. J. & HAMMER, B. W. (1931). Bacteriology of butter. IV. Bacteriological studies on surface taint butter. *Res. Bull. Iowa agric. Exp. Stn*, no. 145.
- DOUDOROFF, M. (1940). The oxidative assimilation of sugars and related substances by *Pseudomonas saccharophila*, with a contribution to the problem of the direct assimilation of di- and polysaccharides. *Enzymologia*, **9**, 59.
- DOUDOROFF, M. & STANIER, R. Y. (1959). Role of poly- β -hydroxybutyric acid in the assimilation of organic carbon by bacteria. *Nature, Lond.* **183**, 1440.
- DOWSON, W. J. (1939). On the systematic position and generic names of the Gram-negative bacterial plant pathogens. *Zentbl. Bakt. ParasitKde* (2. Abt.), **100**, 177.
- DWORKIN, M. & FOSTER, J. W. (1956). Studies on *Pseudomonas methanica* (Söhngen) *nov. comb.* *J. Bact.* **72**, 646.
- FLÜGGE, C. (1886). *Die Mikroorganismen*, 2. Aufl. Leipzig: F. C. W. Vogel.
- FORSYTH, W. G. C., HAYWARD, A. C. & ROBERTS, J. B. (1958). Occurrence of poly- β -hydroxybutyric acid in aerobic Gram-negative bacteria. *Nature, Lond.* **182**, 800.
- FRANCIS, M. J. O., HUGHES, D. E., KORNBERG, H. L. & PHIZACKERLEY, P. J. R. (1963). The oxidation of L-malate by *Pseudomonas* sp. *Biochem. J.* **89**, 430.
- FUCHS, A. (1959). *On the synthesis and breakdown of levan by bacteria*. Thesis, Delft: Waltman.
- GORÉ, S. N. (1921). The cotton wool plug test for indole. *Indian J. med. Res.* **8**, 505.
- GRAY, P. H. H. (1928). The formation of indigotin from indole by soil bacteria. *Proc. R. Soc. B*, **102**, 263.
- GRAY, P. H. H. & THORNTON, H. G. (1928). Soil bacteria that decompose certain aromatic compounds. *Zentr. Bakt. ParasitKde* (2. Abt.), **73**, 74.
- GÜNTHER, C. (1894). Über einen neuen, im Erdboden gefundenen Kommabacillus. *Hyg. Rdsch.* **4**, 721.
- HAYNES, W. C. (1951). *Pseudomonas aeruginosa*—its characterization and identification. *J. gen. Microbiol.* **5**, 939.

- HAYNES, W. C. (1957). In *Bergey's Manual of Determinative Bacteriology*, 7th ed., p. 89. Baltimore: Williams and Wilkins.
- HAYNES, W. C. & RHODES, L. J. (1962). Comparative taxonomy of crystallogenic strains of *Pseudomonas aeruginosa* and *Pseudomonas chlororaphis*. *J. Bact.* **84**, 1080.
- HAYWARD, A. C., FORSYTH, W. G. C. & ROBERTS, J. B. (1959). Synthesis and breakdown of poly- β -hydroxybutyric acid by bacteria. *J. gen. Microbiol.* **20**, 510.
- HILLS, G. M. (1940). Ammonia production by pathogenic bacteria. *Biochem. J.* **34**, 1057.
- HUGH, R. (1962). *Comamonas terrigena* comb.nov. with proposal of a neotype and request for an opinion. *Int. Bull. bact. Nomencl. Taxon.* **12**, 34.
- HUGH, R. & IKARI, P. (1964). The proposed neotype strain of *Pseudomonas alcaligenes* Monias 1928. *Int. Bull. bact. Nomencl. Taxon.* **14**, 103.
- HUGH, R. & LEIFSON, E. (1953). The taxonomic significance of fermentative versus oxidative metabolism of carbohydrates by various Gram-negative bacteria. *J. Bact.* **66**, 24.
- HUGH, R. & LEIFSON, E. (1963). A description of the type strain of *Pseudomonas maltophilia*. *Int. Bull. bact. Nomencl. Taxon.* **13**, 133.
- HUGH, R. & RYSCHENKOW, E. (1960). An *Alcaligenes*-like *Pseudomonas* species. *Bact. Proc.* p. 78.
- HUGH, R. & RYSCHENKOW, E. (1961). *Pseudomonas maltophilia*, an *alcaligenes*-like species. *J. gen. Microbiol.* **26**, 123.
- HUGO, W. B. & TURNER, M. (1957). A soil bacterium producing an unusual blue pigment. *J. Bact.* **73**, 154.
- IZUKA, H. & KOMAGATA, K. (1963*a*). An attempt at grouping of the genus *Pseudomonas*. *J. gen. appl. Microbiol.* **9**, 73.
- IZUKA, H. & KOMAGATA, K. (1963*b*). Taxonomy of the genus *Pseudomonas* with special reference to their modes of metabolism of carbon compounds. *J. gen. appl. Microbiol.* **9**, 83.
- IZUKA, H. & KOMAGATA, K. (1964). Microbiological studies on petroleum and natural gas. *J. gen. appl. Microbiol.* **10**, 207.
- IKARI, P. & HUGH, R. (1963). *Pseudomonas alcaligenes* Monias 1928, a polar monotrichous dextrose non-oxidizer. *Bact. Proc.* p. 41.
- JANOTA-BASSALIK, L. & WRIGHT, L. D. (1964). Azelaic acid utilization by a *Pseudomonas*. *J. gen. Microbiol.* **36**, 405.
- JESSEN, O. (1965). *Pseudomonas aeruginosa* and other green fluorescent pseudomonads. *A taxonomic study*. Copenhagen: Munksgaard.
- KILBY, B. A. (1948). The bacterial oxidation of phenol to β -keto-adipic acid. *Biochem. J.* **43**, v.
- KING, E. O., WARD, M. K. & RANEY, D. E. (1954). Two simple media for the demonstration of pyocyanin and fluorescein. *J. Lab. clin. Med.* **44**, 301.
- KLINGE, K. & GRÄF, W. (1959). Hämolyse, Eigelb-Reaktion und Amöbenauflösung durch *Pseudomonas fluorescens*. *Zentbl. Bakt. ParasitKde* (1. Abt. Orig.), **174**, 243.
- KLUYVER, A. J., DE LEY, J. & RIJVEN, A. (1951). The formation and consumption of lactobionic and maltobionic acids by *Pseudomonas* species. *Antonie van Leeuwenhoek*, **17**, 1.
- KÜNNEMANN, O. (1898). Über denitrifizierende Mikroorganismen. *Landw. VersWes.* **50**, 65.
- LAUTROP, H. & JESSEN, O. (1964). On the distinction between polar monotrichous and lophotrichous flagellation in green fluorescent pseudomonads. *Acta path. microbiol. scand.* **60**, 588.
- LEDERBERG, J. & LEDERBERG, E. M. (1952). Replica plating and indirect selection of bacterial mutants. *J. Bact.* **63**, 399.
- LEIFSON, E. (1951). Staining, shape and arrangement of bacterial flagella. *J. Bact.* **62**, 377.
- LEIFSON, E. & HUGH, R. (1954). A new type of polar monotrichous flagellation. *J. gen. Microbiol.* **10**, 68.
- LIU, P. V. (1960). Identification of pathogenic pseudomonads by extracellular antigens. *J. Bact.* **81**, 28.
- LWOFF, A. & AUDUREAU, A. (1941). La nutrition carbonée de *Moraxella kwoffi*. *Ann. Inst. Pasteur*, **66**, 417.

- LYSENKO, O. (1961). *Pseudomonas*. An attempt at a general classification. *J. gen. Microbiol.* **25**, 379.
- MACLEOD, R. A. (1965). The question of the existence of specific marine bacteria. *Bact. Rev.* **29**, 9.
- MANDEL, M. (1966). Deoxyribonucleic acid base composition in the genus *Pseudomonas*. *J. gen. Microbiol.* **43**, 273.
- Manual of Microbiological Methods* (1957). Ed. by Committee on Bacteriological Technique of American Society for Microbiology. New York: McGraw-Hill.
- MARCUS, P. I. & TALALAY, P. (1956). Induction and purification of α - and β -hydroxysteroid dehydrogenases. *J. biol. Chem.* **218**, 661.
- MERRICK, J. M. & DOUDOROFF, M. (1961). Enzymatic synthesis of poly- β -hydroxybutyric acid in bacteria. *Nature, Lond.* **189**, 890.
- MERRICK, J. M. & DOUDOROFF, M. (1964). Depolymerization of poly- β -hydroxybutyrate by an intracellular enzyme system. *J. Bact.* **88**, 60.
- MONIAS, B. L. (1928). Classification of *Bacterium alcaligenes*, *pyocyaneum* and *fluorescens*. *J. infect. Dis.* **43**, 330.
- MORRIS, M. B. & ROBERTS, J. B. (1959). A group of pseudomonads able to synthesize poly- β -hydroxybutyric acid. *Nature, Lond.* **183**, 1538.
- MUDD, S. & WARREN, S. (1923). A readily cultivable vibrio filterable through Berkefeld 'V' candles, *Vibrio percolans* (new species). *J. Bact.* **8**, 447.
- PALLERONI, N. J. & DOUDOROFF, M. (1965). Identity of *Pseudomonas saccharophila*. *J. Bact.* **89**, 264.
- PARK, R. W. A. (1962). A study of certain heterotrophic polarly flagellate water bacteria: *Aeromonas*, *Pseudomonas* and *Comamonas*. *J. gen. Microbiol.* **27**, 121.
- PEEL, D. & QUAYLE, J. R. (1961). Microbial growth on C_1 compounds. I. Isolation and characterization of *Pseudomonas* AM 1. *Biochem. J.* **81**, 465.
- REDFEARN, M. S., PALLERONI, N. J. & STANIER, R. Y. (1966). A comparative study of *Pseudomonas pseudomallei* and *Bacillus mallei*. *J. gen. Microbiol.* **43**, 293.
- RHODES, M. E. (1959). The characterization of *Pseudomonas fluorescens*. *J. gen. Microbiol.* **21**, 221.
- RHODES, M. E. (1961). The characterization of *Pseudomonas fluorescens* with the aid of an electronic computer. *J. gen. Microbiol.* **25**, 331.
- ROSENBERG, H., ENNOR, A. H. & MORRISON, V. F. (1956). The estimation of arginine. *Biochem. J.* **63**, 153.
- SEBALD, M. & VÉRON, M. (1963). Teneur en bases de l'ADN et classification des vibrions. *Ann. Inst. Pasteur*, **105**, 897.
- SHERRIS, J. C., PRESTON, N. W. & SHOESMITH, J. G. (1957). The influence of oxygen on the motility of a strain of *Pseudomonas* sp. *J. gen. Microbiol.* **16**, 86.
- SHERRIS, J. C., SHOESMITH, J. G., PARKER, M. T. & BRECKON, D. (1959). Tests for the rapid breakdown of arginine by bacteria: their use in the identification of pseudomonads. *J. gen. Microbiol.* **21**, 389.
- SHILO, M. (1957). The enzymic conversion of the tartaric acids to oxalacetic acid. *J. gen. Microbiol.* **16**, 472.
- SIERRA, G. (1957). A simple method for the detection of lipolytic activity of microorganisms and some observations on the influence of the contact between cells and fatty substrates. *Antonie van Leeuwenhoek*, **23**, 15.
- SKERMAN, V. B. D. (1959). *A Guide to the Identification of the Genera of Bacteria*. Baltimore: Williams and Wilkins Co.
- SLADE, H. D., DOUGHTY, C. C. & SLAMP, W. C. (1954). The synthesis of high-energy phosphate in the citrulline ureidase reaction by soluble enzymes of *Pseudomonas*. *Arch. Biochem. Biophys.* **48**, 338.
- SMITH, L. (1961). Cytochrome systems in aerobic electron transport. In *The Bacteria*. Ed. by I. C. Gunsalus and R. Y. Stanier, vol. 2, p. 365. New York: Academic Press.
- SNEATH, P. H. A. (1956). Cultural and biochemical characteristics of the genus *Chromobacterium*. *J. gen. Microbiol.* **15**, 70.

- SNEATH, P. H. A. (1960). A study of the bacterial genus *Chromobacterium*. *Iowa St. J. Sci.* **34**, 243.
- SOKAL, R. R. & SNEATH, P. H. A. (1963). *Principles of Numerical Taxonomy*. San Francisco: Freeman.
- STANIER, R. Y. (1952). Indole formation from tryptophan: a correction. *J. Bact.* **64**, 893.
- STANIER, R. Y. & HAYAISHI, O. (1951). The bacterial oxidation of tryptophan: a study in comparative biochemistry. *Science*, **114**, 326.
- STANIER, R. Y., HAYAISHI, O. & TSUCHIDA, M. (1951). The bacterial oxidation of tryptophan. I. A general survey of the pathways. *J. Bact.* **62**, 355.
- STARR, M. P. (1959). Bacteria as plant pathogens. *A. Rev. Microbiol.* **13**, 211.
- STARR, M. P., BLAU, W. & COSENS, G. (1960). The blue pigment of *Pseudomonas lemonnieri*. *Biochem. Z.* **333**, 328.
- STARR, M. P. & STEPHENS, W. L. (1964). Pigmentation and taxonomy of the genus *Xanthomonas*. *J. Bact.* **87**, 293.
- STEPHENSON, M. (1939). *Bacterial Metabolism*. London: Longmans, Green and Co.
- STOCKS, P. K. & McCLESKEY, C. S. (1964). Identity of the pink-pigmented methanol-oxidizing bacteria as *Vibrio extorquens*. *J. Bact.* **88**, 1065.
- STOLP, H. & STARR, M. P. (1963). *Bdellvibrio bacteriovorus* gen. et sp.n., a predatory, ectoparasitic, and bacteriolytic microorganism. *Antonie van Leeuwenhoek*, **29**, 217.
- THIBAUT, P. (1961). A propos d'*Alcaligenes faecalis*. *Ann. Inst. Pasteur*, **100** (suppl.), 59.
- THORNLEY, M. J. (1960). The differentiation of *Pseudomonas* from other Gram-negative bacteria on the basis of arginine metabolism. *J. appl. Bact.* **23**, 37.
- TSUCHIDA, M., HAYAISHI, O. & STANIER, R. Y. (1952). The bacterial oxidation of tryptophan. IV. Analysis of two blocked oxidations. *J. Bact.* **64**, 49.
- VAN ITERSON, G. (1902). Accumulation experiments with denitrifying bacteria. *Proc. Sect. Sci. K. ned. Akad. Wet.* **5**, 148.
- VAN NIEL, C. B. & ALLEN, M. B. (1952). A note on *Pseudomonas stutzeri*. *J. Bact.* **64**, 413.
- VÉRON, M. (1961). *Pseudomonas* pigmentées. *Ann. Inst. Pasteur*, **100** (suppl.), 16.
- WILLIAMSON, D. H. & WILKINSON, J. F. (1958). The isolation and estimation of poly- β -hydroxybutyrate inclusions of *Bacillus* species. *J. gen. Microbiol.* **19**, 198.