Numerical Classification of the Tribe Klebsiellae

By SHOSHANA BASCOMB*, S. P. LAPAGE, W. R. WILLCOX
AND M. A. CURTIS

*Computer Trials Department, National Collection of Type Cultures,
Central Public Health Laboratory, Colindale Avenue, London N.W. 9*

(Accepted for publication 31 March 1971)

SUMMARY

A numerical classification study was carried out on 177 strains of *Klebsiella* and related groups. Three methods of numerical classification were used: median sorting, single linkage cluster analysis and minimum spanning tree. All three contributed to the final decision on the taxa, but yielded substantially the same results. Of the three, the median sorting, if used alone, would have provided the most information.

The validity of the genus *Klebsiella* was confirmed but the inclusion of the three recognized species of *Enterobacter* in one genus was not confirmed. The genus *Klebsiella* was divided into six taxa, one of which is proposed as *K. mobilis* synon. *Enterobacter aerogenes*. *E. cloacae* occupied a rank similar to that of the genus *Klebsiella*, while *E. liquefaciens* was most closely related to the genus *Serratia* and it is proposed to include it as *S. liquefaciens*. *Enterobacter* 'pigmentés' was found to be closely related to *Chromobacterium typhiflavum*.

INTRODUCTION

During the development of a scheme for the identification of bacteria by computer (Lapage, Bascomb, Willcox & Curtis, 1970) it was found necessary to divide the Gram-negative rod-shaped bacteria under study into taxa that would differ by at least two characters. The classification of the tribe Klebsiellae (Ewing, 1963) did not lend itself easily to such a treatment as the tribe has been divided differently by several authors.

Cowan, Steel, Shaw & Duguid (1960) studied 172 *Klebsiella* strains and proposed a division into six categories, of which *Klebsiella aerogenes*, *K. pneumoniae*, *K. ozaenae* and *K. rhinoscleromatis* were assigned specific rank, and subspecific rank was assigned to *K. edwardsii* var. *atlantae* and *K. edwardsii* var. *edwardsii*. Their study did not include the so-called 'K. oxytoca' group, and they agreed with Lautrop (1956) that this group should be removed from *Klebsiella*.

Slopek & Durlakowa (1967) studied 851 *Klebsiella* strains and their grouping was generally in agreement with that of Cowan *et al.* (1960), though the names and ranks of the groups were different as they placed the two varieties of *Klebsiella edwardsii* into different species, i.e. *K. aerogenes* var. *edwardsii* and *K. pneumoniae* var. *atlantae*. Kaluzewski (1967) studied 82 indole-positive *Klebsiella* strains and found them to be sufficiently different from *K. pneumoniae* to merit species rank as *K. oxytoca*, and he would include in this species all indole-positive strains of *Klebsiella* regardless of their ability to liquefy gelatin.

In the most recently published Report (1963) of the International Subcommittee on

* Present address: Department of Biochemistry, Imperial College, Imperial Institute Road, London S.W.7.

Taxonomy of the Enterobacteriaceae, the *Klebsiella* group (genus) is divided into four biochemical types (typical, ozaenae, rhinoscleroma and oxytocolum), while *Enterobacter* is divided into three biochemical subgroups A, B and C which are respectively synonymous with *Enterobacter cloacae*, *E. aerogenes* and *E. liquefaciens*. *Hafnia* and *Serratia* rank as separate groups (genera).

Wilson & Miles (1964) included two groups of *Klebsiella*: *K. aerogenes*, divided into five biochemical varieties (typical, oxytocolum, scleroma, ozaenae and Friedländer); and *K. cloacae*, divided into varieties A, B and C which are respectively equivalent to Hormaeche & Edwards's (1960) *Enterobacter cloacae*, *E. aerogenes* and *E. liquefaciens*. They placed *Hafnia* and *Serratia* in separate groups.

On the other hand, Fife, Ewing & Davis (1965) studied 1758 strains of the tribe Klebsielleae and divided them into the following taxa: *Klebsiella pneumoniae*, *K. ozaenae*, *K. rhinoscleromatis*, *Enterobacter cloacae*, *E. aerogenes*, *E. liquefaciens*, *E. alvei* (equivalent to *Hafnia*) and two subspecies of *Serratia marcescens*. Their circumscription of *K. pneumoniae*, which we refer to as *K. pneumoniae sensu lato*, is wider than that of Cowan *et al.* (1960), referred to as *K. pneumoniae sensu stricto*, and would include strains considered by Cowan *et al.* (1960) as *K. aerogenes*, *K. pneumoniae*, both varieties of *K. edwardsii* and Lautrop's (1956) *K. oxytocolum*.

In general, disregarding taxonomic rank, the above authors agree that the *Klebsiella* group contains 'ozaenae' and 'scleroma' groups as two distinctive taxa, but disagree about the division within the 'aerogenes/edwardsii/oxytocolum/pneumoniae' complex.

The 177 strains of Enterobacteriaceae under study did not fall conveniently into any of the above schemes, and a numerical classification was thought necessary, because previously published numerical classifications of Enterobacteriaceae have been either too general (Focht & Lockhart, 1965; Krieg & Lockhart, 1966; Goodfellow, 1967) or too specific (Lockhart & Koenig, 1965), and have not included the range of strains studied here.

The numerical classification comprised two stages, calculation of a similarity matrix and cluster analysis. Many different methods have been proposed for cluster analysis starting from a similarity matrix. At present there does not seem to be a generally agreed definition of a cluster which would lead to a single accepted method of clustering. Different methods have been applied to the same data, and the results compared to suggest the deficiencies and relative merit of the methods (Lange, Stenhouse & Offer, 1965; 't Mannetje, 1967); methods have been compared on theoretical grounds (Sokal & Sneath, 1963; Proctor, 1966; Gower, 1967), and by applying them to artificially generated data (Lange *et al.* 1965; Sneath, 1966).

Three methods of cluster analysis were used in the present study, and the approach adopted was to regard the different methods as different ways of exploring the original multidimensional data as represented by the similarity matrix. The final classification was then determined by considering the results of the different cluster analyses bearing in mind the known theoretical properties of the methods. This procedure might be criticized as introducing a subjective element into the objective approach of numerical taxonomy, but at present it does not seem possible to accept uncritically the results of any single classification method.

METHODS

Organisms. The original designation of the 177 strains and their sources are listed in Table 1; their reference numbers are shown in Fig. 2. Of these strains, 74 were named reference cultures of the family Enterobacteriaceae, and a representative selection was in-

cluded of both reference and freshly isolated strains belonging to *Klebsiella*, *Chromobacterium typhiflavum*, *Enterobacter*, *Hafnia* and *Serratia*.

Methods of testing. All strains were examined for purity, Gram reaction and ability to grow aerobically on nutrient agar, then subjected to 50 cultural and biochemical tests (Table 2). The test methods largely followed those described by Cowan & Steel, 1965 (the particular method followed when alternatives are given is shown in parentheses): motility

Table 1. *Strains used and their source**

Original designation	No. of strains tested	Source	
		NCTC	Others
<i>Chromobacterium typhiflavum</i>	5	1	4
<i>Citrobacter ballerupensis</i>	1	1	—
<i>Citrobacter freundii</i>	2	2	—
<i>Edwardsiella tarda</i>	3	3	—
<i>Enterobacter aerogenes</i>	3	3	—
<i>Enterobacter cloacae</i>	27	6	21
<i>Enterobacter liquefaciens</i>	6	—	6
<i>Enterobacter</i> 'pigmentés'†	4	—	4
<i>Enterobacteria</i>	6	—	6
<i>Escherichia adecarboxylata</i> ‡	1	—	1
<i>Escherichia alkalescens</i>	1	1	—
<i>Escherichia coli</i>	4	1	3
<i>Hafnia alvei</i>	15	3	12
<i>Klebsiella aerogenes</i>	10	5	5
<i>Klebsiella edwardsii</i> var. <i>atlantae</i>	4	2	2
<i>Klebsiella edwardsii</i> var. <i>edwardsii</i>	3	2	1
<i>Klebsiella oxytocolum</i>	9	—	9
<i>Klebsiella ozaenae</i>	5	5	—
<i>Klebsiella pneumoniae</i>	5	5	—
<i>Klebsiella rhinoscleromatis</i>	5	5	—
<i>Klebsiella</i> spp.	36	2	34
<i>Kluyvera</i> ‡	2‡	—	—
<i>Salmonella typhimurium</i>	2	—	2
<i>Salmonella typhi</i>	1	1	—
<i>Serratia marcescens</i>	12	3	9
<i>Shigella boydii</i>	2	2	—
<i>Shigella dysenteriae</i>	1	1	—
<i>Shigella flexneri</i>	1	1	—
<i>Shigella sonnei</i>	1	1	—
Total	177	56	119

* The numbers of the individual strains are given in the dendrogram, Fig. 2.

† These taxa were described by Leclerc (1962).

‡ Simulated results based on the description of the species (Asai, Okumura & Tsunoda, 1957).

NCTC = National Collection of Type Cultures.

(method 1, but examined 'wet film' preparation); catalase; oxidase; liquefaction of nutrient gelatin; Hugh & Leifson's oxidation-fermentation test; utilization of citrate (method 2, Simmons); KCN tolerance; oxidation of gluconate (method 2); utilization of malonate; deamination of phenylalanine (i.e. PPA test); urease activity (method 1); production of indole (method 2); production of hydrogen sulphide in triple sugar iron media (TSI, method 1); production of hydrogen sulphide from nutrient broth (method 3); decarboxylase reactions with arginine, lysine and ornithine (method 1); methyl red reaction; Voges-Proskauer reaction (O'Meara's method); nitrate reduction; presence of β -galactosidase (ONPG test); production of acid from carbohydrates in peptone water (PWS) with

Andrade's indicator (final concentration of glucose was 0.6 % (w/v), other carbohydrates 0.4 % (w/v)); production of gas from glucose PWS.

Tests and scoring. The tests used are given in Table 2 in three columns, and the methods of coding the test results head the columns. The first column shows the tests in which all the strains gave either a positive or a negative result. These tests were neither used in the calculation of similarities nor coded. The second column gives the tests in which four levels of scoring, +, ±, ∓, and −, were adopted, and the third column gives those tests that

Table 2. *List of tests used and method of coding*

	Tests coded on the basis of	
	Four levels of scoring	Three levels of scoring
Tests not coded	+ coded 9 ± coded 8 ∓ coded 7 − coded 0	+ coded 9 ± coded 8 − coded 0
All strains positive in	Growth on MacConkey agar	ONPG
Growth at 37°	Simmons' citrate	Gas from glucose PW
Growth at RT	KCN tolerance	Acid from PWS
Hugh & Leifson fermentative	Motility at 37°	adonitol
Catalase	Motility at RT	arabinose
Acid from glucose PW	Malonate	cellobiose
Nitrate reduction	Gluconate	dulcitol
	Indole	glycerol
	H ₂ S (nutrient broth)	inositol
	H ₂ S (TSI agar)	lactose
	Arginine	maltose
	Lysine	mannitol
All strains negative in	Ornithine	raffinose
Oxidase	Urease	rhamnose
PPA	MR at RT	salicin
Hugh & Leifson oxidative	MR at 37°	sorbitol
Hugh & Leifson alkaline	VP at RT	starch
	VP at 37°	sucrose
	Gelatin liquefaction within 28 days	trehalose
		xylose

PW = peptone water; PWS = peptone water plus appropriate sugar; RT = room temperature or incubator at 22° or 30°; PPA = deamination of phenylalanine; MR = methyl red test; VP = Voges-Proskauer test; TSI = triple sugar iron medium.

required only three levels, +, ± and −. The same coding was used in both cases, i.e. 9 for +, 8 for ±, 7 for ∓ and 0 for −. A positive result whether strong or weak was thus scored with a high relative value compared with a negative result, as it was considered that a positive reaction, even if weak (±) or doubtful (∓), indicates the presence of the enzyme systems involved while a negative result suggests their absence. The results of two tests, gelatin liquefaction after 5 days and pigment production, were used in the conventional identification but not in the numerical classification.

Calculation of similarities. The similarity between strains *i* and *j* in test *t* is calculated as

$$S_{tij} = 1 - \frac{|r_{ti} - r_{tj}|}{R_t},$$

where r_{ti} and r_{tj} are the coded test results and R_t is the range of the test, equal to 9 for a test

coded 0 to 9. If either r_{ii} or r_{ij} is unknown, $S_{tij} = 0$ and a 'no comparison' is counted. Then the overall similarity, S_{ij} , is the average of the similarities for all the individual tests,

$$S_{ij} = \frac{1}{N_{ij}} \sum_{\text{All tests } t} S_{tij},$$

where N_{ij} is the number of comparisons for the pair of strains, i.e. the total number of tests minus the number of no comparisons counted.

This measure is the similarity equivalent of the mean character difference measure of distance (Sokal & Sneath, 1963, p. 146). The measure includes negative matches, and the resulting similarities (Table 3a) are almost the same as those given by a simple matching coefficient (Sokal & Sneath, 1963, p. 133) in which $\pm$ and $\mp$ are equated with $+$ (Table 3b). However, the small allowances made for the differences between $+$, $\pm$ and $\mp$ results were judged reasonable considering the interpretation of $\pm$ and $\mp$ results for this material.

Table 3. *Similarities between two strains in a single test*

(a)					(b)				
Quantitative measure					Matching coefficient				
Result of first strain					Result of first strain				

clusters formed at a series of threshold levels. Fig. 1 shows how dendrograms derived by single linkage cluster analysis and directly from the minimum spanning tree may differ. In this paper the dendrogram derived directly from the minimum spanning tree is considered to be the result of a minimum spanning tree cluster analysis. Any differences between the two dendrograms can be reduced by using smaller threshold level steps in the single linkage clustering, but if the steps are too small there will be computational difficulties. The two methods were included in the present study to see whether the extra information provided by the minimum spanning tree method was of any significance in the final classification.

Fig. 1. Dendrograms derived from the same hypothetical data: (a) by single linkage cluster analysis using steps of 2.5%; (b) using exact similarities from the minimum spanning tree.

The median sorting method of cluster analysis is related to the average linkage clustering methods of Sokal & Michener (1958). The method is most easily explained by a geometrical representation (Gower, 1967). Objects are considered as points in a multidimensional space such that the distance between two objects with similarity S_{ij} is $\sqrt{2(1 - S_{ij})}$. Initially the two closest objects (the objects having the highest similarity), say i and j , are joined. The cluster (ij) is represented by a single point halfway between i and j , and the distances of all other points from the cluster are found on this representation. The distances of other points from i and j individually are not considered further. The smallest distance is again found and the process repeated. Thus a cluster is always represented as a single point, and when two clusters join they are again replaced by a single point halfway between them irrespective of the number of individual objects in each cluster. In similarity terms when two points i and j (which may be objects or clusters) join, the similarity between another point k and the new cluster point (ij) is given by

$$S_{k,ij} = \frac{1}{2}[S_{ki} + S_{kj} + \frac{1}{2}(1 - S_{ij})].$$

The program used presented the results of the median sorting as clusters at a number of threshold levels as in the usual form of single linkage cluster analysis.

Sokal & Sneath (1963, p. 192) have noted that single linkage cluster analysis can produce 'long' clusters in which the extreme members may have much lower similarities than the level at which the cluster forms. On the other hand, the division of the clusters formed is more definite since no two members of different clusters can have a similarity greater than the clustering level. Median sorting and similar methods tend to produce more compact

clusters and may give evidence of clusters which do not occur in single linkage cluster analysis. Such clusters may be considered to be acceptable taxa or it may be considered that the continuous spectrum suggested by single linkage cluster analysis is more realistic.

RESULTS

As a preliminary, conventional identification of all the strains was made in the Computer Laboratory based on the results of the 50 tests and other available information. In the case of collection strains the original names were maintained, whereas the identification of freshly isolated strains was based mainly on the identification tables in Cowan & Steel (1965). Strains whose test results partially agreed with the definition of a particular species of *Klebsiella* in the tables were classed as *Klebsiella* spp. The results of all strains were then subjected to the three clustering methods.

The dendrograms formed by the three methods were in general similar to each other both in the major clusters formed and in the positioning of the majority of the individual strains. Only the dendrogram (Fig. 2) derived by the median sorting method is reproduced in detail since it represents most fully the clusters which were finally chosen after comparison of all three dendrograms. The analysis of this dendrogram is discussed first, followed by an examination of the differences between the median sorting and the other two methods of clustering.

Median sorting. As could be expected (Sneath, 1966), the dendrogram produced by this method is stretched along a wider similarity range than the other two. The level of similarity at which all the strains join is as low as 62.5 %. At this level two branches exist, one of a single strain of *Serratia*, the second of a cluster containing all the remaining strains. This illustrates that a branch at any level may indicate either a single strain or that the computer has formed a cluster at a higher similarity level.

The clusters formed at the 80.0 % level may be equated with the traditional tribes within the Enterobacteriaceae, except that the strains of *Hafnia* formed a cluster distinct from those of any other tribe. The formation of clusters which might be equated with traditional genera occurs at different levels in these different tribes, and the tribe *Klebsielleae* appears to be the most homogeneous. Many of the clusters formed at the 87.5 % level can be equated with traditional species, but some clusters include strains belonging to two or more different species, and strains of some species are included in two or more different clusters. The *Klebsiella* strains of the 'aerogenes/oxytoca/edwardsii/pneumoniae' complex appear as one cluster with a similarity level of 92.5 %. Of the *Klebsiella* clusters formed at the 95.0 % level, that of *Klebsiella pneumoniae* is the most homogeneous one and the only one which can be equated with a traditional taxon, i.e. Friedländer's bacillus.

Comparison with the other two clustering methods. As can be seen from the simplified versions of the three dendrograms in Fig. 3*a*, *b* and *c*, there was overall agreement between the clusters formed by all three methods. However, as can be expected, there were differences first in the similarity levels at which all strains joined and secondly in the number of branches at each level of similarity. The number of branches appearing in each dendrogram at each level of similarity is given in Table 4. The dendrogram formed by median sorting extends over a wider similarity range than the dendrograms formed by the other two methods. The level where all the strains are joined is 62.5 % in the median sorting, 82.5 % in the single linkage cluster analysis and 83.5 % in the minimum spanning tree method. At each level of similarity the largest number of branches is found in the median sorting.

The number of strains included in the important taxa by each of the three methods of cluster analysis, the minimum level of similarity within each cluster and the maximum levels

Fig. 3. Schematic form of the dendrograms derived by (a) median sorting, (b) single linkage cluster analysis and (c) the minimum spanning tree method.

of similarity when joining other clusters are given in Table 5. Most of the definite clusters formed by median sorting were also formed by the other two methods. The level of similarity at which clusters were formed, and at which they joined other clusters, was lower for the median sorting method. The minimum levels of similarity within each cluster and the similarity levels at which clusters join are slightly higher for the minimum spanning tree method than for single linkage cluster analysis because the construction of clusters is continuous in the minimum spanning tree while it proceeds stepwise in levels of 2.5 % in the single linkage. However, the clusters formed by these two methods include a similar number of strains. Though the strains of most taxa were clustered similarly by the three methods, those of *Enterobacter liquefaciens*, *Klebsiella ozaenae*, *K. pneumoniae (sensu stricto)* and of *Klebsiella* 'unnamed group' were clustered differently.

Table 4. *Comparison of the number of branches appearing at each level of similarity in the three different methods of cluster analysis*

Level of similarity (%)	No. of branches by		
	Median sorting	Single linkage	Minimum spanning tree
62.5	2	1	1
75.0	3	1	1
77.5	3	1	1
80.0	7	1	1
82.5	10	3	1
85.0	17	10	3
87.5	31	25	18
90.0	49	44	25
92.5	83	72	44
95.0	132	114	76
97.5	177	177	177
100	177	177	177

Final construction of taxa. After comparison of the three dendrograms, taxa were formed mainly on the basis of the clusters existing in the median sorting dendrogram at the similarity level of 87.5 %. In the case of *Enterobacter cloacae*, *Hafnia alvei* and *Klebsiella ozaenae* a few more strains were added to those already included in the clusters. *Klebsiella pneumoniae (sensu stricto)* was treated as a separate taxon, though it does not exist as a separate cluster at this level.

The validity of the taxa thus formed was checked by calculating the mean similarities within and between the taxa (Fig. 4). The mean similarity within a taxon is the average of the similarities between all the pairs that can be formed among the strains of that taxon. The mean similarity between two taxa is the average of the similarities between those pairs obtained by pairing each strain of the first taxon with all the strains of the second taxon. The level of similarity within each taxon was always higher than the level of similarity between taxa. However, the taxa are not of equivalent hierarchical level as the similarities within taxa vary between 99.0 % for *Klebsiella pneumoniae (sensu stricto)* and 88.7 % for *Shigella* and the *Alcalescens-Dispar* group. The highest intertaxa similarity was found between *K. pneumoniae (sensu stricto)* and *K. pneumoniae (sensu lato)* where it was 88.0 %. The biochemical characteristics of these taxa are summarized in Table 6, which contains only 40 tests, because the tests in which all the strains gave constant results (Table 2, first column) have been omitted.

Table 5. Comparison of the levels of similarity of the clusters formed by the different clustering methods

Name of taxon	Median sorting			Single linkage sorting			Minimum spanning tree		
	Level of similarity (%)			Level of similarity (%)			Level of similarity (%)		
	No. of strains included	Minimum within cluster	Maximum when joining other clusters	No. of strains included	Minimum within cluster	Maximum when joining other clusters	No. of strains included	Minimum within cluster	Maximum when joining other clusters
<i>Chromobacterium typhiflavum</i> and <i>Enterobacter</i> 'pigmentés'	6	92·5	87·5	7	90·0	87·5	7	90·5	87·5
<i>Edwardsiella tarda</i>	3	95·0	82·5	3	95·0	82·5	3	95·5	83·5
<i>Enterobacter aerogenes</i>	3	92·5	87·5	3	92·5	90·0	3	93·5	91·5
<i>Enterobacter cloacae</i>	27	87·5	85·0	28	90·0	87·5	28	91·5	88·5
<i>Enterobacter liquefaciens</i>	5	90·0	87·5	3; 3	92·5; 90·0	87·5	3; 3	93·5; 90·5	89·5
<i>Escherichia coli</i>	4	90·0	82·5	4	90·0	85·0	4	91·5	86·0
<i>Hafnia alvei</i>	18	87·5	80·0	18	90·0	85·0	18	91·5	85·5
<i>Klebsiella ozaenae</i>	4	92·5	87·5	2; 2	92·5; 92·5	90·0	2; 2	94·0; 93·0	91·5
<i>Klebsiella pneumoniae</i>									
<i>Klebsiella pneumoniae sensu stricto</i>	6	97·5	95·0	—	—	—	4	99·5	97·5
<i>Klebsiella pneumoniae sensu lato</i>	62	92·5	87·5	67	95·0	90·0	67*	95·0	94·0
<i>Klebsiella rhinoscleromatis</i>	5	90·0	87·5	5	95·0	87·5	5	95·5	89·5
<i>Klebsiella</i> 'unnamed group'	4	90·0	87·5	2	97·5	95·0	2	99·5	95·5
<i>Serratia</i> biotype I	9	90·0	87·5	8	92·5	87·5	8	93·5	89·5
<i>Serratia</i> biotype II	2	97·5	85·0	2	95·0	87·5	2	97·5	87·5
<i>Shigella</i> spp. + A-D group	6	90·0	82·5	6	87·5	85·0	6	87·5	86·0

* This cluster includes the four strains of *K. pneumoniae* (*sensu stricto*).

Fig. 4. Mean similarities within and between the taxa formed.

DISCUSSION

The relationship between the important taxa as revealed by the three different sorting methods is shown in a shortened schematic form in Fig. 3*a, b, c*. In all three the tribe Klebsiellae appears as a separate cluster and includes the following traditional taxa: *Enterobacter*, *Klebsiella*, *Chromobacterium typhiflavum* and *Serratia*. Similarly, the strains of *Hafnia alvei* form another separate and distinct taxon with a rank similar to the tribe Klebsiellae. The strains of *Escherichia coli*, *Shigella* spp. and the Alkaescens-Dispar group form the third distinct cluster. Strains of *Edwardsiella tarda* are included in this cluster by median sorting but appear separately from all the other strains in the other two dendrograms. The other strains of *Salmonella*, *Citrobacter*, and the two simulated strains of *Kluyvera* did not form similar discrete groups. However, as very few strains of these taxa were included for comparison in this study no further conclusions are justifiable.

Divisions within the tribe Klebsiellae. In all dendrograms the tribe contains the following separate clusters: *Klebsiella*, *Enterobacter cloacae*, and a cluster formed from the strains of *Chromobacterium typhiflavum* and *Enterobacter 'pigmentés'*. Two strains of *Serratia* form a distinct cluster which appears in all dendrograms and is of the same rank as these three clusters. However, the position of strains of *E. liquefaciens* and of *Serratia* differ in the three dendrograms.

The *Klebsiella* cluster contains *Klebsiella rhinoscleromatis* and *Enterobacter aerogenes* as separate clusters in all dendrograms, while *K. ozaenae*, though separate from the remaining strains of *Klebsiella*, appears as one cluster in the median sorting dendrogram and as two in the other dendrograms.

Table 6. Characteristics of the taxa formed

Tests	<i>Klebsiella aerogenes</i> / <i>oxytoca/edwardsii</i>	<i>Klebsiella pneumoniae</i>	<i>Klebsiella</i> 'unnamed group'	<i>Klebsiella ozaenae</i>	<i>Enterobacter aerogenes</i> *	<i>Klebsiella rhinoscleromatis</i>	<i>Enterobacter cloacae</i>	<i>Enterobacter liquefaciens</i> †	<i>Serratia</i> biotype I	<i>Serratia</i> biotype II	<i>Chromobacterium typhiflavum</i> <i>Enterobacter</i> 'pigmentés'	<i>Hafnia alvei</i>	<i>Edwardsiella tarda</i>
Motility 37°	-	-	-	-	+	-	d	d	+	d	d	d	+
Motility RT	-	-	-	-	+	-	d	+	+	+	+	d	+
Growth on MacConkey agar	+	+	+	d	+	d	+	+	+	+	+	+	+
Liquefaction gelatin { 1 to 5 days 28 days	d	-	-	-	-	-	d	d	+	+	+	-	-
Growth on Simmons' citrate	+	+	-	d	+	-	+	+	+	+	d	+	-
KCN	+	-	d	d	+	+	+	+	+	+	d	+	-
Gluconate	+	d	d	-	+	-	+	+	+	+	-	+	-
Malonate	d	+	+	-	+	+	d	-	-	-	+	d	-
Urease	+	+	-	-	-	-	-	-	d	+	-	-	-
Indole	d	-	-	-	-	-	d	-	-	-	-	-	+
H ₂ S, TSI	-	-	-	-	-	-	-	-	-	-	-	-	+
H ₂ S, lead acetate paper	-	-	-	-	d	-	d	-	d	-	-	d	+
Arginine dihydrolase	-	-	-	d	-	-	+	d	-	-	-	d	-
Lysine decarboxylase	+	+	d	d	+	-	-	+	+	-	-	+	+
Ornithine decarboxylase	-	-	-	-	+	-	+	+	+	-	-	+	+
MR 37°	d	+	d	+	d	+	d	+	d	+	+	d	+
MR RT	d	+	d	+	-	+	-	d	d	-	d	-	+
VP 37°	d	-	d	-	-	-	+	+	+	+	-	d	-
VP RT	d	-	d	-	d	-	+	d	+	+	d	+	-
Gas from glucose	d	+	+	d	+	-	+	+	d	-	-	+	+
Acid from:													
adonitol	+	+	+	+	+	+	d	-	d	+	-	-	-
arabinose	+	+	+	+	+	+	+	d	-	+	+	+	d
cellobiose	+	+	+	+	+	d	+	d	-	+	d	d	-
dulcitol	d	+	-	d	-	-	d	-	-	-	+	-	-
glycerol	+	+	+	+	+	+	d	+	+	+	d	+	d
inositol	+	+	+	+	+	d	-	+	d	+	-	-	-
lactose	+	+	+	+	+	-	d	-	-	+	-	-	-
(ONPG)	+	+	+	+	+	-	+	d	+	+	+	+	-
maltose	+	+	+	+	+	+	+	d	+	+	d	+	+
mannitol	+	+	+	+	+	+	+	+	+	+	+	+	-
raffinose	+	+	+	+	+	+	+	d	-	+	-	-	-
rhamnose	+	+	+	+	+	d	+	-	-	-	+	+	-
salicin	+	+	+	+	+	+	+	+	+	+	+	-	-
sorbitol	+	+	+	+	+	+	d	d	+	+	-	d	-
starch	d	+	+	d	d	-	-	-	-	-	-	-	d
sucrose	+	+	+	+	+	+	+	+	+	+	+	-	-
trehalose	+	+	+	+	+	+	+	+	+	+	+	+	-
xylose	+	+	+	+	+	+	+	+	d	+	+	d	-

Key: + = 90 to 100% of strains yield positive results; d = 11 to 89% of strains yield positive results;
- = 0 to 10% of strains yield positive results.

* Proposed name: *Klebsiella mobilis*.

† Proposed name: *Serratia liquefaciens*.

The strains of *Klebsiella aerogenes*, *K. edwardsii*, *K. oxytoca* and *K. pneumoniae* (*sensu stricto*) appear in all three dendrograms as one large cluster. The formation of three of these traditional taxa was based (Cowan *et al.* 1960) mainly on the reactions to 13 tests, 11 of which were included in our study, while fimbriation and production of gas from lactose at 44° were not. Moreover, the incubation conditions for the methyl red and Voges–Proskauer tests as well as the method for detecting acetoin were different in the two studies. We found that strains belonging to the traditional taxa gave different results in the methyl red and Voges–Proskauer tests and in the production of gas from glucose from those expected for these taxa. Our results did not confirm the use of these three tests in the separation of the strains into the traditional species. It is possible that if fimbriation and production of gas from lactose at 44° had been included in our study the separation would have been different. However, these additions would only have increased the number of tests from 49 to 51 and these two tests were not necessarily applicable to the differentiation of the other taxa in our study. Attempts to form taxa within this cluster of klebsiellas, either on the basis of traditional classification or on the basis of clusters formed by our numerical techniques, were successful only in the case of *K. pneumoniae* (*sensu stricto*). All other taxa were not homogeneous enough in their test results and therefore not sufficiently different from each other. It would appear that the strains fall more easily into a continuous spectrum and do not form discrete groups. This is particularly applicable to strains of *K. oxytoca* and *K. aerogenes*. *Klebsiella oxytoca* differs from *K. aerogenes* in that the former produces indole and liquefies gelatin while the latter is unable to do either. However, strains were found which were similar to *K. aerogenes* yet gave a positive result in one or other, or both of these tests. Hence there does not appear to be a distinction between these two species.

The four freshly isolated strains that formed a distinct cluster in the median sorting dendrogram (*Klebsiella* ‘unnamed group’) do not form distinct clusters in the other two. However, the within-group and between-group mean similarity values (93.6 % and 86.6 % respectively) would lend support to the validity of this taxon. We therefore suggest that the genus *Klebsiella* should contain the following six taxa:

- Klebsiella* I = *K. aerogenes*/oxytoca/edwardsii
- II = *K. pneumoniae*
- III = *Klebsiella* ‘unnamed group’
- IV = *K. ozaenae*
- V = *Enterobacter aerogenes*
- VI = *K. rhinoscleromatis*

and postpone the naming of group III until its validity has stood the test of time. We consider that *Enterobacter aerogenes* should be classified in the genus *Klebsiella* and propose for this species the name *K. mobilis nomen novum* since the specific epithet ‘aerogenes’ is already in use for *K. aerogenes* (Kruse) Taylor *et al.* 1956. Strain CDC 819–56 (NCTC 10006, ATCC 13048), which was included in this study, will thus be the type strain as it is the proposed neotype for *Enterobacter aerogenes* (Hormaeche & Edwards, 1960). We regard the name *Aerobacter aerogenes* (Kruse) Beijerinck 1900 as a *nomen dubium* as it has been widely used for two different organisms and hence in making our proposal no question of priority with *Aerobacter aerogenes* arises. A discussion of the confusion caused by the name *Aerobacter* is given by Carpenter *et al.* (1970). The characteristics of *K. mobilis* are given in Table 6 and can be compared with those of *K. aerogenes*.

The validity of the inclusion of *Enterobacter aerogenes*, *E. cloacae* and *E. liquefaciens* in the genus *Enterobacter* is not brought out by our classification. *Enterobacter aerogenes* was included in *Klebsiella* in spite of the fact that motility was recorded twice for each strain, i.e. at 37° and at room temperature. *Enterobacter liquefaciens* seems to be more related to *Serratia* biotype I in two of the dendrograms. We therefore propose that *E. liquefaciens* be included in *Serratia*, which would therefore contain at least the two species *S. marcescens* and *S. liquefaciens* comb.nov. A type strain is not suggested since the type strain for *Enterobacter liquefaciens* (ATCC 14460, NCIB 9321; Grimes & Hennerty, 1931), was not included in this study. The characteristics of the taxon are given in Table 6. This proposal is supported by the findings of Hamon, Le Minor & Peron (1970) that strains of *E. liquefaciens* and *S. marcescens* show sensitivity to the same bacteriolytic agents. It would also appear that *E. cloacae* is sufficiently distinct from the other two species and might merit genus rank.

The inclusion in *Serratia* of all red-pigmented Gram-negative rods is also shown to be doubtful, as we could discern at least two biotypes: biotype I which seems to be related to *Enterobacter liquefaciens*, and biotype II which shows different reaction patterns. The characteristics of these biotypes are given in Table 6. *Serratia* biotype I resembles *S. marcescens* subsp. *marcescens* of Fife *et al.* (1965); it also resembles pattern 1 of Fulton, Forney & Leifson (1959) though less closely. The strains of *Serratia* biotype II differ from the description of *Serratia marcescens* subsp. *kiliensis* (Fife *et al.* 1965), by giving positive results in the Voges-Proskauer test at both 37° and room temperature, and by producing acid from arabinose and raffinose. *Serratia* biotype II also differs from pattern 2 of Fulton *et al.* (1959) in producing urease, and acid from adonitol. Not enough strains of this organism have been studied to discuss its relationship to other described species of *Serratia*.

Hafnia alvei is sufficiently distinct to be excluded from the tribe Klebsiellae and perhaps should assume a tribe rank on its own, similar to Escherichiaeae and Klebsiellaeae.

Although in our tests *Chromobacterium typhiflavum* and *Enterobacter* 'pigmentés' appear to be the same and to belong to *Erwinia herbicola*, we have kept the original strain designations throughout this paper because there is no published comparison of the two species, *C. typhiflavum* and *Enterobacter* 'pigmentés', although Graham & Hodgkiss (1967) have shown that *C. typhiflavus* is the same as *E. herbicola*. The one strain of *Escherichia adecarboxylata*, also a yellow-pigmented species, included in our study appeared within the *Chromobacterium typhiflavum* cluster in the median sorting dendrogram but as a separate cluster in the other two. A more detailed study of the yellow-pigmented enterobacteria might show that these three species are closely related to each other and might occupy a genus rank, equivalent to that of *Klebsiella*.

The computer work was done at Rothamsted Experimental Station, Harpenden, Hertfordshire, using the Orion classification program named CLASP, and we should like to thank Mr J. C. Gower and Mr G. J. S. Ross for their help and advice; and the late Dr K. P. Carpenter of the Dysentery Reference Laboratory, Colindale, for providing reference strains. We also particularly wish to thank Mrs M. Della-Valle and Miss Joyce Stevens for the preparation of the manuscript.

Our thanks are due to the Department of Health and Social Security whose grant for the identification of bacteria using a computer enabled us to carry out this work.

REFERENCES

- ASAI, T., OKUMURA, S. & TSUNODA, T. (1957). On the classification of the α -ketoglutaric acid-accumulating bacteria in aerobic fermentation. *Journal of General and Applied Microbiology, Tokyo* **3**, 13-33.
- BEIJERINCK, M. W. (1900). Schwefelwasserstoffbildung in den Stadtgräben und Aufstellung der Gattung *Aërobacter*. *Zentralblatt für Bakteriologie, Parasitenkunde, Infektionskrankheiten und Hygiene II Abteilung* **6**, 193-206.
- CARPENTER, K. P. *et al.* (1970). Request to the Judicial Commission that *Aerobacter* Beijerinck 1900 and *Aerobacter* Hormaeche & Edwards 1958 be declared rejected generic names. *International Journal of Systematic Bacteriology* **20**, 221-224.
- COWAN, S. T. & STEEL, K. J. (1965). *Manual for the Identification of Medical Bacteria*. Cambridge University Press.
- COWAN, S. T., STEEL, K. J., SHAW, C. & DUGUID, J. P. (1960). A classification of the Klebsiella group. *Journal of General Microbiology* **23**, 601-612.
- EWING, W. H. (1963). An outline of nomenclature for the family Enterobacteriaceae. *International Bulletin of Bacteriological Nomenclature and Taxonomy* **13**, 95-110.
- FIFE, M. A., EWING, W. H. & DAVIS, R. B. (1965). *The Biochemical Reactions of the Tribe Klebsiellae* CDC publications, Communicable Disease Center, Atlanta, Georgia.
- FOCHT, D. D. & LOCKHART, W. R. (1965). Numerical survey of some bacterial taxa. *Journal of Bacteriology* **90**, 1314-1319.
- FULTON, M., FORNEY, C. E. & LEIFSON, E. (1959). Identification of *Serratia* occurring in man and animals. *Canadian Journal of Microbiology* **5**, 269-275.
- GOODFELLOW, M. (1967). Numerical taxonomy of some named bacterial cultures. *Canadian Journal of Microbiology* **13**, 1365-1374.
- GOWER, J. C. (1967). A comparison of some methods of cluster analysis. *Biometrics* **23**, 623-637.
- GOWER, J. C. & ROSS, G. J. S. (1969). Minimum spanning trees and single linkage cluster analysis. *Applied Statistics* **18**, 54-64.
- GRAHAM, D. C. & HODGKISS, W. (1967). Identity of Gram-negative, yellow pigmented, fermentative bacteria isolated from plants and animals. *Journal of Applied Bacteriology* **30**, 175-189.
- GRIMES, M. & HENNERTY, A. J. (1931). A study of bacteria belonging to the subgenus *Aerobacter*. *Scientific Proceedings of the Royal Dublin Society* **20**, 89-97.
- HAMON, Y., LE MINOR, L. & PERON, Y. (1970). Les bactériocines d'*Enterobacter liquefaciens*. Intérêt taxonomique de leur étude. *Compte rendu hebdomadaire des séances de l'Académie des sciences, Paris* **270**, série D, 886-889.
- HORMAECHÉ, E. & EDWARDS, P. R. (1960). A proposed genus, *Enterobacter*. *International Bulletin of Bacteriological Nomenclature and Taxonomy* **10**, 71-74.
- KALUZEWSKI, S. (1967). Taxonomic position of indole-positive strains of Klebsiella. *Experimental Medicine and Microbiology* (formerly *Medycyna Doswiadczalna i Mikrobiologia*) **19**, 350-359.
- KREIG, R. E. & LOCKHART, W. R. (1966). Classification of enterobacteria based on overall similarity. *Journal of Bacteriology* **92**, 1275-1280.
- LANGE, R. T., STENHOUSE, N. S. & OFFLER, C. E. (1965). Experimental appraisal of certain procedures for the classification of data. *Australian Journal of Biological Sciences* **18**, 1189-1205.
- LAPAGE, S. P., BASCOMB, S., WILLCOX, W. R. & CURTIS, M. A. (1970). Computer identification of bacteria. In *Automation, Mechanization and Data Handling in Microbiology*, pp. 1-22. Edited by A. Baillie & R. J. Gilbert. Society for Applied Bacteriology Technical Series no. 4.
- LAUTROP, H. (1956). Gelatin liquefying Klebsiella strains (*Bacterium oxytocolum* (Flügge)). *Acta pathologica et microbiologica scandinavica* **39**, 375-384.
- LECLERC, H. (1962). Étude biochimique d'Enterobacteriaceae pigmentées. *Annales de l'Institut Pasteur, Paris* **102**, 726-741.
- LOCKHART, W. R. & KOENIG, K. (1965). Use of secondary data in numerical taxonomy of the genus *Erwinia*. *Journal of Bacteriology* **90**, 1638-1644.
- PROCTOR, J. R. (1966). Some processes of numerical taxonomy in terms of distance. *Systematic Zoology* **15**, 131-140.
- REPORT of the Subcommittee on Taxonomy of the Enterobacteriaceae (1963). *International Bulletin of Bacteriological Nomenclature and Taxonomy* **13**, 69-93.

- SLOPEK, S. & DURLAKOWA, I. (1967). Studies on the taxonomy of *Klebsiella* bacilli. *Archivum Immunologiae et Therapiae Experimentalis* **15**, 481-487.
- SNEATH, P. H. A. (1957). The application of computers to taxonomy. *Journal of General Microbiology* **17**, 201-226.
- SNEATH, P. H. A. (1966). A comparison of different clustering methods as applied to randomly spaced points. *The Classification Society Bulletin* **1**, 2-18.
- SOKAL, R. R. & MICHENER, C. D. (1958). A statistical method for evaluating systematic relationships. *University of Kansas Science Bulletin* **38**, 1409-1438.
- SOKAL, R. R. & SNEATH, P. H. A. (1963). *Principles of Numerical Taxonomy*. San Francisco & London: W. H. Freeman.
- TAYLOR, E. W., COWAN, S. T., LOVELL, R., TAYLOR, J., THOMAS, S. B., CUTHBERT, W. A., JACOBS, S. E. & CLEGG, L. F. L. (1956). The nomenclature of the coli-aerogenes bacteria. *Journal of Applied Bacteriology* **19**, 108-111.
- 'T MANNETJE, L. (1967). A comparison of eight numerical procedures applied to the classification of some African *Trifolium* taxa based on *Rhizobium* affinities. *Australian Journal of Botany* **15**, 521-528.
- WILSON, G. S. & MILES, A. A. (1964). *Topley and Wilson's Principles of Bacteriology and Immunity*, 5th edn. London: Edward Arnold.