R Plasmids from *Yersinia*

By S. KIMURA, T. IKEDA, T. EDA, YOHKO MITSUI
AND KAZUE NAKATA

Department of Bacteriology, Teikyo University School of Medicine, 2-11-1,
Itabashiku, Tokyo 173, Japan

(Received 25 April 1976; revised 24 June 1976)

INTRODUCTION

The recipient ability of *Yersinia pestis* and *Yersinia pseudotuberculosis* for R plasmids was demonstrated in the primary period of R plasmid research (Ginoza & Matney, 1963). Afterwards, transfer of R plasmids to *Y. enterocolitica* was confirmed by several workers (Knapp & Lebek, 1967; Rusu, Baron & Lazaroae, 1970; Kimura *et al.*, 1975*a*) and transfer to *Y. pseudotuberculosis* was shown by Zarembo (1972). However, there are few reports of detection of R plasmids in naturally occurring *Y. enterocolitica* and *Y. pseudotuberculosis* (Cornelis, Wauters & Bruynoghe, 1973; Kimura *et al.*, 1975*b*; Ikeda, Eda & Kimura, 1975).

Classification of R plasmids by incompatibility was initiated by several workers (Watanabe *et al.*, 1964; Meynell, Meynell & Datta, 1968; Chabbert *et al.*, 1972). The distribution of incompatibility groups was examined, not only in wild-type strains of *Shigella*, *Escherichia* and *Salmonella*, but also in *Proteus rettgeri* (Coetzee, Datta & Hedges, 1972), *Proteus morganii* (Hedges *et al.*, 1973), *Pseudomonas aeruginosa* (Datta *et al.*, 1971; Datta & Hedges, 1972), *Aeromonas liquefaciens* (Aoki *et al.*, 1971), *Providencia* (Hedges, 1974) and *Serratia marcescens* (Hedges, Rodriguez-Lemoine & Datta, 1975). The epidemiology and classification of these plasmids were summarized by Datta (1975*a, b*).

In this paper, we describe patterns of drug resistance, *fi* types and incompatibility grouping of plasmids from six strains of *Y. enterocolitica* and *Y. pseudotuberculosis* obtained in Japan.

METHODS

Yersinia strains. All 42 strains examined were isolated from the intestinal contents of swine, except one *Y. pseudotuberculosis* strain that came from a human patient with gastroenteritis. Of the 37 strains of *Y. enterocolitica*, of three different serotypes (O-3, O-5 and O-9), 29 were drug-sensitive, four were resistant to streptomycin alone, three to streptomycin and tetracycline, and one to ampicillin, streptomycin and tetracycline. All five *Y. pseudotuberculosis* strains were resistant to streptomycin alone.

Escherichia coli K12 strains. These were: J53, F⁻ *lac*⁺ *met* *pro*; J53 *nal*, a nalidixic-acid-resistant mutant of J53; W3104, F⁻ *gal* λ ⁻; W3104 *rif*, a rifampicin-resistant mutant of W3104; W1895, Hfr *met*.

Plasmids. Standard plasmids, one of each of 16 incompatibility groups (Hedges, 1974), were kindly supplied by Drs Yoshikawa and Arai (who had received them from Dr N. Datta). The standard N group plasmid, N3 (Datta & Hedges, 1971), determined resistance to streptomycin (Sm), tetracycline (Tc) and sulphonamides (Su). R plasmids from *Yersinia* strains are shown in Table 1.

Table 1. Nature of *R* plasmids from *Yersinia* strains

Plasmid designation	Strain of origin	Resistance markers*	<i>fi</i> type	Exclusion of N ₃	<i>hspII</i>	Incompatibility group
pTE101	<i>Y. pseudotuberculosis</i> Tateishi	Sm	—	—	—	N
pTE102	<i>Y. pseudotuberculosis</i> TP1004	Sm	—	+	+	N
pTE201	<i>Y. enterocolitica</i> Y-42	Sm	—	+	+	N
pTE211	<i>Y. enterocolitica</i> TP83	Sm,Tc	—	NT	+	Undesignated
pTE211-1†	<i>Y. enterocolitica</i> TP83	Sm	—	+	+	N
pTE212	<i>Y. enterocolitica</i> TP83	Tc	—	—	—	Undesignated
pTE213	<i>Y. enterocolitica</i> TP247	Sm,Tc	—	+	+	N
pTE214	<i>Y. enterocolitica</i> TP248	Sm,Tc	—	+	+	N

* Sm, Streptomycin; Tc, tetracycline.

† Segregant obtained after ultraviolet irradiation.

NT, Not tested.

Phages. P1kc and λ were used for the *hspII* restriction test and *f*₂ for the *fi* determination test.

Media. Liquid medium was Bacto-Penassay broth (Difco). Plating medium was Heart Infusion nutrient agar (Eiken).

Transfer experiments. Donor culture (0.1 ml) and recipient culture (0.9 ml) were added to 4.0 ml Penassay broth. After 3 or 24 h incubation at 37°C, 0.1 ml portions of the mixture were inoculated on agar plates containing streptomycin (25 μ g ml⁻¹) or tetracycline (12.5 μ g ml⁻¹) and nalidixic acid (25 μ g ml⁻¹).

Plasmid incompatibility and entry exclusion. Methods were as described by Coetzee *et al.* (1972).

fi determination. The method of Watanabe, Fukasawa & Takano (1962) was used. Hfr strains of *E. coli* K12 W1895 (NR84) and W1895 (NR45) were used as *fi*⁺ and *fi*⁻ controls respectively.

Restriction and modification of phages. The *hspII* restriction by *R* plasmids was determined as described by Bannister & Glover (1968). Phages P1kc and λ were each propagated on W3104 R⁻ and on W3104 (N₃). *R* plasmid N₃ determines *hspII* restriction and modification specificity (Hedges, 1972). Each phage preparation was titrated on W3104 R⁻ and on the same strain carrying N₃ and each of the *Yersinia* *R* plasmids; efficiencies of plating (e.o.p.) were compared.

RESULTS AND DISCUSSION

Transfer of drug resistance to Escherichia coli K12. The 13 resistant strains were examined for transfer of *R* plasmids to *E. coli* K12. Such plasmids were detected in six strains (Table 1). These *R* plasmids could also re-transfer to *E. coli* K12 W3104 *rif*. One strain (TP83) of *Y. enterocolitica* transferred Tc resistance and Sm,Tc resistance separately to the recipients (Table 1).

fi character of *Yersinia* plasmids. All *R* plasmids derived from *Yersinia* were found to be *fi*⁻.

Compatibility of R plasmids from Yersinia among the standard plasmids. Because the *Yersinia* plasmids were all *fi*⁻ they were not tested for compatibility with F-like plasmids. In the first tests, *Yersinia* strains were used as donors, the recipients being *E. coli* K12 strains carrying standard plasmids of known compatibility groups with distinguishable resistance markers. By this means it was shown that the *Yersinia* *R* plasmids all co-existed stably with plasmids of groups I α , I γ , P, A, C, J, H, L and S, i.e. they belonged to none of these groups. However, the compatibility of the *Yersinia* plasmids with plasmids of groups N, W and M

could not be tested in this way, since we had no standard plasmids of those groups that lacked Sm or Tc resistance. We decided to test for phage restriction and modification (r-m) because it has been reported that all R plasmids with r-m specificity *hspII* fall into compatibility group N (Hedges, 1972) and many of the *fi*⁻ plasmids previously identified in Japan belonged to group N (Datta & Hedges, 1971).

Restriction and modification of R plasmids from Yersinia. Plasmid N₃ and six of the *Yersinia* plasmids, when present in the indicator strain w3104, reduced the e.o.p. of P1kc or λ at least 100-fold. No such restriction was seen when the phage had been propagated on an N₃⁺ strain. No greater than threefold differences in e.o.p. were found between titrations where there was no restriction. This result indicated that these six *Yersinia* plasmids determined *hspII* (Table 1) and suggested that they belonged to incompatibility group N.

Compatibility of R plasmids from Yersinia with N₃ (Su, Sm, Tc). The frequency of transfer of N₃ (per donor bacterium in 24 h mating) was measured using, as recipients, w3104 *rif*^R and w3104 *rif* carrying each of the *Yersinia* plasmids. With five of the plasmids so tested, the frequency of transfer was reduced by at least 200-fold compared with that of the R⁻ control, i.e. entry exclusion of N₃ was demonstrated (Table 1).

The w3104 *rif*^R transconjugants from the above experiments were then used as donors, the recipient being J53 *nal* and selection being for Sm and Tc resistance separately. Only the whole N₃ resistance pattern was observed to be transferred, never the resistance pattern of the *Yersinia* plasmid without N₃. This was taken as evidence that the *Yersinia* plasmids had been eliminated from the w3104 *rif* strains on the introduction of N₃, i.e. that they were incompatible with N₃.

The nature of the R plasmids derived from *Yersinia* is summarized in Table 1.

Strains of *Escherichia* (Oyata *et al.*, 1975) and *Shigella* (Tanaka, Tsunoda & Mitsunashi, 1975) isolated in Japan during the past 4 to 7 years have most frequently been resistant to four drugs (streptomycin, tetracycline, chloramphenicol and sulphonamide) and most plasmids from those strains belonged to incompatibility group FII (Datta, 1975*b*). However, drug-resistance patterns of strains from *Salmonella* were mostly Sm, Tc and plasmids from them were mostly *fi*⁻ (Nakaya, Yoshida & Terawaki, 1975) and belonged to incompatibility group N (Datta, 1975*a*). It seems that plasmids from *Yersinia* isolated in Japan resemble those from *Salmonella*.

We thank Dr Kanazawa, Niigata Railway Hospital, Drs Zen-Yoji and Maruyama, Tokyo Metropolitan Research Laboratory of Public Health, and Drs Sasagawa and Ikemura, Niigata Prefectural Institute of Health, for providing several strains which were used in this investigation.

REFERENCES

- AOKI, T., EGUSA, S., OGATA, Y. & WATANABE, T. (1971). Detection of resistance factors in fish pathogen *Aeromonas liquefaciens*. *Journal of General Microbiology* **65**, 343-349.
- BANNISTER, D. & GLOVER, S. W. (1968). Restriction and modification of bacteriophages by R⁺ strains of *Escherichia coli* K12. *Biochemical and Biophysical Research Communications* **30**, 735-739.
- CHABBERT, Y. A., SCAVIZZI, M. R., WITCHITZ, J. L., GERBAUD, G. R. & BOUANCHAUD, D. H. (1972). Incompatibility groups and the classification of *fi*⁻ resistance factors. *Journal of Bacteriology* **112**, 666-675.
- COETZEE, J. N., DATTA, N. & HEDGES, R. W. (1972). R factors from *Proteus rettgeri*. *Journal of General Microbiology* **72**, 543-552.
- CORNELIS, G., WAUTERS, G. & BRUYNOGHE, G. (1973). Resistances transferables chez des souches sauvages de *Yersinia enterocolitica*. *Annales de Microbiologie (Institut Pasteur)* **124A**, 299-309.
- DATTA, N. (1975*a*). Classification of plasmids into compatibility groups: Distribution of genes among the groups. *Microbial Drug Resistance*, pp. 83-91. Edited by S. Mitsunashi and H. Hashimoto. Tokyo, Japan: University of Tokyo Press.

- DATTA, N. (1975*b*). Epidemiology and classification of plasmids. *Microbiology—1974*, pp. 9–15. Edited by D. Schlessinger. Washington, D.C.: American Society for Microbiology.
- DATTA, N. & HEDGES, R. W. (1971). Compatibility groups among *fi*⁻ R factors. *Nature, London* **234**, 222–223.
- DATTA, N. & HEDGES, R. W. (1972). R factors identified in Paris, some conferring gentamycin resistance, constitute a new compatibility group. *Annales de l'Institut Pasteur* **123**, 779–883.
- DATTA, N., HEDGES, R. W., SHAW, E., SYKES, R. B. & RICHMOND, M. H. (1971). Properties of an R factor from *Pseudomonas aeruginosa*. *Journal of Bacteriology* **108**, 1244–1249.
- GINOZA, H. S. & MATNEY, T. S. (1963). Transmission of a resistance transfer factor from *Escherichia coli* to two species of *Pasteurella*. *Journal of Bacteriology* **85**, 1177–1178.
- HEDGES, R. W. (1972). Phenotypic characterization of *fi*⁻ R factors determining the restriction and modification *hspII* specificity. *Molecular and General Genetics* **115**, 225–233.
- HEDGES, R. W. (1974). R factors from Providence. *Journal of General Microbiology* **81**, 171–181.
- HEDGES, R. W., DATTA, N., COETZEE, J. N. & DENNISON, S. (1973). R factors from *Proteus morganii*. *Journal of General Microbiology* **77**, 249–259.
- HEDGES, R. W., RODRIGUEZ-LEMOINE, V. & DATTA, N. (1975). R factors from *Serratia marcescens*. *Journal of General Microbiology* **86**, 88–92.
- IKEDA, T., EDA, T. & KIMURA, S. (1975). Detection of R plasmids from Sm,Tc resistant strains of *Yersinia enterocolitica*. *Medicine and Biology* **91**, 365–369.
- KIMURA, S., EDA, T., IKEDA, T. & MIZUNO, T. (1975*a*). Transfer of R plasmids to *Yersinia enterocolitica*. *Medicine and Biology* **91**, 35–40.
- KIMURA, S., EDA, T., IKEDA, T. & SUZUKI, M. (1975*b*). Detection of conjugative R plasmids in genus *Yersinia*. *Japanese Journal of Microbiology* **19**, 449–451.
- KNAPP, W. & LEBEK, G. (1967). Übertragung der infektiösen Resistenz auf Pasteurellen. *Pathologie und microbiologie* **30**, 103–121.
- MEYNELL, E., MEYNELL, G. G. & DATTA, N. (1968). Phylogenic relationships of drug-resistance factors and other transmissible bacterial plasmids. *Bacteriological Reviews* **32**, 55–83.
- NAKAYA, R., YOSHIDA, Y. & TERAWAKI, Y. (1975). Antibiotic resistance and R plasmids in *Salmonella* isolated from humans in Japan (1966–1972). *Microbial Drug Resistance*, pp. 237–252. Edited by S. Mitsuhashi and H. Hashimoto. Tokyo, Japan: University of Tokyo Press.
- OTAYA, H., MACHIARA, S., YOSHIMURA, M. & YAMAMOTO, M. (1975). Multiply drug-resistant *Escherichia coli* isolated from clinical specimens. *Microbial Drug Resistance*, pp. 201–208. Edited by S. Mitsuhashi and H. Hashimoto. Tokyo, Japan: University of Tokyo Press.
- RUSU, V., BARON, O. & LAZAROE, D. (1970). Transfert du facteur de resistance (R) des *Enterobacteriaceae* à *Yersinia enterocolitica*. *Archives roumaines de pathologie expérimentale et de microbiologie* **29**, 571–581.
- TANAKA, T., TSUNODA, M. & MITSUHASHI, S. (1975). Drug resistance in *Shigella* strains isolated in Japan from 1965 to 1973. *Microbial Drug Resistance*, pp. 187–199. Edited by S. Mitsuhashi and H. Hashimoto. Tokyo, Japan: University of Tokyo Press.
- WATANABE, T., FUKASAWA, T. & TAKANO, T. (1962). Conversion of male bacteria of *Escherichia coli* K12 to resistance to *f* phages by infection with the episome 'resistance transfer factor'. *Virology* **17**, 218–219.
- WATANABE, T., NISHIDA, H., OGATA, C., ARAI, T. & SATO, S. (1964). Episome-mediated transfer of drug resistance in *Enterobacteriaceae*. VII. Two types of naturally occurring R factors. *Journal of Bacteriology* **88**, 716–726.
- ZAREMBA, M. (1972). *In vitro* and *in vivo* transfer of the R factor to *Yersinia pseudotuberculosis* cells. *Bacterial Plasmids and Antibiotic Resistance*, pp. 37–42. Edited by T. Watanabe. Prague: Avicenum Czechoslovak Medical Press, Prague. Berlin, Heidelberg and New York: (Springer Verlag).